

Topic:- DU\_J19\_MPHIL\_MATHS

**1) Which of the following journals is published by Indian Mathematical Society**

**[Question ID = 13918]**

1. Indian Journal of Pure and Applied Mathematics. [Option ID = 25669]
2. Indian Journal of Mathematics. [Option ID = 25671]
3. Ramanujan Journal of Mathematics. [Option ID = 25670]
4. The Mathematics Students . [Option ID = 25672]

**Correct Answer :-**

- Indian Journal of Pure and Applied Mathematics. [Option ID = 25669]

**2) Name a Fellow of Royal Society who expired in 2019 [Question ID = 13917]**

1. M. S. Ragunathan. [Option ID = 25665]
2. Manjul Bhargava. [Option ID = 25666]
3. Michael Atiyah. [Option ID = 25667]
4. S. R. Srinivasa Varadhan. [Option ID = 25668]

**Correct Answer :-**

- M. S. Ragunathan. [Option ID = 25665]

**3) Which of the following statements is true? [Question ID = 13973]**

1. Every topological space having Bolzano-Weierstrass property is a compact space. [Option ID = 25890]
2. If  $\{x_n\}$  is a convergent sequence in a topological space  $X$  with a limit  $x$  then  $Y = \{x\} \cup \{x_n : n = 1, 2, \dots\}$  is a compact subset of  $X$ . [Option ID = 25891]
3. The projection map  $p : X \times Y \rightarrow Y$  defined by  $p(x, y) = y$  is a closed map for all topological spaces  $X, Y$ . [Option ID = 25889]
4. Every topological space is a first countable space. [Option ID = 25892]

**Correct Answer :-**

- The projection map  $p : X \times Y \rightarrow Y$  defined by  $p(x, y) = y$  is a closed map for all topological spaces  $X, Y$ . [Option ID = 25889]

**4) Which of the following statements is true for topological spaces? [Question ID = 13927]**

1. Every second countable space is separable. [Option ID = 25706]

2. Every separable space is second countable. [Option ID = 25705]
3. Every first countable space is second countable. [Option ID = 25708]
4. Every first countable space is separable. [Option ID = 25707]

**Correct Answer :-**

- Every separable space is second countable. [Option ID = 25705]

**5) Which of the following statements is not true? [Question ID = 13997]**

1. If  $H$  and  $K$  are normal subgroups of  $G$ , then the subgroup generated by  $H \cup K$  is also a normal subgroup of  $G$ .  
[Option ID = 25987]
2. Let  $G$  be a finite group and  $H$  a subgroup of order  $n$ . If  $H$  is the only subgroup of order  $n$ , then  $H$  is normal in  $G$ .  
[Option ID = 25986]
3. The set of all permutations  $\sigma$  of  $S_n$  ( $n \geq 3$ ) such that  $\sigma(n) = n$  is a normal subgroup of  $S_n$ .  
[Option ID = 25985]
4. For groups  $G$  and  $H$  and  $f : G \rightarrow H$  a group homomorphism. If  $H$  is abelian and  $V$  is a subgroup of  $G$  containing  $\ker f$  then  $N$  is a normal subgroup of  $G$ .  
[Option ID = 25988]

**Correct Answer :-**

- The set of all permutations  $\sigma$  of  $S_n$  ( $n \geq 3$ ) such that  $\sigma(n) = n$  is a normal subgroup of  $S_n$ .  
[Option ID = 25985]

**6) Which one of the following fellowship is based on merit in M.A/M.Sc. of the University [Question ID = 13920]**

1. NBHM-JRF. [Option ID = 25679]
2. INSPIRE-JRF [Option ID = 25677]
3. UGC-JRF. [Option ID = 25680]
4. CSIR-JRF [Option ID = 25678]

**Correct Answer :-**

- INSPIRE-JRF [Option ID = 25677]

**7) The Abel prize 2019 was awarded to [Question ID = 13919]**

1. Lennert Carleson. [Option ID = 25673]
2. Mikhail Gromov. [Option ID = 25676]
3. Karen Keskulla Uhlenbeck. [Option ID = 25674]
4. Peter Lax. [Option ID = 25675]

**Correct Answer :-**

- Lennert Carleson. [Option ID = 25673]

8)

Let  $X$  be a normed space over  $\mathbb{C}$  and  $f$  a non-zero linear functional on  $X$ . Then

[Question ID = 13981]

1.  $f$  is surjective and a closed map. [Option ID = 25922]
2.  $f$  is surjective and open. [Option ID = 25921]
3.  $f$  is continuous and bijective. [Option ID = 25924]
4.  $f$  is open and continuous. [Option ID = 25923]

Correct Answer :-

- $f$  is surjective and open. [Option ID = 25921]

9)

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$ .

Then which of the following statements is not true?

[Question ID = 13968]

1.  $f$  is bounded above on  $(a, \infty)$ . [Option ID = 25869]
2.  $f'$  is not continuous at 0. [Option ID = 25871]
3.  $f$  is infinitely differentiable at every non zero  $x \in \mathbb{R}$ . [Option ID = 25870]
4.  $f$  is neither convex nor concave on  $(0, \delta)$ . [Option ID = 25872]

Correct Answer :-

- $f$  is bounded above on  $(a, \infty)$ . [Option ID = 25869]

10)

The principal part of the Laurent series of  $f(z) = \frac{1}{z(z-1)(z-3)}$  in the annulus  $\{z : 0 < |z| < 1\}$  is

[Question ID = 13988]

1.  $-\frac{1}{3z}$  [Option ID = 25951]
2.  $\frac{1}{z}$ . [Option ID = 25949]
3.  $\frac{1}{3z}$  [Option ID = 25952]
4.  $\frac{1}{3z^2}$  [Option ID = 25950]

Correct Answer :-

- $\frac{1}{z}$ . [Option ID = 25949]

11) The general solution of the differential equation

$$\frac{dy}{dx} = \frac{y}{x} + c \cot \frac{y}{x}$$

where  $c$  is a constant, is

[Question ID = 14009]

1.  $\operatorname{cosec}(y/x) = c/x$ . [Option ID = 26036]
2.  $\operatorname{cosec}(y/x) = cx$ . [Option ID = 26035]
3.  $\sec(y/x) = cx$ . [Option ID = 26033]
4.  $\sec(y/x) = c/x$ . [Option ID = 26034]

**Correct Answer :-**

- $\sec(y/x) = cx$ . [Option ID = 26033]

12)

Velocity potential for the uniform stream flow with velocity  $\bar{q} = -Ui$ , where  $U$  is constant and  $i$  is the unit vector in  $x$ -direction, past a stationary sphere of radius  $a$  and centre at origin, for  $r \geq a$  is

[Question ID = 14008]

1.  $U \cos \theta \left( r + \frac{1}{2} \frac{a^2}{r^3} \right)$ . [Option ID = 26029]
2.  $U \cos \theta \left( r^2 + \frac{a^2}{r^3} \right)$ . [Option ID = 26032]
3.  $U \cos \theta \left( r^2 + \frac{1}{2} \frac{a^2}{r^3} \right)$ . [Option ID = 26031]
4.  $U \cos \theta \left( r + \frac{a^2}{r^3} \right)$ . [Option ID = 26030]

**Correct Answer :-**

- $U \cos \theta \left( r + \frac{1}{2} \frac{a^2}{r^3} \right)$ . [Option ID = 26029]

13)

Let  $X = P[a, b]$  be the linear space of all polynomials on  $[a, b]$ . Then which of the following statements is not true?

[Question ID = 13979]

1.  $X$  is dense in  $C[a, b]$  with  $\| \cdot \|_p$ -norm,  $1 \leq p \leq \infty$ . [Option ID = 25916]
2.  $X$  is a Banach space with  $\| \cdot \|_p$ -norm,  $1 \leq p \leq \infty$ . [Option ID = 25913]
3.  $X$  has a denumerable basis. [Option ID = 25915]
4.  $X$  is incomplete with  $\| \cdot \|_\infty$ -norm. [Option ID = 25914]

**Correct Answer :-**

- $X$  is a Banach space with  $\|\cdot\|_p$ - norm,  $1 \leq p \leq \infty$ . [Option ID = 25913]

**14)**

Let  $W = \{(x, x, x) : x \in \mathbb{R}\}$  be a subspace of the inner product space  $\mathbb{R}^3$  over  $\mathbb{R}$ . The orthogonal complement of  $W$  in  $\mathbb{R}^3$  is the plane

**[Question ID = 13995]**

1.  $2x + y + z = 0$ . [Option ID = 25979]
2.  $x + 2y + z = 0$ . [Option ID = 25978]
3.  $x + y + z = 0$ . [Option ID = 25980]
4.  $x + y + 2z = 0$ . [Option ID = 25977]

**Correct Answer :-**

- $x + y + 2z = 0$ . [Option ID = 25977]

**15)**

The integral surface of the partial differential equation  $x^2p + y^2q + z^2 = 0$ ,  $p = \frac{\partial z}{\partial x}$ ,  $q = \frac{\partial z}{\partial y}$  which passes through the hyperbola  $xy = x + y$ ,  $z = 1$  is

**[Question ID = 14007]**

1.  $\frac{1}{x} + \frac{2}{y} + \frac{1}{z} = 3$ . [Option ID = 26027]
2.  $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 3$ . [Option ID = 26028]
3.  $\frac{2}{x} + \frac{1}{y} + \frac{1}{z} = 3$ . [Option ID = 26026]
4.  $\frac{1}{x} + \frac{1}{y} + \frac{2}{z} = 3$ . [Option ID = 26025]

**Correct Answer :-**

- $\frac{1}{x} + \frac{1}{y} + \frac{2}{z} = 3$ . [Option ID = 26025]

**16)**

The value of  $\oint_C x^2 dx + (xy + y^2) dy$ , where  $C$  is the boundary of the region  $R$  bounded by  $y = x$  and  $y = x^2$  and is oriented in positive direction is

**[Question ID = 13969]**

1.  $1/15$  [Option ID = 25876]
2.  $2$  [Option ID = 25875]
3.  $1/10$  [Option ID = 25874]
4.  $1/5$  [Option ID = 25873]

**Correct Answer :-**

- $1/5$  [Option ID = 25873]

**17)**

Let  $W = \{(x, y, 0) : x, y \in \mathbb{R}\}$  be a subspace of  $\mathbb{R}^3$ . The cosets of  $W$  in  $\mathbb{R}^3$  are

**[Question ID = 13994]**

1. lines parallel to z-axis. [Option ID = 25975]
2. lines perpendicular to z-axis. [Option ID = 25976]
3. planes perpendicular to xz- plane. [Option ID = 25973]
4. planes parallel to yz- plane. [Option ID = 25974]

**Correct Answer :-**

- planes perpendicular to xz- plane. [Option ID = 25973]

**18)**

Let  $R$  be a ring with unity. An element  $a$  of  $R$  is called nilpotent if  $a^n = 0$  for some positive integer  $n$ . An element  $a$  of  $R$  is called unipotent if and only if  $1 - a$  is nilpotent. Consider the following statements:

- (I) In a commutative ring with unity, product of two unipotent elements is invertible.
- (II) In a ring with unity, every unipotent element is invertible.

Then

**[Question ID = 14001]**

1. Neither (I) nor (II) is correct. [Option ID = 26004]
2. Both (I) and (II) are correct. [Option ID = 26003]
3. Only (I) is correct. [Option ID = 26001]
4. Only (II) is correct. [Option ID = 26002]

**Correct Answer :-**

- Only (I) is correct. [Option ID = 26001]

**19)** Which of the following statements is not true?

**[Question ID = 13970]**

$$g_n(x) = \frac{1}{n(1+x^2)} \rightarrow 0, n \rightarrow \infty \text{ uniformly on } \mathbb{R}.$$

1. [Option ID = 25877]

2.  $h_n(x) = \frac{\sin nx}{n}$  converges uniformly on  $\mathbb{R}$ . [Option ID = 25879]

3.  $f_n(x) = \frac{x^2 + nx}{x}$  converges uniformly on  $\mathbb{R}$ . [Option ID = 25878]

4.  $u_n(x) = \frac{x^n}{n}$  converges uniformly on  $[0, 1]$ . [Option ID = 25880]

**Correct Answer :-**

$$g_n(x) = \frac{1}{n(1+x^2)} \rightarrow 0, n \rightarrow \infty \text{ uniformly on } \mathbb{R}.$$

[Option ID = 25877]

20)

The value of the integral  $\int_C \frac{dz}{z^2+4}$  where  $C$  is the anticlockwise circle  $|z - i| = 2$  is

[Question ID = 13984]

1.  $2\pi$ . [Option ID = 25935]
2. 0 [Option ID = 25933]
3.  $\pi/2$ . [Option ID = 25934]
4.  $\pi$ . [Option ID = 25936]

**Correct Answer :-**

- 0 [Option ID = 25933]

21)

Which of the following statements is true for the product  $\prod_{\alpha \in \Lambda} X_\alpha$  with product topology of a family  $\{X_\alpha\}_{\alpha \in \Lambda}$  of topological spaces?

[Question ID = 13974]

1. If each  $X_\alpha$  is metrizable then  $\prod_{\alpha \in \Lambda} X_\alpha$  is metrizable. [Option ID = 25895]
2. If each  $X_\alpha$  is normal then  $\prod_{\alpha \in \Lambda} X_\alpha$  is normal. [Option ID = 25893]
3. If each  $X_\alpha$  is completely regular then  $\prod_{\alpha \in \Lambda} X_\alpha$  is completely regular. [Option ID = 25896]
4. If each  $X_\alpha$  is locally connected then  $\prod_{\alpha \in \Lambda} X_\alpha$  is locally connected. [Option ID = 25894]

**Correct Answer :-**

- If each  $X_\alpha$  is normal then  $\prod_{\alpha \in \Lambda} X_\alpha$  is normal. [Option ID = 25893]

22) Consider  $\mathbb{R}$  with usual metric and a continuous map  $f : \mathbb{R} \rightarrow \mathbb{R}$  then

[Question ID = 13975]

1.  $f(A)$  is bounded for every bounded subset  $A$  of  $\mathbb{R}$ . [Option ID = 25899]
2.  $f$  is bounded. [Option ID = 25897]
3.  $f^{-1}(A)$  is compact for all compact subset  $A$  of  $\mathbb{R}$ . [Option ID = 25900]
4. Image of  $f$  is an open subset of  $\mathbb{R}$ . [Option ID = 25898]

**Correct Answer :-**

- $f$  is bounded. [Option ID = 25897]

23) Define a sequence of functions  $\{f_n\}$  on  $\mathbb{R}$  as

$$f_n(x) = \begin{cases} 1, & \text{if } x \in [-n-2, -n) \\ 0, & \text{otherwise.} \end{cases}$$

Let  $\alpha = \int_{-\infty}^{\infty} \lim_{n \rightarrow \infty} f_n(x) dx$  and  $\beta = \lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} f_n(x) dx$ . Then

[Question ID = 13986]

1.  $0 < \alpha < 1, \beta = 1$  [Option ID = 25942]
2.  $\alpha = 0, \beta = \infty$ . [Option ID = 25943]
3.  $\alpha = \beta = 0$ . [Option ID = 25941]
4.  $\alpha = 0, \beta = 2$ . [Option ID = 25944]

**Correct Answer :-**

- $\alpha = \beta = 0$ . [Option ID = 25941]

24)

Suppose  $f$  is an entire function with  $f(0) = 0$  and  $u$  be the real part of  $f$  such that  $|u(x, y)| \leq 1$  for all  $(x, y) \in \mathbb{R}^2$ . Then the range of  $u$  is

[Question ID = 13985]

1.  $[-1, 1]$ . [Option ID = 25938]
2.  $[0, 1]$ . [Option ID = 25937]
3.  $\{0\}$ . [Option ID = 25939]
4.  $[-1, 0]$ . [Option ID = 25940]

**Correct Answer :-**

- $[0, 1]$ . [Option ID = 25937]

25)

For the minimal splitting field  $F$  of a polynomial  $f(x)$  of degree  $n$  over a field  $K$ . Consider the following statements:

- (I)  $F$  over  $K$  is a normal extension.
- (II)  $n \mid [F : K]$ .
- (III)  $F$  over  $K$  is a separable extension.

Then

[Question ID = 14002]

1. All (I), (II) and (III) are true. [Option ID = 26007]
2. None of (I), (II) and (III) is true. [Option ID = 26008]
3. Only (I) is true. [Option ID = 26005]
4. Only (I) and (II) are true. [Option ID = 26006]

**Correct Answer :-**

- Only (I) is true. [Option ID = 26005]

26)



Let  $V = \{x + \alpha y : \alpha, x, y \in \mathbb{Q}\}$ . Then  $V$  a vector space over  $\mathbb{Q}$  of dimension

[Question ID = 13991]

1. 2 [Option ID = 25963]
2. 1 [Option ID = 25964]
3. 3 [Option ID = 25962]
4. infinity. [Option ID = 25961]

**Correct Answer :-**

- infinity. [Option ID = 25961]

27)

Let  $X = \mathbb{C}^2$  with  $\|\cdot\|_1$  norm and  $X_0 = \{(x_1, x_2) \in X : x_2 = 0\}$ . Define  $g : X_0 \rightarrow \mathbb{C}$  by  $g(x) = x_1, x = (x_1, 0)$ . Consider the following statements:

- Every  $f \in X'$  (dual space of  $X$ ) is of the form  $f(x_1, x_2) = ax_1 + bx_2$  for some  $a, b \in \mathbb{C}$ .
- Hahn-Banach extensions of  $g$  are precisely of the form  $f(x) = x_1 + bx_2, x = (x_1, x_2) \in X, |b| \leq 1, b \in \mathbb{C}$ .

Then

[Question ID = 13982]

1. (I) is true but (II) is false. [Option ID = 25925]
2. (I) is false but (II) is true. [Option ID = 25926]
3. Neither (I) nor (II) is true. [Option ID = 25927]
4. Both (I) and (II) are true. [Option ID = 25928]

**Correct Answer :-**

- (I) is true but (II) is false. [Option ID = 25925]

28)

Which of the following statements is not true for a subset  $A$  of a metric space  $X$ , whose closure is  $\bar{A}$ ?

[Question ID = 13978]

1. If  $X$  is totally bounded then  $A$  is totally bounded. [Option ID = 25911]
2.  $A$  is connected if and only if  $\bar{A}$  is connected. [Option ID = 25912]
3.  $A$  is bounded if and only if  $\bar{A}$  is bounded. [Option ID = 25909]
4.  $A$  is totally bounded if and only if  $\bar{A}$  is totally bounded. [Option ID = 25910]

**Correct Answer :-**

- $A$  is bounded if and only if  $\bar{A}$  is bounded. [Option ID = 25909]

29) How many pairs of elements are there that generate

$$D_8 = \langle a, b \mid a^4 = b^2 = 1, ab = ba^{-1} \rangle$$

[Question ID = 13998]

1. 2 [Option ID = 25989]
2. 5 [Option ID = 25991]
3. 8 [Option ID = 25992]
4. 4 [Option ID = 25990]

**Correct Answer :-**

- 2 [Option ID = 25989]

**30)**

For each  $n \in \mathbb{N}$ , define  $x_n \in C[0, 1]$  by

$$x_n(t) = \begin{cases} n^2 t, & 0 \leq t \leq 1/n \\ 1/t, & 1/n < t \leq 1 \end{cases}$$

where  $C[0, 1]$  is endowed with sup-norm. Then which of the following is not true:

**[Question ID = 13983]**

1. The sequence  $\{x_n\}_{n \in \mathbb{N}}$  is uniformly bounded on  $[0, 1]$ . [Option ID = 25931]
2. Each  $x_n$  is uniformly continuous on  $[0, 1]$ . [Option ID = 25932]
3. The set  $\{x_n(t) : n \in \mathbb{N}\}$  is bounded for each  $t \in [0, 1]$ . [Option ID = 25929]
4.  $\|x_n\|_\infty \leq n$  for all  $n$ . [Option ID = 25930]

**Correct Answer :-**

- The set  $\{x_n(t) : n \in \mathbb{N}\}$  is bounded for each  $t \in [0, 1]$ . [Option ID = 25929]

**31)**

The eigenvalues of the boundary value problem  $y'' + y' + (1 + \lambda)y = 0$ ,  $y(0) = 0$ ,  $y(1) = 0$  are

**[Question ID = 14005]**

1.  $-\frac{3}{4} + n^2$ ,  $n \in \mathbb{N}$ . [Option ID = 26018]
2.  $\frac{3}{4} + n^2\pi^2$ ,  $n \in \mathbb{N}$ . [Option ID = 26019]
3.  $-\frac{3}{4} + n^2\pi^2$ ,  $n \in \mathbb{N}$ . [Option ID = 26020]
4.  $\frac{3}{4} + n^2$ ,  $n \in \mathbb{N}$ . [Option ID = 26017]

**Correct Answer :-**

- $\frac{3}{4} + n^2$ ,  $n \in \mathbb{N}$ . [Option ID = 26017]

**32)**

Let  $(X, d)$  be a complete metric space. Then which of the following statements holds true?

**[Question ID = 13976]**

1.  $X$  is compact as well as connected. [Option ID = 25902]

2.

If  $\{F_n\}$  is a decreasing sequence of non-empty closed subsets of  $X$  then  $F = \bigcap_{n=1}^{\infty} F_n$  is non-empty.

[Option ID = 25903]

3. Every open subspace of  $X$  is complete. [Option ID = 25904]

4.

If  $X$  is union of a sequence of its subsets then the closure of at least one set in the sequence must have non-empty interior.

[Option ID = 25901]

**Correct Answer :-**

•

If  $X$  is union of a sequence of its subsets then the closure of at least one set in the sequence must have non-empty interior.

[Option ID = 25901]

**33)**

Let  $V$  be the set of all polynomials over  $\mathbb{R}$ . A linear transformation  $D : V \rightarrow V$  is defined by  $D(f(x)) = \frac{d^3}{dx^3}(f(x))$ . Then

**[Question ID = 13993]**

1. dimension of kernel of  $D$  is 2. [Option ID = 25969]
2. dimension of kernel of  $D$  is 4. [Option ID = 25970]
3. range of  $D = V$ . [Option ID = 25972]
4. range of  $D$  is a finite dimensional space [Option ID = 25971]

**Correct Answer :-**

- dimension of kernel of  $D$  is 2. [Option ID = 25969]

**34)** If  $G = \mathbb{Z}_6 \oplus \mathbb{Z}_{20} \oplus \mathbb{Z}_{72}$ , then  $G$  is isomorphic to

**[Question ID = 14000]**

1.  $\mathbb{Z}_8 \oplus \mathbb{Z}_9 \oplus \mathbb{Z}_{40}$ . [Option ID = 25998]
2.  $\mathbb{Z}_2 \oplus \mathbb{Z}_{12} \oplus \mathbb{Z}_{360}$ . [Option ID = 26000]
3.  $\mathbb{Z}_5 \oplus \mathbb{Z}_{27} \oplus \mathbb{Z}_{64}$ . [Option ID = 25997]
4.  $\mathbb{Z}_6 \oplus \mathbb{Z}_{32} \oplus \mathbb{Z}_{45}$ . [Option ID = 25999]

**Correct Answer :-**

- $\mathbb{Z}_5 \oplus \mathbb{Z}_{27} \oplus \mathbb{Z}_{64}$ . [Option ID = 25997]

**35)**

The general solution of the partial differential equation

$$\frac{\partial^2 z}{\partial x \partial y} + \frac{\partial z}{\partial x} - \frac{\partial z}{\partial y} - z = xy$$

is

[Question ID = 14006]

1.  $e^x f_1(y) + e^{-y} f_2(x) + xy + y - x - 1.$  [Option ID = 26023]
2.  $e^x f_1(y) + e^{-y} f_2(x) - xy - y + x + 1.$  [Option ID = 26022]
3.  $e^{-x} f_1(y) + e^y f_2(x) + xy + y - x - 1.$  [Option ID = 26024]
4.  $e^{-x} f_1(y) + e^y f_2(x) - xy - y + x + 1.$  [Option ID = 26021]

**Correct Answer :-**

- $e^{-x} f_1(y) + e^y f_2(x) - xy - y + x + 1.$  [Option ID = 26021]

**36)**

The function  $f : [0, 2\pi] \rightarrow S^1$  defined by  $f(t) = e^{it}$ , where  $S^1$  is the unit circle, is

[Question ID = 13972]

1. continuous, one-one but not onto. [Option ID = 25886]
2. not a continuous map. [Option ID = 25885]
3. a continuous bijection but not an open map. [Option ID = 25887]
4. a homeomorphism. [Option ID = 25888]

**Correct Answer :-**

- not a continuous map. [Option ID = 25885]

**37)** Define  $f$  on  $\mathbb{C}$  by

$$f(z) = \begin{cases} \frac{(\bar{z})^2}{z}, & \text{if } z \neq 0 \\ 0, & z = 0. \end{cases}$$

Let  $u$  and  $v$  denote the real and imaginary parts of  $f$ . Then at the origin

[Question ID = 13990]

1.  $u, v$  do not satisfy the Cauchy Riemann equations but  $f$  is differentiable. [Option ID = 25959]
2.  $u, v$  satisfy the Cauchy Riemann equations but  $f$  is not differentiable [Option ID = 25958]
3.  $f$  is differentiable and  $u, v$  satisfy the Cauchy Riemann equations. [Option ID = 25957]
4.  $f$  is not differentiable and  $u, v$  do not satisfy the Cauchy Riemann equations. [Option ID = 25960]

**Correct Answer :-**

- $f$  is differentiable and  $u, v$  satisfy the Cauchy Riemann equations. [Option ID = 25957]

38)

Let  $V$  be the set of all polynomials over  $\mathbb{R}$ . Define  $W = \{x^n f(x) : f(x) \in V\}$ ,  $n \in \mathbb{N}$  is fixed. Then which of the following statements is not true?

[Question ID = 13992]

1.  $V$  is infinite dimensional over  $\mathbb{R}$ . [Option ID = 25967]
2. The quotient space  $V/W$  is finite dimensional. [Option ID = 25966]
3.  $W$  is not a subspace of  $V$ . [Option ID = 25965]
4.  $V$  has linearly independent set of  $m$  vectors for every  $m \in \mathbb{N}$ . [Option ID = 25968]

Correct Answer :-

- $W$  is not a subspace of  $V$ . [Option ID = 25965]

39)

Navier Stokes equation of motion for steady viscous incompressible fluid flow in absence of body force is (where  $\bar{q}$ ,  $p$ ,  $\rho$ ,  $\bar{\zeta}$  and  $\nu$  are velocity, pressure, density, vorticity, and kinematic coefficient of viscosity respectively)

[Question ID = 14004]

1.  $\nabla\left(\frac{1}{2}\bar{q}^2 - \frac{p}{\rho}\right) + \bar{q} \times \bar{\zeta} = \nu \nabla^2 \bar{q}$ . [Option ID = 26015]
2.  $\nabla\left(\frac{1}{2}\bar{q}^2 + \frac{p}{\rho}\right) - \bar{q} \times \bar{\zeta} = \nu \nabla^2 \bar{q}$ . [Option ID = 26014]
3.  $\nabla\left(\frac{1}{2}\bar{q}^2 + \frac{p}{\rho}\right) + \bar{q} \times \bar{\zeta} = \nu \nabla^2 \bar{q}$ . [Option ID = 26013]
4.  $\nabla\left(\bar{q}^2 + \frac{p}{\rho}\right) - \bar{q} \times \bar{\zeta} = -\nu \nabla^2 \bar{q}$ . [Option ID = 26016]

Correct Answer :-

- $\nabla\left(\frac{1}{2}\bar{q}^2 + \frac{p}{\rho}\right) + \bar{q} \times \bar{\zeta} = \nu \nabla^2 \bar{q}$ . [Option ID = 26013]

40)

Let  $X = C_{00}$  (the space of all real sequences having only finitely many non-zero terms) with  $\|\cdot\|_{\infty}$ -norm. Define  $P : X \rightarrow X$  by

$$P(x)(2j - 1) = x(2j - 1) + jx(2j)$$

$$P(x)(2j) = 0$$

for  $x \in X$ ,  $j \in \mathbb{N}$ . Then which of the following statements is not true?

[Question ID = 13980]

1.  $P$  is closed map. [Option ID = 25918]
2.  $P$  is linear and  $P^2 = P$ . [Option ID = 25917]
3.  $\text{Range}(P)$  is a closed subspace of  $X$ . [Option ID = 25919]
4.  $P$  is a continuous map. [Option ID = 25920]

**Correct Answer :-**

- $P$  is linear and  $P^2 = P$ . [Option ID = 25917]

**41)**

The value of  $\int_C 2x \, ds$ , where  $C$  consists of the arc  $C_1$  of the parabola  $y = x^2$  from  $(0, 0)$  to  $(1, 1)$  followed by the line segment from  $(1, 1)$  to  $(0, 0)$  is

**[Question ID = 13971]**

1.  $\frac{5\sqrt{5} - 1}{6} + 2\sqrt{2}$ . [Option ID = 25882]
2.  $\frac{5\sqrt{5} - 4}{3} + 2\sqrt{2}$ . [Option ID = 25884]
3.  $\frac{5\sqrt{5} - 1}{6} + \sqrt{2}$ . [Option ID = 25881]
4.  $\frac{3\sqrt{5} - 1}{5} + \sqrt{2}$ . [Option ID = 25883]

**Correct Answer :-**

- $\frac{5\sqrt{5} - 1}{6} + \sqrt{2}$ . [Option ID = 25881]

**42)**

For each integer  $n$ , define  $f_n(x) = x + n$ ,  $x \in \mathbb{R}$  and let  $G = \{f_n : n \in \mathbb{Z}\}$ . Then

**[Question ID = 13999]**

1.  $G$  is a cyclic group under composition. [Option ID = 25994]
2.  $G$  is a non-cyclic group under composition. [Option ID = 25995]
3.  $G$  does not form a group under composition. [Option ID = 25993]
4.  $G$  is a non-abelian group under composition. [Option ID = 25996]

**Correct Answer :-**

- $G$  does not form a group under composition. [Option ID = 25993]

**43)**

Suppose  $G$  is an open connected subset of  $\mathbb{C}$  containing 0 and  $f : G \rightarrow \mathbb{C}$  is analytic such that  $f(0) = 0$  and  $|f(z) - 1| = 1$  for all  $z \in G$ . Then the range of  $f$  is

[Question ID = 13989]

1.  $\{0, 2\}$ . [Option ID = 25954]
2.  $\{1 + e^{i\theta} : 0 \leq \theta \leq \pi\}$  [Option ID = 25956]
3.  $\{1 + e^{i\theta} : 0 \leq \theta \leq 2\pi\}$ . [Option ID = 25953]
4.  $\{0\}$  [Option ID = 25955]

**Correct Answer :-**

- $\{1 + e^{i\theta} : 0 \leq \theta \leq 2\pi\}$ . [Option ID = 25953]

- 44) Consider the following statements:  
Dimension of kinematic coefficient of viscosity is  
(I)  $L^2T^{-1}$ .  
(II) same as dimension of stream function.  
(III)  $L^{-2}T^1$ .  
(IV) same as dimension of stokes stream function.  
Then

[Question ID = 14003]

1. Only (III) and (IV) are true. [Option ID = 26012]
2. Only (I) and (II) are true. [Option ID = 26009]
3. Only (II) and (III) are true. [Option ID = 26011]
4. Only (I) and (IV) are true. [Option ID = 26010]

**Correct Answer :-**

- Only (I) and (II) are true. [Option ID = 26009]

45)

Consider a sequence  $\{x_n\}$  defined by  $0 < x_1 < 1$  and  $x_{n+1} = 1 - \sqrt{1 - x_n}$ ,  $n = 1, 2, \dots$ . Then  $\frac{x_{n+1}}{x_n}$  converges to

[Question ID = 13967]

1. 0 [Option ID = 25866]
2. 1/3 [Option ID = 25867]
3. 1/2 [Option ID = 25868]
4. 1 [Option ID = 25865]

**Correct Answer :-**

- 1 [Option ID = 25865]

46)

Which of the following statements about the outer measure  $m^*$  on  $\mathbb{R}$  is true?

[Question ID = 13987]

1. There exists an open subset  $A \subseteq \mathbb{R}$  such that  $m^*A = 0$ . [Option ID = 25945]
2. Every subset of  $\mathbb{R}$  of zero outer measure is at most countable. [Option ID = 25947]
3. If  $B \subseteq \mathbb{R}$  is unbounded, then  $m^*B > 0$ . [Option ID = 25948]
4. Every non empty closed subset  $E$  of  $\mathbb{R}$  has  $m^*E > 0$ . [Option ID = 25946]

**Correct Answer :-**

- There exists an open subset  $A \subseteq \mathbb{R}$  such that  $m^*A = 0$ . [Option ID = 25945]

**47) Which of the following statements is true? [Question ID = 13977]**

1. In a metric space, the image of a Cauchy sequence under a continuous map is a Cauchy sequence . [Option ID = 25906]
2. Every closed and bounded subset of a metric space is compact. [Option ID = 25907]
3. Every infinite subset of the closed unit ball  $B$  in  $\mathbb{R}^n$  has a limit point in  $B$ . [Option ID = 25905]
4. In a metric space, every closed ball of positive radius is connected. [Option ID = 25908]

**Correct Answer :-**

- Every infinite subset of the closed unit ball  $B$  in  $\mathbb{R}^n$  has a limit point in  $B$ . [Option ID = 25905]

**48) Which one of the following statements is not true? [Question ID = 13966]**

1. There is a function  $f$  defined on  $\mathbb{R}$  which is continuous on  $\mathbb{Q}$  (rational numbers) and discontinuous on  $\mathbb{Q}'$  (irrational numbers). [Option ID = 25861]
2. Monotone convergence property is equivalent to completeness of  $\mathbb{R}$ . [Option ID = 25864]
3. Bolzano-Weierstrass theorem is equivalent to completeness of  $\mathbb{R}$ . [Option ID = 25863]
4. Cantor's intersection property of  $\mathbb{R}$  is equivalent to completeness of  $\mathbb{R}$ . [Option ID = 25862]

**Correct Answer :-**

- There is a function  $f$  defined on  $\mathbb{R}$  which is continuous on  $\mathbb{Q}$  (rational numbers) and discontinuous on  $\mathbb{Q}'$  (irrational numbers). [Option ID = 25861]

**49) Present President of the Ramanujan Mathematical Society is [Question ID = 13916]**

1. V. Kumar Murty. [Option ID = 25664]



2. Dinesh Singh [Option ID = 25661]
3. S. Ponnusamy [Option ID = 25662]
4. R. Balakrishnan. [Option ID = 25663]

**Correct Answer :-**

- Dinesh Singh [Option ID = 25661]

**50) The characteristic and the minimal polynomial are same for the matrix**

**[Question ID = 13996]**

1.  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  [Option ID = 25983]
2.  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  [Option ID = 25982]
3.  $\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$  [Option ID = 25981]
4. All of the above matrices [Option ID = 25984]

**Correct Answer :-**

- $\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$  [Option ID = 25981]