SAT Chemistry Practice Test 20

SAT Chemistry Practice Test 3: Part B

- 1. Treatment of the dry solid with a mild oxidizing agent produces a purple solid
- A. N₂
- B. KI
- $C.\;CCI_4$
- D. AgNO₃
- E. CaCO₃
- 2. Is the principle reaction responsible for the energy output of the sun
- A. Gamma decay
- B. Nuclear fusion
- C. Alpha decay
- D. Positron emission
- E. Nuclear fission
- 3. Is a nuclear process that results in no change in the mass number and atomic number of a nuclide
- A. Gamma decay
- B. Nuclear fusion
- C. Alpha decay
- D. Positron emission
- E. Nuclear fission
- 4. Responsible for most helium found on Earth
- A. Gamma decay
- B. Nuclear fusion
- C. Alpha decay
- D. Positron emission
- E. Nuclear fission
- 5. The nuclear process that transmutes uranium-238 into thorium-234

- A. Gamma decay
- B. Nuclear fusion
- C. Alpha decay
- D. Positron emission
- E. Nuclear fission
- 6. Has a pH of 13
- A. 0.1 *M* MgCl₂
- B. 0.1 *M* HCIO₄
- C. 0.1 *M* NH₄OH
- D. 0.1 *M* KOH
- E. 0.1 *M* LiNO₃
- 7. The solution with the lowest freezing point temperature
- A. 0.1 *M* MgCl₂
- B. 0.1 M HCIO₄
- C. 0.1 *M* NH₄OH
- D. 0.1 *M* KOH
- E. 0.1 *M* LiNO₃
- 8. The solution with the highest boiling point temperature
- A. 0.1 *M* MgCl₂
- B. 0.1 *M* HCIO₄
- C. 0.1 *M* NH₄OH
- D. 0.1 *M* KOH
- E. 0.1 *M* LiNO₃
- 9. Indicates a red flame when ionized with a Bunsen burner
- A. 0.1 *M* MgCl₂
- B. 0.1 M HCIO₄
- C. 0.1 *M* NH₄OH

D. 0.1 *M* KOH

E. 0.1 *M* LiNO₃

10. Choose the answer below that accurately describes the correct molecular shape for the molecule $XeOF_4$.

- A. Tetrahedral
- B. Trigonal pyramidal
- C. Trigonal bipyramidal
- D. Square pyramidal
- E. Flat
- **11.** For the radioactive atom ⁹⁹Tc, what is the correct number of protons and neutrons?
- A. 43 protons and 56 neutrons
- B. 43 protons and 99 neutrons
- C. 56 protons and 43 neutrons
- D. 56 protons and 99 neutrons
- E. Cannot be determined
- 12. Which one of the following acids is NOT strong?
- A. HCI
- B. HBr
- C. HNO₃
- D. H₃PO₄
- E. H₂SO₄

13. Identify the equation used to determine the amount of heat required to melt 10 grams of ice.

A. $Q = mC_{sp}\Delta T$ B. $Q = n\Delta H$ C. $KE = \frac{1}{2}mv^2$ D. PE = mghE. PV = nRT 14. Identify the correct ground state electron configuration for Cr.

- A. [Ar] 3s²3d⁴
- B. [Ar] 3s²3d⁵
- C. [Ar] 4s²3d⁵
- D. [Ar] 4s²3d⁴
- E. [Ar] 4s¹3d⁵
- 15. What is the hydroxide concentration for a solution with a pH of 10 at 25°C?
- A. 10⁻¹⁴ M
- B. 10⁻¹⁰ M
- C. 10⁻⁷ M
- D. 10⁻⁴ M
- E. 10⁻¹ M

16. Five hundred milliliters of solution of 0.1 *M* NaBr has how many milligrams of bromine?

- A. 200 mg
- B. 400 mg
- C. 2,000 mg
- D. 4,000 mg
- E. 20,000 mg

17. According to the ideal gas law, what is the approximate volume that will be occupied by 0.5 mole of an ideal gas at 30° C and 3 atm pressure (gas constant R = 0.0821 L?atm/mol?K)?

A. Less than 1 L

B. 5 L

- C. 10 L
- D. 15 L
- E. More than 20 L

18. Given that $\Delta G = \Delta H - T\Delta S$, how is the spontaneity of an endothermic reaction expected to change with decreasing *T*?

A. Becomes less spontaneous

- B. Becomes more spontaneous
- C. Does not change
- D. Decreases at first but then increases
- E. Insufficient information to make a conclusion
- 19. Identify the element with the greatest first ionization energy.
- A. Ce
- B. C
- C. Cl
- D. Ca
- E. Cs

20. Identify the molecule/ion with the greatest potential to act as a Lewis acid.

- A. CH_3^+
- B. CN⁻
- C. NH₃
- D. BF₄⁻
- E. CO₂
- **21.** 2 Ca₃(PO₄)₂ + 6 SiO₂ + 10 C \rightarrow P₄ +...CaSiO₃ + 10 CO

Which coefficient balances the reaction given above?

- A. 2
- B. 4
- C. 5
- D. 6
- E. 8

22. A 100-milliliter solution containing $AgNO_3$ was treated with excess NaCl to completely precipitate the silver as AgCl. If 5.7 g AgCl was obtained, what was the concentration of Ag⁺ in the original solution?

A. 0.03 *M*

B. 0.05 *M*

C. 0.12 M

D. 0.30 M

E. 0.40 M

23. Identify which of the following statements is FALSE.

- A. The vapor pressure of a liquid decreases with increasing atmospheric pressure.
- B. The value of an equilibrium constant is dependent on temperature.
- C. The rate of a spontaneous reaction cannot be determined solely by its Gibbs free energy.
- D. During a phase transition, the temperature of a substance must be constant.
- E. The addition of a catalyst to a reaction at equilibrium has no net effect on the system.