

# MTB 2019

---

## Notation

$\mathbb{Z}$  = the set of integers

$\mathbb{N} = \{n \in \mathbb{Z} : n \geq 1\}$

$\mathbb{R}$  = the set of real numbers

$\mathbb{Q}$  = the set of rational numbers

$\mathbb{C}$  = the set of complex numbers

---

- (1) Let  $A$  be a diagonalisable  $m \times m$  matrix with entries from  $\mathbb{C}$ . Define  $\exp(A) = \sum_{n=0}^{\infty} \frac{A^n}{n!}$  assuming  $A^0$  is the identity matrix. Prove that  $\det(\exp(A)) = e^{\text{Tr}(A)}$ .
- (2) Let  $H = \{1 + 4k : k \in \mathbb{Z}, k \geq 0\}$ . An element  $x \in H$  is called *H-prime* if  $x \neq 1$  and  $x$  cannot be written as product of two strictly smaller elements of  $H$ .
- (i) Show that  $xy \in H$  for all  $x, y \in H$ .
- (ii) Prove that every  $x \in H$  greater than 1 can be factored as a product of *H*-primes but unique factorisation does not hold.
- (3) Find an ideal  $I$  in  $A = \frac{\mathbb{Z}[X]}{(X^4 + X^2 + 1)}$  such that  $\frac{A}{I}$  is a finite field with 25 elements.
- (4) (a) Suppose  $A$  and  $B$  are closed subsets of a topological space such that  $A \cap B$  and  $A \cup B$  are connected. Prove that  $A$  and  $B$  are connected.
- (b) Demonstrate that the conclusion may not hold if the assumption of  $A$  and  $B$  being closed subsets, is dropped.
- (5) Examine whether there is a polynomial  $f(X) \in \mathbb{R}[X]$  such that  $\frac{\mathbb{R}[X]}{(f(X))}$  is isomorphic **as rings** to the product ring  $\mathbb{C} \times \mathbb{C}$ .

- (6) Let  $f(x), g(x) \in \mathbb{Z}[X]$  with  $f(x) = \sum_{j=0}^n a_j x^j$  and  $b(x) = \sum_{j=0}^n b_j x^j$ . For  $m \in \mathbb{N}$ , we say  $f \equiv g \pmod{m}$  if  $a_j = b_j \pmod{m}$  for  $0 \leq j \leq n$ . For an odd prime  $p > 0$ , let

$$f(x) = x^{p-1} - 1 \quad \text{and} \quad g(x) = (x-1)(x-2)\cdots(x-p+1).$$

Prove that

- (i) the polynomial  $f(x) - g(x)$  has degree  $p-2$ , and
- (ii)  $f \equiv g \pmod{p}$ .

- (7) Prove that the following two groups are isomorphic:

- (a)  $\mathbb{Z}[X]$ , the group of polynomials with integer coefficients under addition, and
- (b)  $\mathbb{Q}_{>0}$ , the group of positive rational numbers under multiplication.

Hint: Fundamental theorem of Arithmetic.

- (8) Prove that there cannot be any topological space  $X$  such that  $\mathbb{R}$  is homeomorphic to  $X \times X$  with the product topology.

- (9) Let  $A = \frac{\mathbb{C}[X, Y]}{(X^2 + Y^2 - 1)}$ , and  $x, y$  denote the images of  $X, Y$  in  $A$  respectively. If  $u = x + iy$ , then prove that

- (a)  $u$  is a unit in  $A$ , and
- (b)  $u - i$  generates a maximal ideal of  $A$ .

- (10) Let  $A$  be a real symmetric  $m \times m$  matrix with  $m$  distinct eigenvalues and  $v_1, \dots, v_m$  be the corresponding eigenvectors. Let  $C$  be an  $m \times m$  matrix satisfying  $\langle Cv_j, v_j \rangle = 0$  for  $1 \leq j \leq m$ . Prove that there exists an  $m \times m$  matrix  $X$  such that  $AX - XA = C$ .