

## INSTRUCTIONS FOR CANDIDATES

- Please answer FOUR questions from EACH group.
- Each question carries 10 marks. Total marks : 80.
- $\mathbb{R}, \mathbb{C}, \mathbb{Z}$  and  $\mathbb{N}$  denote respectively the set of real numbers, set of complex numbers, set of all integers and set of all positive integers.

### Group A

1. Let  $a_n \in \mathbb{R}$ , such that  $\sum_{n=1}^{\infty} |a_n| = \infty$  and  $\sum_{n=1}^m a_n \rightarrow a \in \mathbb{R}$  as  $m \rightarrow \infty$ .  
Let  $a_n^+ = \max\{a_n, 0\}$ . Show that  $\sum_{n=1}^{\infty} a_n^+ = \infty$ .
2. Let  $E = \{(x, y, z) \in \mathbb{R}^3 \mid x, y, z > 0, xy + yz + zx = 1\}$ . Prove that there exists  $(a, b, c) \in E$  such that  $abc \geq xyz$ , for all  $(x, y, z) \in E$ .
3. Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be an increasing function. Suppose there are sequences  $(x_n)$  and  $(y_n)$  such that  $x_n < 0 < y_n$  for all  $n \geq 1$  and  $f(y_n) - f(x_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Prove that  $f$  is continuous at 0.
4. Do there exist continuous functions  $P$  and  $Q$  on  $[0, 1]$  such that  $y(t) = \sin(t^2)$  is a solution to  $y'' + Py' + Qy = 0$  on  $[\frac{1}{n}, 1]$  for all  $n \geq 1$ ? Justify your answer.
5. Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \int_{e^{x^3+x}}^{1+e^{x^3+x}} e^{r^2} dr$$

for all  $x \in \mathbb{R}$ . Prove that  $f$  is monotone.

6. Let  $w = \{w(i, j)\}_{1 \leq i, j \leq m}$  be an  $m \times m$  symmetric matrix with non-negative real entries such that  $w(i, j) = 0$  if and only if  $i = j$ . Show that  $d(i, j) = \min\{\sum_{j=0}^{k-1} w(i_j, i_{j+1}) \mid k \geq 1, i_0 = i, i_k = j, i_j \in \{1, \dots, m\}\}$  is a metric on  $\{1, \dots, m\}$ .

## Group B

7. Factory A produces 1 bad watch in 100 and factory B produces 1 bad watch in 200. You are given two watches from one of the factories and you don't know which one.
  - (a) What is the probability that the second watch works?
  - (b) Given that the first watch works, what is the probability that the second watch works?
8. Let  $R$  be a commutative ring containing a field  $k$  as a sub-ring. Assume that  $R$  is a finite dimensional  $k$ -vector space. Show that every prime ideal of  $R$  is maximal.
9. Let  $p, q$  be prime numbers and  $n \in \mathbb{N}$  such that  $p \nmid n - 1$ . If  $p \mid n^q - 1$  then show that  $q \mid p - 1$ .
10. Determine all finite groups which have exactly 3 conjugacy classes.
11. Let  $F$  be a field,  $a \in F$ ,  $p$  a prime integer. Suppose the polynomial  $x^p - a$  is reducible in  $F[x]$ . Prove that this polynomial has a root in  $F$ .
12. Let  $V$  be a finite-dimensional vector space over a field  $F$  and let  $T : V \rightarrow V$  be a linear transformation. Let  $W \subseteq V$  be a subspace such that  $T(W) \subseteq W$ . Suppose  $T$  is diagonalizable. Is  $T$  restricted to  $W$  also diagonalizable?