

PMB 2019

1. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function satisfying

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow -\infty} f(x) = 0.$$

Show that f is a bounded function on \mathbb{R} and attains a maximum or a minimum. Give an example to show that it attains a maximum but not a minimum.

2. Let $g : [0, 1] \rightarrow \mathbb{R}$ be a continuous function such that $g(1) = 0$. Show that

$$\sup_{x \in [0, 1]} |x^n g(x)| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

3. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a twice continuously differentiable function and suppose $f(0) = f'(0) = 0$. If $|f''(x)| \leq 1$ for all $x \in \mathbb{R}$, then prove that $|f(x)| \leq 1/2$ for all $x \in [-1, 1]$.

4. Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a function defined by

$$f(x, y) = \begin{cases} \frac{x^2 y^3}{x^4 + y^6}, & \text{if } x \neq 0, y \in \mathbb{R}, \\ 0, & \text{if } x = 0, y \in \mathbb{R}. \end{cases}$$

- (a) Find all $(a, b) \in \mathbb{R}^2 \setminus \{(0, 0)\}$ such that f has a nonzero directional derivative at $(0, 0)$ with respect to the direction (a, b) .
- (b) Is f continuous at $(0, 0)$? Justify your answer.
5. Let C be a subset of a compact metric space (X, d) . Assume that for every continuous function $h : X \rightarrow \mathbb{R}$, the restriction of h to C attains a maximum on C . Prove that C is compact.

Please turn over

6. Let G be a non-abelian group of order pq , where $p < q$ are primes.
- How many elements of G have order q ?
 - How many elements of G have order p ?
7. Prove or disprove the following statement: The ring $\mathbb{Q}[X]/(X^4 - 1)$ is isomorphic to a product of fields.
8. Let M be a symmetric matrix with real entries such that $M^k = 0$ for some $k \in \mathbb{N}$. Show that $M = 0$.
9. Suppose A and B are two $n \times n$ matrices with real entries such that the sum of their ranks is strictly less than n . Show that there exists a nonzero column vector $\mathbf{x} \in \mathbb{R}^n$ such that $A\mathbf{x} = B\mathbf{x} = \mathbf{0}$.
10. Suppose there are n persons in a party. Every pair of persons meet each other with probability $p \in (0, 1)$ independently of the other pairs. Let $N(i)$ be the number of people the i^{th} person meets in the party. For all $i, j \in \{1, 2, \dots, n\}$ with $i \neq j$ and for all $k, l \in \{1, 2, \dots, n-2\}$, show that

$$P[N(i) = k, N(j) = l] = \binom{n-2}{k-1} \binom{n-2}{l-1} p^{k+l-1} (1-p)^{2n-k-l-2} \\ + \binom{n-2}{k} \binom{n-2}{l} p^{k+l} (1-p)^{2n-k-l-3}.$$