

ANSWER KEY (AIPMT-2009)

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Ans	2	4	2	3	2	3	3	2	3	1	4	4	2	2	1	3	4	1	4	1
Ques.	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
Ans	3	4	1	3	4	2	4	2	2	2	2	1	4	2	2	3	3	2	4	3
Ques.	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
Ans	3	1	1	3	2	1	3	1	4	1	4	2	2	4	4	2	2	2	2	2
Ques.	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
Ans	3	2	2	2	3	1	4	4	1	2	3	1	3	3	2	2	2	4	4	3
Ques.	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
Ans	4	2	2	1	1	2	1	1	2	4	2	2	3	2	1	1	4	2	4	1
Ques.	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
Ans	4	3	4	2	2	3	4	4	3	1	2	1	4	1	4	1	4	4	2	2
Ques.	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140
Ans	3	2	3	2	4	3	1	1	2	1	3	4	3	4	1	1	4	3	4	2
Ques.	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160
Ans	1	1	1	1	4	4	1	4	1	1	3	2	3	3	3	3	2	3	3	2
Ques.	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180
Ans	1	1	3	4	4	2	3	3	2	3	4	2	3	3	2	3	1	2	3	2
Ques.	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200
Ans	2	1	3	4	4	1	3	3	2	4	4	3	3	2	3	2	4	3	1	4

HINTS & SOLUTIONS

$$1. [P] = \frac{F}{A} = \left[\frac{MLT^{-2}}{L^2} \right] = [ML^{-1}T^{-2}]$$

$$2. s = \frac{1}{2} at^2$$

$$\frac{s_2}{s_1} = \left(\frac{20}{10} \right)^2 = 4$$

$$s_2 = 4s_1$$

$$3. S_r = v_r t$$

$$1000 = (v - 10) \times 100$$

$$v = 20 \text{ m/s}$$

$$4. \Sigma F = ma$$

$$\Rightarrow T - mg = ma$$

$$a = \frac{T - mg}{m} = 4 \text{ m/s}^2$$

$$5. \vec{P}_1 + \vec{P}_2 + \vec{P}_3 = 0$$

$$|\vec{P}_3| = |\vec{P}_1 + \vec{P}_2|$$

$$m \times 4 = \sqrt{P_1^2 + P_2^2} = \sqrt{12^2 + 16^2}$$

$$m = 5 \text{ kg}$$

6. Loss in grav. PE = gain in spring PE
At maximum elongation

$$Mgx = \frac{1}{2} kx^2 \Rightarrow x = \frac{2Mg}{k}$$

$$7. \vec{R}_{cm} = \frac{m_1 \vec{r}_1 + m_2 \vec{r}_2}{m_1 + m_2} = -2\hat{i} - \hat{j} + \hat{k}$$

$$8. I = 4 \times \left[\frac{M\ell^2}{12} + M \left(\frac{\ell}{2} \right)^2 \right] \text{ the parallel axis theorem} = \frac{4}{3} M\ell^2$$

9. Apply conservation of angular momentum.

$$L_i = L_r$$

$$MR^2\omega = (M + 2m)R^2\omega'$$

$$\omega' = \frac{M\omega}{M + 2m}$$

$$10. \vec{F} = 6\hat{i} - 8\hat{j} + 10\hat{k}$$

$$|\vec{F}| = \sqrt{6^2 + 8^2 + 10^2} = 10\sqrt{2}$$

$$m = \frac{|\vec{F}|}{a} = \frac{10\sqrt{2}}{1} = 10\sqrt{2} \text{ kg}$$

11. $\vec{\tau} = \vec{r} \times \vec{F}$
 $\vec{\tau}$ is perpendicular to \vec{r} and \vec{F} .

12. Kepler's 2nd law

$$\Rightarrow \left(\frac{\Delta A}{\Delta t} \right)_{\text{planet}} = \text{constant}$$

$$\frac{A_1}{t_1} = \frac{A_2}{t_2}$$

$$\Rightarrow \frac{2A}{t_1} = \frac{A}{t_2}$$

$$\Rightarrow t_1 = 2t_2$$
13. $mv = \frac{dm}{du} \times \frac{du}{dt} = \frac{dm}{dt} = \text{Rate of flowing mass}$

$$F_{av} = \frac{dm}{dt} \times \frac{v}{2} = \frac{(mv)v}{2} = \frac{mv^2}{2}$$

$$p = \frac{dK}{dt} = \frac{mv^2}{2} \times v = \frac{mv^3}{3}$$
14. Loss of energy = $\frac{1}{2} mv^2 - mgh = 20 \text{ J}$
15. For steady state $\frac{dQ}{dt} = \frac{kA(T_1 - T_2)}{L}$
16. Isochoric \rightarrow Volume constant
17. $P \propto T^4$

$$\frac{P_2}{P_1} = \left(\frac{1000}{500} \right)^4$$

$$P_2 = 16P_1 = 112$$
18. $dU = Q - W = 8400 - 500 = 7900 \text{ J}$
19. $n' = v \left(\frac{c+u}{c-u} \right)$

$$= 600 \left(\frac{330+30}{330-30} \right) = 720 \text{ Hz}$$
20. $v = \omega \sqrt{A^2 - x^2}$

$$= \frac{2\pi}{T} \sqrt{a^2 - \frac{a^2}{4}} = \frac{\pi a \sqrt{3}}{T}$$
21. In SHM, $F_{\text{restoring}} \propto -x$
22. As the coefficient of x is negative, it is moving along +ve x -axis and equating the equation

$$E_y = 2.5 \cos[(2\pi \times 10^6)t - (\pi \times 10^{-2})x]$$
 with $y = A \cos(\omega t - kx)$

$$\omega = 2\pi \times 10^6$$

$$\Rightarrow f = \frac{\omega}{2\pi} = 10^6 \text{ Hz}$$

$$k = \pi \times 10^{-2}$$

$$\Rightarrow \lambda = \frac{2\pi}{k}$$
23. $5\lambda = 4$

$$\lambda = \frac{4}{5}$$

$$k = \frac{2\pi}{\lambda} = \frac{10\pi}{4} = 7.85$$
 wave moves along positive X -direction
24. $\Delta v = \frac{V}{2\ell_1} - \frac{V}{2\ell_2} = \frac{V}{2} \left[\frac{1}{\ell_1} - \frac{1}{\ell_2} \right]$

$$= \frac{1}{2} \sqrt{\frac{T}{\mu}} \left\{ \frac{1}{\ell_1} - \frac{1}{\ell_2} \right\}$$
25. In series, $C_{eq} = \frac{C}{3}$,

$$V_{eq} = 3V$$
26. Total resistance of wire = $12 \Omega \times 2\pi \times 10^{-1}$

$$= 2.4\pi$$
 Resistance of each half = $\frac{2.4\pi}{2} = 1.2\pi$
 and as about diameter both parts are in parallel

$$\text{Req.} = \frac{1.2\pi}{2} = 0.6 \pi \Omega$$
27. $W = MB (\cos\theta_1 - \cos\theta_2)$

$$= 2 \times 10^4 \times 6 \times 10^{-4} (\cos\theta - \cos 60^\circ)$$

$$= 12 \times \frac{1}{2} = 6 \text{ J}$$
28. $\vec{F} = q(\vec{v} \times \vec{B})$

$$= -2 \times 10^{-6} [(2\hat{i} + 3\hat{j}) \times 10^6 \times 2\hat{j}] = -(8N) \hat{k}$$
29. $e = \frac{d\phi}{dt} = \frac{d}{dt} (B\pi r^2)$

$$= 2\pi r B \frac{dr}{dt}$$


$$= 2 \times \pi \times 2 \times 10^{-2} \times 4 \times 10^{-2} \times 2 \times 10^{-3}$$

$$= 3.2 \times 10^{-6} \pi \text{ Vol} = 3.2 \pi \mu \text{V}$$
30. $V = -x^2y - xz^3 + 4$

$$\vec{E} = -\nabla V = - \left(\hat{i} \frac{\partial}{\partial x} + \hat{j} \frac{\partial}{\partial y} + \hat{k} \frac{\partial}{\partial z} \right) (-x^2y - xz^3 + 4)$$

$$= (2xy + z^3) \hat{i} + x^2 \hat{j} + 3xz^2 \hat{k}$$

31.



For loop (3)

$$E_1 - (i_1 + i_2)R - i_1 r_1 = 0$$

For loop (4)

$$-E_1 + i_1 r_1 - i_2 r_2 + E_2 = 0$$

For loop (1)

$$E_2 - (i_1 + i_2)R - i_2 r_2 = 0$$

32. $I_g = 1.0\text{A}$, $G = 60\Omega$, $I = 5.0\text{A}$

$$S = \frac{I_g}{I - I_g} G$$

$$G = \frac{1.0}{5.0 - 1.0} = 60 = 15\ \Omega \text{ in parallel}$$

33. $T = \frac{2\pi m}{2B}$

T is time period

34. $P_{av} = E_{rms} \cdot I_{rms} \cos \phi$

$$= \epsilon \cdot \frac{\epsilon}{Z} \cdot \frac{R}{Z} = \frac{\epsilon^2 R}{Z^2} = \frac{\epsilon^2 R}{R^2 + \left(\omega L - \frac{1}{\omega C}\right)^2}$$

35. $q_A = 4\pi a^2 \sigma$,

$$q_B = -4\pi b^2 \sigma,$$

$$q_C = 4\pi c^2 \sigma, \quad c = a + b$$

$$V_A = \frac{1}{4\pi \epsilon_0} \left(\frac{q_A}{a} + \frac{q_B}{b} + \frac{q_C}{c} \right) = \frac{2\sigma a}{\epsilon_0}$$

$$V_B = \frac{1}{4\pi \epsilon_0} \left(\frac{q_A}{a} + \frac{q_B}{b} + \frac{q_C}{c} \right) = \frac{\sigma}{\epsilon_0} \left(\frac{a^2}{b} - b + c \right) = \frac{\sigma}{\epsilon_0} \left(a + \frac{a^2}{b} \right)$$

$$V_C = \frac{1}{4\pi \epsilon_0} \left(\frac{q_A}{a} + \frac{q_B}{b} + \frac{q_C}{c} \right) = \frac{\sigma}{\epsilon_0} \left(\frac{a^2 - b}{c} + c \right) = \frac{2\sigma a}{\epsilon_0}$$

So, $V_C = V_A \neq V_B$

36. $E = V + Ir$

$$\Rightarrow V = E - Ir$$

Comparing with $y = mx + c$

Slope = $-r$, intercept = E

37. Out of the four structures, when the circular and elliptical loops come out from the field, equal area is not traced in equal interval of time. So any induced in both is not constant.

38. As diamagnetic substances have negative intensity of magnetisation, they are weakly repelled by the external field.

39. No. of photoelectrons emitted is independent of frequency but depends on intensity.

40. No. of photons = $\frac{E}{(hc/\lambda)}$

$$= \frac{9 \times 10^{-3} \times 6.67 \times 10^{-7}}{6.6 \times 10^{-34} \times 3 \times 10^8} = 3 \times 10^{16}$$

41. As a and b have same stopping potential and c has greater stopping potential, then $v_c > v_a = v_b$ as b and c have same saturation current and a has lesser value.

So $I_a < I_b = I_c$

42. $ZX^A \xrightarrow{\alpha} Z-2Y^{A-4} \xrightarrow{2\beta} ZP^{A-4}$

As the resulting daughter and parent nucleus has same atomic number. So they are isotope.

43. $\frac{n(n-1)}{2} = 6$

$$\Rightarrow n = 4$$

For maximum wavelength energy difference between states should be minimum because

$$\lambda = \frac{hc}{\Delta E}$$

So, transition state in $n = 4$ to $n = 3$

44. Energy = $\frac{1}{4\pi \epsilon_0} \frac{Z_1 Z_2}{r_0}$

45. $ZX^A \xrightarrow{\beta} Z+1Y^A \xrightarrow{\alpha} Z-1B^{A-4} \xrightarrow{\gamma} Z-1B^{A-4}$

46. $qV = 2eV$

$$\Rightarrow 1.6 \times 10^{-19} V = 2 \times 1.6 \times 10^{-19} V$$

$$\Rightarrow V = 2V$$


$$E = \frac{V}{d}$$

$$\Rightarrow E = \frac{2V}{4 \times 10^{-8}} = 5 \times 10^7$$

47. $\frac{\sqrt{3}a}{2} = 3.7 \text{ \AA}$


$\Rightarrow a = \frac{2 \times 3.7}{\sqrt{3}} = 4.3 \text{ \AA}$

48. $\lambda_{\text{max}} = \frac{hc}{eV}$
 $= \frac{1242 \text{ eV \AA}}{2.5 \text{ eV}} = 4968 \text{ \AA}$



50. $\Delta I_B = 100 \text{ \mu A}$
 $\Delta I_C = 5 \text{ mA}$

$\beta = \frac{\Delta I_C}{\Delta I_B} = \frac{5 \times 10^{-3}}{100 \times 10^{-6}} = 50$






10g H₂ required O₂ = 80 which is not present 64 g O₂ required 8 g of H₂ and H₂ left = 2g. Thus, O₂ is the limiting reactant and H₂ is excess reactant.

Hence, H₂O formed from 64 of O₂

$= \frac{18}{16} \times 64$

$= 72 \text{ g} = \frac{72}{18} \text{ mole} = 4 \text{ mole}$




53. Maximum no. of electrons in any subshell
 $= 4l + 2$

54. m value $-l$ to $+l$

55. $\Delta H = \text{dissociation energy of reactant} - \text{Bond dissociation of energy of product.}$

$\Delta H = (606.10 + 4 \times 410.5 + 431.37) - (6 \times 410.50 + 336.49)$
 $= -120.0 \text{ kJ/mol}$


56. $K_h = \frac{K_w}{K_b} = \frac{10^{-14}}{1.77 \times 10^{-5}} = 5.65 \times 10^{-10}$



$E^\circ = 0.337 \text{ V}$

$\Delta G = -nFE^\circ_{\text{cell}}$


$= -2 \times F \times 0.337 = -0.674$



$E^\circ = -0.153 \text{ V}$

$\Delta G = +1 \times F \times 0.153$

Final



$\Delta G = -0.52 \text{ V}$

$\Delta G = -nFE^\circ_{\text{cell}}$

$E^\circ_{\text{cell}} = 0.52 \text{ V}$

58. 20 mL of 0.50 M HCl = 20 × 0.050 m mol
 = 1.0 m mol = 1.0 meq. of HCl

30 mL of 0.10 M Ba(OH)₂
 = 30 × 0.1 m mol
 = 3 m mol = 3 × 2 meq
 = 6 meq Ba(OH)₂

1 meq of HCl will neutralize 1 meq of Ba(OH)₂

Ba(OH)₂ left = 5 meq.


Total volume = 50 mL

Ba(OH)₂ conc. in final solution

$= \frac{5}{50} \text{ N} = 0.1 \text{ N} = 0.05 \text{ M}$

$[\text{OH}^-] = 2 \times 0.05 \text{ M} = 0.10 \text{ M}$

Alternatively,



2 m mol of HCl neutralize 1 m mole of Ba(OH)₂

1 m mol of HCl neutralize 0.5 m mol of Ba(OH)₂

Ba(OH)₂ left = 3 - 0.5 m mol = 2.5 m mol

$[\text{Ba(OH)}_2] = \frac{2.5}{50} \text{ M} = 0.05 \text{ M}$

or $[\text{OH}^-] = 2 \times 0.05 = 0.1 \text{ M}$

59. $\text{KE} = 4.4 \times 10^{-19} - 4.0 \times 10^{-19}$

$\text{KE/molecule} = 0.4 \times 10^{-19}$

$\text{KE/atom} = \frac{0.4 \times 10^{-19}}{2}$
 $= 2 \times 10^{-20} \text{ J}$

60. $\frac{1}{3} \frac{-d[\text{H}_2]}{dt} = \frac{1}{2} \frac{d[\text{NH}_3]}{dt}$

$\frac{-d[\text{H}_2]}{dt} = \frac{3}{2} \frac{d[\text{NH}_3]}{dt}$

$\frac{-d[\text{H}_2]}{dt} = \frac{3}{2} \times 2 \times 10^{-4} = 3 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$

61. $\text{Rate} = k[\text{A}][\text{B}]^2$
 $= k[2\text{A}][2\text{B}]^2 = k \times 8[\text{A}][\text{B}]^2$

62. $\alpha = \frac{A_c}{A_\infty} = \frac{8}{400}$

$K_a = C\alpha^2$

$= \frac{1}{32} \times \frac{8}{400} \times \frac{8}{400} = 1.25 \times 10^{-5}$

63. $\Delta T_f = i k_f \cdot m$
 $i = \frac{\Delta T_f}{k_f \cdot m} = \frac{0.00732}{1.86 \times 0.002} = \frac{0.00732}{0.00372}$
 $i = 2$

Compound will be $[\text{Co}(\text{NH}_3)_5]\text{NO}_2\text{NO}_2[\text{Cl}]$
 Total possible ions = 2

64. $-\frac{1}{4} \frac{d(\text{Br}^-)}{dt} = -\frac{1}{3} \frac{d(\text{Br}_2)}{dt}$
 $\frac{d(\text{Br}_2)}{dt} = -\frac{3}{5} \frac{d(\text{Br}^-)}{dt}$

65. for bcc type of unit cell
 $\sqrt{3}a = 4r$
 $r = \frac{\sqrt{3}}{4} a = \frac{1.732 \times 351}{4} = 151.98$

66. $K_c = K_a(\text{CH}_3\text{COOH}) \times \frac{1}{K_a(\text{HCN})}$
 $= 1.5 \times 10^{-5} \times \frac{1}{4.5 \times 10^{-10}}$
 $\cong 3 \times 10^4$

67. For a spontaneous reaction
 $\Delta G = -ve$
 Or at eq. $\Delta G = 0$
 $\Delta H = T\Delta S$
 $T = \frac{\Delta H}{\Delta S} = \frac{170 \times 10^3}{170} = 1000 \text{ K}$

68. $K = \frac{0.693}{t_{1/2}} = \frac{0.693}{1386} = 0.5 \times 10^{-3} \text{ S}^{-1}$

69. Both BF_3 and NO_2^- is sp^2 hybridized.

70. $\text{F}_2 \rightarrow$ reduction potential very high so strongest oxidizing agent.

71. $\text{N}_2 = 14e = \text{B.O.} = 3$
 $\text{N}_2^- = 15e = \text{B.O.} = 2.5$
 $\text{N}_2^{2-} = 16e = \text{B.O.} = 2$

72. $\text{MI} > \text{MBr} > \text{MCl} > \text{MF}$


Down the group increases covalent character

73. Since CaO itself is basic, It will not react with NaOH

74. $W = \frac{E}{96500} \times I \times t$
 $W = \frac{9}{96500} \times 4.0 \times 10^4 \times 6 \times 3600$
 $= 8.1 \times 10^4 \text{ g}$

75. +I stability down the group increase due to inert pair effect
 $\text{Al} < \text{Ga} < \text{In} < \text{Tl}$



76. $\text{Cu} \rightarrow$ fcc lattice



$4r = \sqrt{2} a$
 $r = \frac{1}{2\sqrt{2}} a$
 $r = \frac{1}{2\sqrt{2}} \times 361 = 128$

77. Inter molecular force in alcohol is mainly H-bonding

78. $\text{Cr}^{3+} \rightarrow \text{Is}^2, 2s^2 2p^6, 3s^2 3p^6 3d^3$



Unpaired electron shows colour so absorb visible light.





79. In both TiF_6^{2-} and Cu_2Cl_2 , these no d-electrons or no unpaired electrons, so, these are colourless.

80. As complexes of the type $[\text{MA}_3\text{B}_3]$ can show geometrical isomerism known as facmer isomerism and not optical isomerism. So here $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]^0$ can not show optical isomerism.




81. 7th group \rightarrow largest number of oxidation state.

82. $(\text{CH}_3)_3\text{B} \rightarrow$ is electron deficient compound, so behaves as Lewis acid.


83. $[\text{Ne}]3s^2 3p^3$ has highest ionization energy (half-filled)



$$\Delta G = \Delta H - T\Delta S$$








90. HNO_3 on nitrating mixture acts as a base.



92. Equanil is a diasaccharide [Everyday life]

93. Neoprene is



98. C₂ having → 2-σ bond → sp
C₃ having → 4-σ bond → sp³
C₅ having → 3-σ bond → sp²
C₆ having → 4-σ bond → sp³

99. Gene

100. Thyroxine contains iodine. Its structure is

