

ANSWER KEY (AIPMT-2011)


Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Ans.	4	3	3	4	3	2	1	4	3	3	3	1	2	1	4	4	1	4	3	3
Ques.	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
Ans.	3	4	2	2	3	2	1	3	1	1	2	2	4	4	2	1	3	4	4	3
Ques.	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
Ans.	3	2	3	1	4	4	2	4	1	1	3	3	3	1	3	3	4	4	1	1
Ques.	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
Ans.	4	3	2	1	3	3	4	1	4	2	4	3	2	2	2	2	3	4	3	2
Ques.	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
Ans.	4	4	4	3	2	1	2	2	4	3	3	4	3	3	2	2	3	3	4	3
Ques.	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
Ans.	1	3	3	2	1	1	3	1	2	1	3	1	1	3	2	3	3	2	3	1
Ques.	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140
Ans.	2	4	3	3	2	4	3	2	4	3	4	1	2	4	1	3	4	1	1	3
Ques.	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160
Ans.	2	3	2	1	1	1	3	3	2	3	3	2	3	4	1	3	3	1	1	1
Ques.	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180
Ans.	3	4	4	3	1	2	3	3	3	2	3	3	1	1	1	2	2	2	3	1
Ques.	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200
Ans.	2	3	2	1	2	3	4	4	4	2	3	1	2	2	2	4	1	1	4	1

HINTS & SOLUTIONS

PHYSICS

1. Velocity of light $c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$
So dimension of given expression is equal to velocity $\Rightarrow [LT^{-1}]$

2.




Moment of inertia about an axis passing through one end $= I_{cm} + md^2$

$$= I_0 + M \left(\frac{L}{2} \right)^2 = I_0 + \frac{ML^2}{4}$$

3. $v = \sqrt{2gh} = \sqrt{2 \times 10 \times 20} = 20 \text{ m/sec}$

4.



For (man + lift)
 $T - (60 + 940)g = (60 + 940) \times 1$
 $T = (60 + 940)(10 + 1) = 11000 \text{ N}$

5.

$$P = \vec{F} \cdot \vec{V} = FV \cos \theta$$

Power will be maximum when velocity and $\cos \theta$ will be maximum.

6.

$$\theta = 2t^3 - 6t^2$$

$$\omega = \frac{d\theta}{dt} = 6t^2 - 12t$$

$$\alpha = \frac{d\omega}{dt} = 12t - 12$$

$$\tau = I\alpha$$

Torque will be zero when α is zero

$$\text{so } \alpha = 12t - 12 = 0$$

$$t = 1 \text{ sec}$$

7. If particle move in a circular path with constant speed, the acceleration of the particle is centripetal acceleration

$$a_c = \omega^2 R = \left(\frac{2\pi}{T}\right)^2 R$$

$$a_c = \frac{4\pi^2 R}{T^2} = \frac{4\pi^2}{(0.2\pi)^2} \times 5 \times 10^{-2}$$


$$a_c = 5 \text{ m/sec}^2$$

8. Impulse = $mv_2 - mv_1$
 $= -mv - mv = -2mv$

9. $v_1 r_1 = v_2 r_2$

$$\frac{v_1}{v_2} = \frac{r_2}{r_1}$$

- 10.



$$F_{\text{ex}} = \frac{dP}{dt} = 0 \Rightarrow dP = 0 \Rightarrow P = \text{constant}$$

$$\vec{P}_i = \vec{P}_f$$

$$0 = \vec{P}_{\text{Nu}} + \vec{P}_{\text{Ph}}$$

$$|\vec{P}_{\text{Nu}}| = |\vec{P}_{\text{Ph}}| = \frac{h}{\lambda} = \frac{h\nu}{c}$$

$$\text{Recoil K.E. of nucleus } K.E_{\text{Nu}} = \frac{P_{\text{Nu}}^2}{2M_{\text{Nu}}}$$

$$K.E_{\text{Nu}} = \frac{(h\nu/c)^2}{2M} = \frac{h^2 \nu^2}{2Mc^2}$$

11. Potential energy will increase when work is done by the system against a conservative force.

12. Average acceleration

$$\vec{A}_{\text{avg}} = \frac{\vec{v}_f - \vec{v}_i}{t} = \frac{40\hat{j} - 30\hat{i}}{10}$$

$$\vec{A}_{\text{avg}} = \frac{\sqrt{40^2 + 30^2}}{10} = 5 \text{ m/sec}^2$$

13. Maximum Range

$$R_{\text{max}} = \frac{u^2}{g} = \frac{(20)^2}{10} = 40 \text{ m}$$

14. Work done = area between force v/s displacement curve and displacement axis

$$= (2 \times 4) + \frac{2 \times 5}{2} = 13 \text{ J}$$

15. $\phi_{\text{net}} = \frac{\Sigma q}{\epsilon_0}$

\therefore Net flux does not depend on size of Gaussian surface
 \Rightarrow Flux remains unchanged.

16. $V_A = \frac{kq}{L} + \frac{kq}{L} - \frac{kq}{\sqrt{5}L} - \frac{kq}{\sqrt{5}L}$
 $= \frac{2kq}{L} \left(1 - \frac{1}{\sqrt{5}}\right)$

17. $U = \frac{1}{2} CV^2$

$$= \frac{1}{2} \left(\frac{\epsilon_0 A}{d}\right) V^2$$

$$= \frac{1}{2} (\epsilon_0 Ad) \left(\frac{V}{d}\right)^2 = \frac{1}{2} \epsilon_0 E^2 Ad$$

18. $\therefore \rho = I_g^2 R$
 $36 = I^2(9)$
 $\Rightarrow I_g = 2A$

\therefore In parallel $I \propto \frac{1}{R}$

$$\frac{I_9}{I_6} = \frac{6}{9} \quad \frac{2}{I_6} = \frac{6}{9}$$

$$I_6 = 3A$$

$$\Rightarrow I_{\text{ckt}} = 2 + 3 = 5A$$

$$\Rightarrow V_{2\Omega} = IR = (5)(2) = 10 \text{ volt}$$

19. $I = \frac{E}{R+r}$

$$2 = \frac{E}{2+r} \quad \dots(1)$$

$$0.5 = \frac{E}{9+r} \quad \dots(2)$$

(1) divided by (2)

$$4 = \frac{9+r}{2+r}$$

$$8 + 4r = 9 + r \text{ or } 3r = 1$$

$$\therefore r = \frac{1}{3} \Omega$$


20. At neutral temperature

$$\frac{dE}{dT} = 0$$

21. From $\vec{F} = I(\vec{\ell} \times \vec{B})$

$$\vec{F}_{BC} = -\vec{F}_{AC}$$

$$\vec{F}_{AC} = -\vec{F}$$



22. $E = Pt = mc^2$

$$m = \frac{Pt}{c^2} = \frac{10^6 \times 3600}{(3 \times 10^8)^2}$$

$$m = 40 \mu\text{gm}$$

23. Diamagnetic will be feebly repelled Paramagnetic will be feebly attracted Ferromagnetic will be strongly attracted.

24. $\hat{v} = \hat{E} \times \hat{B}$ or (direction of propagation of waves is $\hat{E} \times \hat{B}$)

25. B will not apply force E field will apply a force opposite to velocity of the electron hence speed will decrease.

26. $e = -\frac{d\phi}{dt}$

27. $I_{\text{rms}} = \frac{E_0 / \sqrt{2}}{1/\omega C}$

28. $\tan \phi = \frac{X_L}{R} = 1, \phi = 45^\circ$

29. $dS = \frac{\Delta Q}{T} = \frac{80 \times 1000}{273} \approx 293 \text{ cal/K}$

30. In isothermal expansion work done against surrounding is negative but work done by gas is positive.

$$\Delta W = +150 \text{ J}$$

$$dU = 0$$

From F.L.O.T.

$$\Delta Q = \Delta W + dU$$

$$\Delta Q = +150 \text{ J}$$

heat is +ve it means heat absorb by gas

31. Motion start from extreme position and for small displacement it is SHM $y = A \cos(\omega t + \phi)$

32. $Y_1 = a \sin(\omega t + kx + 0.57)$

$$Y_2 = a \sin(\omega t + kx + \pi/2)$$

$$\text{Phase difference} = \frac{\pi}{2} - 0.57 = 1 \text{ radian}$$

33. Any function which is converted into single $y = A \sin(\omega t + \phi)$ or $y = A \cos(\omega t + \phi)$ is considered SHM.

34. Frequency is same in both medium

$$n_1 = n_2$$

$$\frac{v_1}{\lambda_1} = \frac{v_2}{\lambda_2}$$

$$\frac{\lambda_2}{\lambda_1} = \frac{v_2}{v_1} = \frac{3500}{350} = 10$$

36. $\frac{1}{\lambda} = RZ^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$

$$R(1)^2 \left(\frac{1}{1^2} - \frac{1}{2^2} \right) = RZ^2 \left(\frac{1}{2^2} - \frac{1}{4^2} \right)$$

$$Z = 2$$

38. Focal length of the lens

$$\frac{1}{f} = (1.5 - 1) \left(\frac{1}{20} - \frac{1}{-20} \right) = \frac{1}{20}$$

$$f = 20 \text{ cm}$$

From lens formula

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$$

$$\frac{1}{v} - \frac{1}{-30} = \frac{1}{20}$$

$$v = 60 \text{ cm}$$

$$\frac{I}{O} = m = \frac{v}{u} = \frac{60}{-30} = -I$$

$$I = -2(O) = -2 \times 2 = -4 \text{ cm}$$

so image will be real inverted and of size 4 cm.

39. $K.E._{\text{max}} = eV_0$

$$V_0 = \frac{K.E._{\text{max}}}{e} = \frac{0.5 \text{ eV}}{e}$$

$$V_0 = 0.5 \text{ volt}$$

40. $\lambda = \frac{h}{\sqrt{2mq\Delta V}} \propto \frac{1}{\sqrt{\Delta V}}$

$$\frac{\lambda_2}{\lambda_1} = \sqrt{\frac{\Delta V_1}{\Delta V_2}} = \sqrt{\frac{25}{100}} = \frac{1}{2}$$

$$\lambda_2 = \frac{\lambda_1}{2}$$

41. $v_{\max} = \sqrt{\frac{2}{m} K.E_{\max}}$
 $v_{\max} = \sqrt{\frac{2}{m} (E_{\text{ph}} - W)}$
 $\frac{v_1}{v_2} = \sqrt{\frac{E_{\text{ph}_1} - W}{E_{\text{ph}_2} - W}} = \sqrt{\frac{1 - 0.5}{2.5 - 0.5}}$
 $\frac{v_1}{v_2} = \frac{1}{2}$
42. Velocity of electron emitted from the electron gun can be increased by potential difference between the anode and filament.
43. $X \rightarrow Y$
 $X : Y = 1 : 15$
A.P. = $\frac{1}{16} = \frac{1}{2^n}$
No. of half life $n = 4$
 $t = nT_{1/2} = 4 \times 50 = 200$ yr.
44. Photoelectron emission take place when certain minimum "frequency" light fall on metal surface.
45. Thermal K.E. \geq Electrostatic P.E.
46. ${}_nX^m \xrightarrow{1\alpha} {}_{n-2}Y^{m-4} \xrightarrow{2\beta^-} {}_nZ^{m-4}$
 α emission decreases mass no. by 4 and atomic no. by 2 and β^- emission increases atomic number by one but leaves mass no. unchanged.
47. $\beta = \frac{\Delta I_C}{\Delta I_B} = \frac{(20-10) \times 10^{-3}}{(300-100) \times 10^{-6}} = 50$
48. By addition of pentavalent impurity only n-type of semiconductor are constructed
49. In FB width of depletion layer is decreased.
50. From theory
- CHEMISTRY**
51. Number of atomic orbitals in an orbit
 $= n^2 = 4^2 = 16$
52. $\Delta G_3 = \Delta G_1 + \Delta G_2$
 $\Rightarrow -2 FE^\circ = -1F \times 0.15 + (-1F \times 0.50)$
 $\Rightarrow -2 FE^\circ = -0.15F - 0.50 F$
 $\Rightarrow -2 FE^\circ = -F (0.15 + 0.50)$
 $\therefore E^\circ = \frac{0.65}{2} = 0.325$ volt
53. Mole fraction of solute = $\frac{1}{56.55} = 0.0177$
54. Average velocity = $\sqrt{\frac{8RT}{\pi M}}$
55. $\text{pOH} = \text{pK}_b + \log \frac{[\text{Salt}]}{[\text{Base}]}$
 $= 4.74 + \log \frac{0.20}{0.30} = 4.74 + (0.301 - 0.477)$
 $= 4.74 - 0.176 = 4.56$
 $\therefore \text{pH} = 14 - 4.56 = 9.44$
56. $\frac{r_A}{r_B} = \sqrt{\frac{M_B}{M_A}}$
 $\Rightarrow \frac{v_A}{t_A} \times \frac{t_B}{v_B} = \sqrt{\frac{M_B}{M_A}} \Rightarrow \frac{10}{20} = \sqrt{\frac{M_B}{49}}$
 $\Rightarrow \frac{1}{4} = \frac{M_B}{49} \quad \therefore M_B = \frac{49}{4} = 12.25$
57. For an ideal gas, for free expansion
 $q = 0 ; \Delta T = 0$ and $w = 0$
58. $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g}) ; K_1$
 $2\text{NO}(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g}) ; K_2$

 $\text{N}_2(\text{g}) + 2\text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g}) ; K = K_1 \times K_2$
 $\therefore \text{For } \text{NO}_2(\text{g}) \rightleftharpoons \frac{1}{2} \text{N}_2(\text{g}) + \text{O}_2(\text{g}) ;$
 $K' = \left[\frac{1}{K_1 \cdot K_2} \right]^{1/2}$
59. $x/m = P \times T$ is the incorrect relation.
60. $\Delta S_{\text{vap}} = \frac{\Delta H_{\text{vap}}}{T} = \frac{30\text{KJmol}^{-1}}{300\text{K}} = 100 \text{ J mol}^{-1} \text{ K}^{-1}$
61. Fact
62. $E^\circ_{\text{cell}} = E^\circ_{\text{cathode(RP)}} - E^\circ_{\text{anode(RP)}}$
 $= 0.15 - (-0.74)$
 $= +0.89 \text{ V}$
63. Fact
64. $\therefore \Delta G^\circ = -nFE^\circ$
and $\Delta G^\circ = -RT \log_e K_{\text{eq}}$

