COMMON ENTRANCE TEST - 2011

DATE	SUBJECT	TIME
28-04-2011	CHEMISTRY	02.30 PM to 03.50 PM

MAXIMUM MARKS	TOTAL DURATION	MAXIMUM TIME FOR ANSWERING
60	80 MINUTES	70 MINUTES

Г	MENTION YOUR	QUESTION BOOKLET DETAILS	
	CET NUMBER	VERSION CODE	SERIAL NUMBER
		A-1	727393

DOs:

- 1. Check whether the CET No. has been entered and shaded in the respective circles on the OMR answer sheet.
- 2. This Question Booklet is issued to you by the Invigilator after the 2nd Bell, i.e., after 02.30 p.m.
- 3. The Serial Number of this question booklet should be entered on the OMR answer sheet.
- 4. The Version Code of this question booklet should be entered on the OMR answer sheet and the respective circles should be shaded completely.
- 5. Compulsory sign at the bottom portion of the OMR answer sheet in the space provided.

DON'Ts:

- 1. The timing and marks printed on the OMR answer sheet should not be damaged/mutilated/spoiled.
- 2. The 3rd Bell rings at 02.40 p.m. till then;
 - Do not remove the seal/staple present on the right hand side of this question booklet.
 - Do not look inside this question booklet.
 - Do not start answering on the OMR answer sheet.

IMPORTANT INSTRUCTIONS TO CANDIDATES

- This question booklet contains 60 questions and each question will have one statement and four distracters (four different options / choices).
- After the 3rd Bell is rung at 02.40 p.m., remove the seal/staple present on the right hand side of this question booklet and start answering on the OMR answer sheet.
- 3. During the subsequent 70 minutes:
 - Read each question carefully.
 - Choose the correct answer from out of the four available distracters (options/choices) given under each question/statement.
 - Completely darken/shade the relevant circle with a BLUE OR BLACK INK BALLPOINT PEN against the question number on the OMR answer sheet.

CORRECT METHOD OF SHADING THE CIRCLE ON THE OMR SHEET IS AS SHOWN BELOW:



- 4. Please note that even a minute unintended ink dot on the OMR sheet will also be recognized and recorded by the scanner. Therefore, avoid multiple markings of any kind on the OMR answer sheet.
- 5. Use the space provided on each page of the question booklet for Rough Work. Do not use the OMR answer sheet for the same.
- 6. After the last bell is rung at 03.50 p.m., stop writing on the OMR answer sheet and affix your LEFT HAND THUMB IMPRESSION on the OMR answer sheet as per the instructions.
- 7. Hand over the OMR answer sheet to the room Invigilator as it is.
- 8. After separating and retaining the top sheet (KEA Copy), the Invigilator will return the bottom sheet replica (Candidate's copy) to you to carry home for self-evaluation.
- 9. Preserve the replica of the OMR answer sheet for a minimum period of ONE year.

SR - 49 Turn Over



CHEMISTRY

- Which one of the following statements is FALSE?
 - During roasting, moisture is removed from the ore.
 - The ore is freed from almost all nonmetallic impurities.
 - Calcination of ore is carried out in the absence of any blast of air.
 - The concentrated zinc blende is subjected to calcination during its extraction by pyrometallurgy.
- Which one of the following sets of quantum numbers represents the highest energy level in an atom?

1)
$$n = 4$$
, $l = 0$, $m = 0$, $s = +\frac{1}{2}$ 2) $n = 3$, $l = 1$, $m = 1$, $s = +\frac{1}{2}$

2)
$$n=3$$
, $l=1$, $m=1$, $s=+\frac{1}{2}$

3)
$$n=3$$
, $l=2$, $m=-2$, $s=+\frac{1}{2}$ 4) $n=3$, $l=0$, $m=0$, $s=+\frac{1}{2}$

4)
$$n=3$$
, $l=0$, $m=0$, $s=+\frac{1}{2}$

- When O_2 is converted into O_2^+ ;
 - both paramagnetic character and bond order increase
 - bond order decreases
 - paramagnetic character increases
 - 4) paramagnetic character decreases and the bond order increases
- In chromite ore, the oxidation number of iron and chromium are respectively

$$1) +3, +2$$

$$2) +3, +6$$

$$3) +2, +6$$

$$(4)$$
 +2, +3

- The number of naturally occurring p-block elements that are diamagnetic is
 - 1) 18

3)

- 6. If the energies of the two photons are in the ratio of 3:2, their wavelengths will be in the ratio of
 - 1) 9:4

2) 2:3

3) 1:2

4) 3:2


- 7. Which one of these is NOT TRUE for benzene?
 - 1) There are three carbon-carbon single bonds and three carbon-carbon double bonds.
 - 2) It forms only one type of monosubstituted product.
 - 3) The bond angle between carbon-carbon bonds is 120°.
 - 4) Heat of hydrogenation of benzene is less than the theoretical value.
- 8. Generally, the first ionization energy increases along a period. But there are some exceptions. The one which is NOT an exception is
 - 1) Na and Mg

Be and B

3) N and O

4) Mg and Al

9. Out of the given two compounds, the vapour pressure of B at a particular temperature is




2) higher than that of A

3) same as that of A

4) higher or lower than A depending on the size of the vessel





. .

10. Increasing order of carbon-carbon bond length for the following is

 C_2H_4

 C_2H_2

 C_6H_6

 C_2H_0

(A)

(A)

(B)

(C

(D)

1) B < C < A < D

2) C < B < A < D

3) B < A < C < D

4) D < C < A < B

- - 1) 31.5

2) 75

3) 25

- 4) 40.2
- 12. 50 cm³ of 0.2 N HCl is titrated against 0.1 N NaOH solution. The titration was discontinued after adding 50 cm³ of NaOH. The remaining titration is completed by adding 0.5 N KOH. The volume of KOH required for completing the titration is
 - 1) 10 cm^3

2) 12 cm^3

3) $16.2 \, \text{cm}^3$

- 4) 21.0 cm³
- 13. The rms velocity of hydrogen is $\sqrt{7}$ times the rms velocity of nitrogen. If T is the temperature of the gas, which of the following is true?
 - 1) $T_{N_2} = T_{H_2}$

 $2) \quad T_{H_2} = \sqrt{7} \, T_{N_2}$

3) $T_{N_2} = 2T_{H_2}$

- 4) $T_{N_2} = \sqrt{7} T_{H_2}$
- 14. 25 g of each of the following gases are taken at 27°C and 600 mm pressure. Which of these will have the least volume?
 - 1) HBr

2) HCl

3) HF

- 4) HI
- 15. The amount of heat evolved when 500 cm³ of 0.1 M HCl is mixed with 200 cm³ of 0.2 M NaOH is
 - 1) 1.292 kJ

2) 2.292 kJ

3) 0.292 kJ

4) 22.9 kJ



- The enthalpy of vaporization of benzene is +35.3 kJ/mol at its boiling point of 80°C. The entropy change in the transition of vapour to liquid at its boiling point is [in J mol-1 K-1].
 - 1) -100

2) +100

3) +342

- -342
- Based on the first law of thermodynamics, which one of the following is correct?
 - 1) For an isothermal process, q = +w
 - 2) For an isochoric process, $\Delta U = -q$
 - 3) For an adiabatic process, $\Delta U = -w$
 - For a cyclic process, q = -w
- Consider the following gaseous equilibria with equilibrium constants K_1 and K_2 respectively.

$$SO_{2(g)} + \frac{1}{2}O_{2(g)} \Longrightarrow SO_{3(g)}$$

$$2SO_{3(g)} \Longrightarrow 2SO_{2(g)} + O_{2(g)}$$

The equilibrium constants are related as

1) $2K_1 = K_2^2$


2) $K_1^2 = \frac{1}{K_0}$

3) $K_2^2 = \frac{1}{K}$

- 4) $K_2 = \frac{2}{K^2}$
- 19. During the adsorption of Krypton on activated charcoal at low temperature;

 - 1) $\Delta H < 0$ and $\Delta S < 0$ 2) $\Delta H > 0$ and $\Delta S < 0$
 - 3) $\Delta H > 0$ and $\Delta S > 0$
- 4) $\Delta H < 0$ and $\Delta S > 0$
- 20. For the reversible reaction, $A_{(g)} + B_{(g)} = C_{(g)} + D_{(g)} \Delta G^0 = -350 \,\mathrm{kJ}$, which one of the following statements is true?
 - The reaction is thermodynamically nonfeasible.
 - The entropy change is negative.
 - Equilibrium constant is greater than one.
 - The reaction should be instantaneous.

21. Identify B and D in the following sequence of reactions.



- 1) Methanol and bromoethane
- 2) Ethyl hydrogen sulphate and alcoholic KOH
- 3) Ethyl hydrogen sulphate and aqueous KOH
- 4) Ethanol and alcoholic KOH
- 22. The compound which gives turbidity immediately with Lucas reagent at room temperature is
 - 1) butan-1-ol

- 2) butan-2-ol
- 3) 2-methyl propan-2-ol
- 4) 2-methyl propan-1-ol
- 23. Ethyl benzene CANNOT be prepared by
 - 1) Wurtz reaction

- 2) Wurtz-Fittig reaction
- 3) Friedel-Crafts reaction
- 4) Clemmensen reduction
- 24. 1.2 g of organic compound on Kjeldahlization liberates ammonia which consumes 30 cm³ of 1 N HCl. The percentage of nitrogen in the organic compound is
 - 1) 30

2) 35

3) 46.67

- 4) 20.8
- 25. Carbon cannot reduce Fe_2O_3 to Fe at a temperature below 983 K because
 - 1) free energy change for the formation of CO is more negative than that of Fe_2O_3
 - 2) CO is thermodynamically more stable than Fe_2O_3
 - 3) carbon has higher affinity towards oxygen than iron
 - 4) iron has higher affinity towards oxygen than carbon



26.	The yellow precipitate formed d	uring the chromyl chloride test is chemically	
	1) chromic acid	2) lead chromate	
	3) lead acetate	4) sodium chromate	
27.		ed between 10 cm³ of molten zinc and 100 cm³ of molten	
	lead at 800°C. The percentage of	f silver still left in the lead layer is approximately	
	1) 2	2) 5	
	3) 3	4) 1	
28.	Which one of the following is tr	ue?	
	1) NaOH is used in the concentration of bauxite ore.		
	2) NaOH is a primary standard in volumetric analysis.		
	3) Manganous hydroxide	is soluble in excess of NaOH solution.	
	4) NaOH solution does no	ot react with Cl_2 .	
29.	In Ramsay and Rayleigh's isola	ation of noble gases from air, the nitrogen of the air	
	is finally converted into		
	1) NaNO ₂ only	2) NO and NO ₂	
	3) NaNO ₃ only	4) NaNO ₂ and NaNO ₃	
30.	The spin only magnetic moment	of Fe^{2+} ion (in BM) is approximately	
.,	1) 4	2) 7	
	3) 5	4) 6	

- 31. The IUPAC name of the complex $\left[Co(NH_3)_4 Cl_2 \right] Cl$ is
 - . 1) dichloro tetraammine cobalt (III) chloride
 - 2) tetraammine dichloro cobalt (III) chloride
 - 3) tetraammine dichloro cobalt (II) chloride
 - 4) tetraammine dichloro cobalt (IV) chloride
- - 1) 287×10^{-3}

2) 143.5×10^{-3}

3) 143.5×10^{-2}

- 4) 287×10^{-2}
- 33. The following data were obtained during the first order decomposition of $2A_{(g)} \to B_{(g)} + C_{(s)}$ at a constant volume and at a particular temperature.

Sr. No.	Time	Total pressure in Pascal
1	At the end of 10 min	300
2	After completion	200

The rate constant in min⁻¹ is

1) 0.0693

2) 69.3

3) 6.93

- 4) 6.93×10^{-4}
- 34. The time required for 100% completion of a zero order reaction is
 - 1) al

 $2) \frac{a}{2k}$

3) $\frac{a}{k}$

- $\frac{2k}{a}$
- - 1) 0.01

2) 0.1

3) 0.02

4) 0.001



- pH value of which one of the following is NOT equal to one?
 - 1) 0.1 M CH, COOH
 - 2) 0.1 M HNO₃
 - 3) $0.05 \text{ M} H_2 SO_4$
 - 4) 50 cm³ 0.4 M HCl + 50 cm³ 0.2 M NaOH
- A buffer solution contains 0.1 mole of sodium acetate dissolved in 1000 cm3 of 0.1 M acetic acid. To the above buffer solution, 0.1 mole of sodium acetate is further added and dissolved. The pH of the resulting buffer is
 - 1) pK_a

3) $pK_a - Log 2$

- 2) $pK_a + 2$ 4) $pK_a + Log 2$
- 38. H_0S is passed into one dm³ of a solution containing 0.1 mole of Zn^{2+} and 0.01 mole of Cu^{2+} till the sulphide ion concentration reaches 8.1 × 10-19 moles. Which one of the following statements is true?

 $[K_{sp} \text{ of } ZnS \text{ and } CuS \text{ are } 3 \times 10^{-22} \text{ and } 8 \times 10^{-36} \text{ respectively}]$

- 1) Only ZnS precipitates
- 2) Both CuS and ZnS precipitate
- 3) Only CuS precipitates
- 4) No precipitation occurs
- 39. E_1 , E_2 and E_3 are the emfs of the following three galvanic cells respectively:
 - (i) $Zn(s) | Zn^{2+}(0.1M) | Cu^{2+}(1M) | Cu(s)$
 - (ii) $Zn(s) | Zn^{2+}(1M) | Cu^{2+}(1M) | Cu(s)$
 - (iii) $Zn(s) | Zn^{2+}(1M) | Cu^{2+}(0.1M) | Cu(s)$

Which one of the following is true?

1) $E_2 > E_1 > E_3$

2) $E_1 > E_2 > E_3$

3) $E_3 > E_1 > E_2$

- 4) $E_3 > E_2 > E_1$
- 40. 0.023 g of sodium metal is reacted with 100 cm3 of water. The pH of the resulting solution is
 - 10

- A 1 41. The standard emf of a galvanic cell involving 2 moles of electrons in its redox reaction is 0.59 V. The equilibrium constant for the redox reaction of the cell is $2) 10^{5}$ $1) - 10^{20}$ $4) 10^{10}$ 3) 10 42. 9.65 coulombs of electric current is passed through fused anhydrous MgCl2. The magnesium metal thus obtained is completely converted into a Grignard reagent. The number of moles of Grignard reagent obtained is 2) 1 × 10⁻⁴ 1) 5×10^{-4} 4) 1×10^{-5} 3) 5×10^{-5} 43. The empirical formula of a nonelectrolyte is CH2O. A solution containing 3 g of the compound exerts the same osmotic pressure as that of 0.05 M glucose solution. The molecular formula of the compound is 2) $C_2H_4O_2$ 1) CH₂O 4) $C_3H_6O_3$ 3) $C_4 H_8 O_4$ 44. Which one of the following is a covalent crystal? 2) Ice 1) Rock salt Dry ice 3) Quartz Which one of the following DOES NOT involve coagulation? 1) Clotting of blood by the use of ferric chloride
 - (Space for Rough Work)

2) Formation of delta region

Peptization

3) Treatment of drinking water by potash alum

0

- 47. Which one of the nitrogen atoms in $H_2N-NH-C-NH_2$ is the most nucleophilic? I
 - 1) III
 - 2) 1
 - 3) II
 - 4) All three nitrogen atoms are equally strong nucleophilic centers
- 48. The maximum number of possible optical isomers in 1-bromo-2-methyl cyclobutane is ...
 - 1) 4

2) 2

3) 8

- 4) 16
- 49. Which one of the following is the most energetic conformation of cyclohexane?
 - 1) Boat


2) Twisted boat

3) Chair

- 4) Half chair
- 50. Which one of the following is an intermediate in the reaction of benzene with CH_3Cl in the presence of anhydrous $AlCl_3$?
 - 1) Cl+

2) CH.

 CH_3^+



- 51. Which one of the following is NOT TRUE for the hydrolysis of t-butyl bromide with aqueous NaOH?
 - 1) Reaction occurs through the S_N1 mechanism.
 - 2) The intermediate formed is a carbocation.
 - 3) Rate of the reaction doubles when the concentration of alkali is doubled.
 - 4) Rate of the reaction doubles when the concentration of t-butyl bromide is doubled.
- **52.** Following is the substitution reaction in which -CN replaces -Cl.

$$R-Cl + KCN \xrightarrow{\Delta} R-CN + KCl$$
 $(alcoholic)$

To obtain propanenitrile, R-Cl should be

1) chloroethane

2) 1-chloropropane

3) chloromethane

- 4) 2-chloropropane
- 53. The conversion of m-nitrophenol to resorcinol involves respectively
 - 1) hydrolysis, diazotization and reduction
 - 2) diazotization, reduction and hydrolysis
 - 3) hydrolysis, reduction and diazotization
 - 4) reduction, diazotization and hydrolysis
- 54. Formic acid is a stronger acid than acetic acid. This can be explained using
 - 1) +M effect

2) -I effect

3) +I effect

- 4) -M effect
- 55. The reagent with which both acetaldehyde and acetone react is
 - 1) Fehling's solution
- 2) I₂ / NaOH

3) Tollens' reagent

4) Carbonic acid

(Space for Rough Work)

collegedunia

- 56. Which of the following gives an aldehyde on dry distillation?
 - 1) Calcium formate + calcium acetate
 - 2) Calcium acetate + calcium benzoate
 - 3) Calcium acetate
 - 4) Calcium benzoate
- 57. α-maltose consists of
 - 1) one α -D-glucopyranose unit and one β -D-glucopyranose unit with 1-2 glycosidic linkage
 - 2) two α-D-glucopyranose units with 1-2 glycosidic linkage
 - 3) two β -D-glucopyranose units with 1-4 glycosidic linkage
 - 4) two α-D-glucopyranose units with 1-4 glycosidic linkage
- 58. Which one of the following DOES NOT correctly match with each other?
 - 1) Silk-polyamide

2) Lipase-enzyme

3) Butter-fat


- 4) Oxytocin-enzyme
- 59. In an alkaline medium, glycine predominantly exists as/in a/an
 - 1) cation

2) anion

3) zwitterion

4) covalent form

60. The IUPAC name of



1) but-3-enoic acid

- 2) but-1-enoic acid
- 3) pent-4-enoic acid
- 4) prop-2-enoic acid