



Marking scheme Compartment – 2019

CHEMISTRY (043)/ CLASS XII



56/1/1







Q.No	Value Points	Marks
	SECTION A	
1	2-3% solution of iodine in alcohol-water mixture. It acts as an antiseptic.	½, ½
2	No unpaired electron.	1
	OR	
2	$[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-}$; $\text{C}_2\text{O}_4^{2-}$ is a didentate / chelating ligand so it is more stable	½, ½
3		1
	OR	
3	$\text{CH}_3\text{-CH}_2\text{-CH}_3$	1
4	Glycogen	1
5	$[\text{CF}_2\text{-CF}_2]_n$	1
	SECTION B	
6	a) Conductivity of a solution at any given concentration is the conductance of one unit volume of solution kept between two platinum electrodes with unit area of cross-section. Molar conductivity is the conductivity of solution for 1M solution. b) Because number of ions per unit volume that carry the current in a solution decreases.	½, ½ 1
7	Since its a first order reaction, a) Unit of rate constant is $\text{s}^{-1} / \text{time}^{-1}$ b) $t_{1/2} = \frac{0.693}{k}$ $= \frac{0.693}{5.5 \times 10^{-14}}$ $= 1.26 \times 10^{13} \text{ s}$ (or any other unit of time)	1 ½, ½
8	a)  or b) Phosphine is formed, $\text{P}_4 + 3\text{NaOH} + 3\text{H}_2\text{O} \rightarrow \text{PH}_3 + 3\text{NaH}_2\text{PO}_2$ (Award full marks if only balanced equation is given)	1 ½, ½
9	a) 5f electrons in actinoids have poorer shielding effect than 4f electrons in lanthanoids. b) Fe, due to more unpaired electrons leading to stronger metallic bonding.	1 ½, ½
10.	$\text{CH}_3\text{-C}=\text{CH}_2$ $\quad \quad $ $\quad \quad \text{CH}_3$; 2--Methylpropene	1,1
	OR	



10.	$\begin{array}{c} \text{H} \\ \\ -\text{C}-\text{C}^+ \\ \end{array} + \text{H}_2\ddot{\text{O}} \rightleftharpoons \begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ -\text{C}-\text{C}-\text{O}^+-\text{H} \\ \quad \end{array}$ <p>a)</p> <p>b) 2,6-dimethylphenol</p>	1
11	<p>For fcc,</p> $r = \frac{a}{2\sqrt{2}}$ $a = 2r \times \sqrt{2}$ $= 2 \times 125\text{pm} \times 1.414$ $= 353.5 \text{ pm}$	<p>½</p> <p>½</p> <p>1</p>
OR		
11	$d = \frac{zM}{a^3 N_A}$ $a^3 = \frac{4 \times 99 \text{ g mol}^{-1}}{3.04 \text{ g cm}^{-3} \times 6.022 \times 10^{23} \text{ mol}^{-1}}$ $a^3 = 21.6 \times 10^{-23} \text{ cm}^3 \quad (\text{Deduct half marks if correct unit is not given})$	<p>½</p> <p>½</p> <p>1</p>
12	<p>a) X₂Y₃</p> <p>b) Metal excess defect due to anionic vacancies / F-centres</p>	<p>1</p> <p>1</p>
SECTION C		
13	$\text{Zn}^{2+} + 2e \rightarrow \text{Zn}(s)$ $E_{\text{Zn}^{2+}/\text{Zn}} = E^\circ_{\text{Zn}^{2+}/\text{Zn}} - \frac{0.059}{2} \log \frac{1}{[\text{Zn}^{2+}]}$ $E_{\text{Zn}^{2+}/\text{Zn}} = -0.76 - \frac{0.059}{2} \log \frac{1}{[0.01]}$ $= -0.76 - \frac{0.059}{2} \log 10^2$ $= -0.76 - 0.059 \text{ V}$ $E_{\text{Zn}^{2+}/\text{Zn}} = -0.819 \text{ V} \quad (\text{Deduct half marks if correct unit is not given})$	<p>1</p> <p>1</p> <p>1</p>
14	$\log \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$ $\log \frac{6 \times 10^{-2}}{2 \times 10^{-2}} = \frac{E_a}{2.303 \times 8.314 \text{ J K}^{-1} \text{ mol}^{-1}} \left[\frac{1}{300} - \frac{1}{320} \right] \text{ K}^{-1}$ $\log 3 = \frac{E_a}{19.15 \text{ J mol}^{-1}} \left[\frac{320-300}{300 \times 320} \right]$ $0.4771 = \frac{E_a}{19.15 \text{ J mol}^{-1}} \left[\frac{20}{300 \times 320} \right]$ $E_a = 43855 \text{ J mol}^{-1} \text{ or } 43.855 \text{ kJ mol}^{-1} \quad (\text{Deduct half marks if correct unit is not given})$	<p>½</p> <p>1</p> <p>½</p> <p>1</p>
15	<p>a) Because acid formed in the reaction provides H⁺ which acts as a catalyst in hydrolysis.</p> <p>b) The solution becomes colourless because the molecules of methylene blue / dye get adsorbed on the surface of charcoal.</p> <p>c) Milk / Vanishing cream (or any other suitable example)</p>	<p>1</p> <p>1</p> <p>1</p>
OR		
15	<p>a) Colloids which act as electrolytes at low concentration and show colloidal</p>	1




	behaviour at high concentration. b) The movement of colloidal particles towards a particular electrode under the influence of an electric field. c) The potential difference between fixed layer and the diffused layer of opposite charges.	1 1								
16	a) Calamine is an ore of Zn while malachite is an ore of copper. / Calamine is $ZnCO_3$ while malachite is $CuCO_3 \cdot Cu(OH)_2$ b) Zn is more reactive than Cu, so reduction will be faster in case of Zn. c) Cryolite makes alumina a good conductor of electricity and lowers melting point of the mixture.	1 1 1								
OR										
16	a) <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;">Cast iron</th> <th style="width: 50%;">Pig iron</th> </tr> </thead> <tbody> <tr> <td>1. It contains low percentage of carbon and other impurities</td> <td>1. It contains high percentage of carbon and other impurities</td> </tr> <tr> <td>2. It is moulded pig iron</td> <td>2. It is directly obtained from blast furnace</td> </tr> <tr> <td>3. It is less brittle</td> <td>3. It is more brittle</td> </tr> </tbody> </table> b) Zone refining – impurities are more soluble in the melt than the solid state of the metal. (Any two)	Cast iron	Pig iron	1. It contains low percentage of carbon and other impurities	1. It contains high percentage of carbon and other impurities	2. It is moulded pig iron	2. It is directly obtained from blast furnace	3. It is less brittle	3. It is more brittle	1+ 1 1
Cast iron	Pig iron									
1. It contains low percentage of carbon and other impurities	1. It contains high percentage of carbon and other impurities									
2. It is moulded pig iron	2. It is directly obtained from blast furnace									
3. It is less brittle	3. It is more brittle									
17	a) i) $Na_2Cr_2O_7 + 2 KCl \rightarrow K_2Cr_2O_7 + 2 NaCl$ ii) $5SO_3^{2-} + 2MnO_4^- + 6H^+ \rightarrow 2Mn^{2+} + 3H_2O + 5SO_4^{2-}$ b) Orange colour of $Cr_2O_7^{2-}$ changes to yellow due to formation of chromate ion / $Cr_2O_7^{2-} + 2 OH^- \rightarrow 2 CrO_4^{2-} + H_2O$ (yellow)	1 1 1								
18	 a) b) In the presence of strong field ligand, CO, the unpaired d-electrons of Ni pair up so $[Ni(CO)_4]$ is diamagnetic but Cl^- being a weak ligand is unable to pair up the unpaired electrons, so $[Ni(Cl_4)]^{2-}$ is paramagnetic.	1,1 1								
OR										
18	a) Ligand that can ligate through two different atoms. b) Ligands can be arranged in a series in the order of increasing field strength. / An experimentally determined series based on absorption of light by complexes with different ligands. c) Complexes in which a metal is bound to more than one kind of ligands / donor groups	1 1 1								
19.	a) i) Propene  ii)	1 1								

	b) Phenyl isocyanides / 	1
20.	a) i)  ii) $\text{RCOR} + \text{R}'\text{MgX} \longrightarrow \text{R}-\overset{\text{R}'}{\underset{\text{R}}{\text{C}}}-\text{OMgX} \xrightarrow{\text{H}_2\text{O}} \text{R}-\overset{\text{R}'}{\underset{\text{R}}{\text{C}}}-\text{OH} + \text{Mg(OH)X}$ (R, R' = CH ₃) b) Due to hydrogen bonding in propanol	1 1 1
21	a) X =  / benzene diazonium chloride , Y =  / Cyanobenzene b) Aniline gets protonated and is converted to anilinium ion which is m-directing	1+ 1 1
22	a) i)  or n-Hexane is formed ii)  or Saccharic acid is formed b) Starch is a polymer of α-glucose while cellulose is a polymer of β-glucose.	1 1 1
23	a) Ethane-1,2-diol and benzene-1,6-dicarboxylic acid / ethylene glycol and phthalic acid b) It represents 6 carbon atoms present in both the monomer units. c) PHBV (poly-β-hydroxybutyrate-co-β-hydroxyvalerate) / Nylon-2-Nylon-6	1 1 1
24	a) Antiseptics – chemicals applied on living tissues to prevent the growth of microorganisms while disinfectants are applied on non-living tissues. Example: Antiseptic- Dettol , Disinfectants- 1% phenol (or any other suitable example) b) It is needed by diabetic persons as it is excreted from the body in urine unchanged. / Reduces calories intake	½ + ½ ½ + ½ 1
OR		
24	i) Chemical compounds used for the treatment of stress and mental diseases. ii) Chemical compounds which stop overproduction of acid in stomach. iii) Chemical compounds which reduce or abolish pain without disturbing nervous system.	1 1 1
SECTION D		
25	a) $\Delta T_f = T_f^0 - T_f = 273.15 - 271 \text{ K} = 2.15 \text{ K}$ $\Delta T_f = K_f m$	½

	$\Delta T_f = K_f \times \frac{w_b \times 1000}{M_B \times w_A}$ $K_f = \frac{2.15K \times 342 \text{ g mol}^{-1} \times 95 \text{ g}}{5 \text{ g} \times 1000 \text{ g Kg}^{-1}}$ $= 13.97 \text{ K kg mol}^{-1}$ <p>For 5% glucose in water,</p> $\Delta T_f = K_f m$ $\Delta T_f = K_f \times \frac{w_b \times 1000}{M_B \times w_A}$ $= \frac{13.97 \text{ K Kg mol}^{-1} \times 5 \text{ g} \times 1000 \text{ g Kg}^{-1}}{180 \text{ g mol}^{-1} \times 95 \text{ g}}$ $= 4.08 \text{ K}$ $T_f = T_f^{\circ} - \Delta T_f = 273.15 - 4.08 \text{ K} = 269.07 \text{ K}$ <p>b) It is due to the fact that KCl dissociates to give K^+ and Cl^- ions whereas urea does not dissociate into ions</p> <p>c) Liquids having similar nature and polarities / which obey Raoult's law</p>	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1</p>
	OR	
25	$\Delta T_f = K_f m$ $\Delta T_f = K_f \times \frac{w_b \times 1000}{M_B \times w_A}$ $M_b = K_f \times \frac{w_b \times 1000}{\Delta T_f \times w_A}$ $= 5.12 \times \frac{1 \times 1000}{0.40 \times 50}$ $= 256 \text{ g mol}^{-1}$ <p>b) Higher the value of K_H, lower will be the solubility of a gas in the liquid.</p> <p>c) Low level of oxygen in the blood and tissues of people at high altitudes leads to the condition of anoxia.</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
26	<p>A= S_8 / Sulphur</p> $S_8 + 8 O_2 \rightarrow 8SO_2 \quad / \quad S + O_2 \rightarrow SO_2$ <p>B= SO_2</p> $Ca(OH)_2 (aq) + SO_2 (g) \rightarrow CaSO_3 (s) + H_2O$ <p style="text-align: center;">(milky)</p> <p>Decolourises $KMnO_4$</p> $2KMnO_4 + 5 SO_2 + 2H_2O \rightarrow 2 H_2SO_4 + 2MnSO_4 + K_2SO_4 \quad / \quad 2 MnO_4^- + 5 SO_2 + 2H_2O \rightarrow 4 H^+ + 2Mn^{2+} + 5 SO_4^{2-}$ <p>Reduces Fe^{3+} to Fe^{2+}</p> $2Fe^{3+} + SO_2 + 2 H_2O \rightarrow 2 Fe^{2+} + SO_4^{2-} + 4H^+$	<p>1</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1</p>
	OR	
26	<p>a) $H_2Te > H_2Se > H_2S > H_2O$</p> <p>b) PCl_4^-, as phosphorous has 10 e which cannot be accommodated in sp^3 orbitals.</p> <p>c) Rhombic sulphur</p> <p>d) H_3PO_4</p>	<p>1</p> <p>1/2, 1/2</p> <p>1</p> <p>1</p>



	e) PCl_3 hydrolyses in presence of moisture to give fumes of HCl / $\text{PCl}_3 + 3\text{H}_2\text{O} \rightarrow \text{H}_3\text{PO}_3 + 3\text{HCl}$	1
27	<p>a) Compound = Benzaldehyde or $\text{C}_6\text{H}_5\text{CHO}$</p> <p>Reactions</p> <p>Reaction with 2,4-DNP</p>  <p>With Tollens reagent</p> $\text{RCHO} + 2[\text{Ag}(\text{NH}_3)_2]^+ + 3\bar{\text{O}}\text{H} \longrightarrow \text{RCO}\bar{\text{O}} + 2\text{Ag} + 2\text{H}_2\text{O} + 4\text{NH}_3 \text{ (Where R= -C}_6\text{H}_5\text{)}$ <p>Cannizzaro</p> $2 \text{C}_6\text{H}_5\text{-CHO} + \text{Conc. NaOH} \longrightarrow \text{C}_6\text{H}_5\text{-CH}_2\text{OH} + \text{C}_6\text{H}_5\text{-COONa}$ <p>b) i) Add neutral FeCl_3 to both the compounds, phenol will give violet colour. ii) Add NaHCO_3 to both the compounds, benzoic acid acid will give brisk effervescence of CO_2</p>	<p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>1</p> <p>1</p>
	OR	
27	<p>a) A= CH_3COOH B= CH_3COCl C= CH_3CONH_2 D= CH_3NH_2</p> <p>b) $\text{HCOOH} < \text{ClCH}_2\text{COOH} < \text{CCl}_3\text{COOH} < \text{CF}_3\text{COOH}$</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>