SUBJECT: CHEMISTRY	DAY-2			
SESSION: AFTERNOON	TIME: 02.30 P.M. TO 03.50 P.M.			

MAXIMUM MARKS	TOTAL DURATION	MAXIMUM TIME FOR ANSWERING		
60	80 MINUTES	70 MINUTES		

MENTION YOUR	QUESTION BOOKLET DETAILS				
CET NUMBER	VERSION CODE	SERIAL NUMBER			
	A - 1	729873			

DOs:

- 1. Check whether the CET No. has been entered and shaded in the respective circles on the OMR answer sheet.
- 2. This Question Booklet is issued to you by the invigilator after the 2nd Bell i.e., after 2.30 p.m.
- 3. The Serial Number of this question booklet should be entered on the OMR answer sheet.
- 4. The Version Code of this question booklet should be entered on the OMR answer sheet and the respective circles should also be shaded completely.
- 5. Compulsorily sign at the bottom portion of the OMR answer sheet in the space provided.

DON'TS:

- 1. THE TIMING AND MARKS PRINTED ON THE OMR ANSWER SHEET SHOULD NOT BE DAMAGED/MUTILATED/SPOILED.
- 2. The 3rd Bell rings at 2.40 p.m., till then;
 - Do not remove the paper seal present on the right hand side of this question booklet.
 - Do not look inside this question booklet.
 - Do not start answering on the OMR answer sheet.

IMPORTANT INSTRUCTIONS TO CANDIDATES

- 1. This question booklet contains 60 questions and each question will have one statement and four distracters. (Four different options / choices.)
- 2. After the 3rd Bell is rung at 2.40 p.m., remove the paper seal on the right hand side of this question booklet and check that this booklet does not have any unprinted or torn or missing pages or items etc., if so, get it replaced by a complete test booklet. Read each item and start answering on the OMR answer sheet.
- 3. During the subsequent 70 minutes:
 - Read each question carefully.
 - Choose the correct answer from out of the four available distracters (options / choices) given under each question / statement.
 - Completely darken / shade the relevant circle with a BLUE OR BLACK INK BALL POINT PEN against the question number on the OMR answer sheet.

Correct Method of shading the circle on the OMR answer sheet is as shown below:



- 4. Please note that even a minute unintended ink dot on the OMR answer sheet will also be recognised and recorded by the scanner. Therefore, avoid multiple markings of any kind on the OMR answer sheet.
- 5. Use the space provided on each page of the question booklet for Rough Work. Do not use the OMR answer sheet for the same.
- 6. After the last bell is rung at 3.50 p.m., stop writing on the OMR answer sheet and affix your LEFT HAND THUMB IMPRESSION on the OMR answer sheet as per the instructions.
- 7. Hand over the OMR ANSWER SHEET to the room invigilator as it is.
- 8. After separating the top sheet (Our Copy), the invigilator will return the bottom sheet replica (Candidate's copy) to you to carry home for self-evaluation.
- 9. Preserve the replica of the OMR answer sheet for a minimum period of ONE year.





- 1. The unit cell with crystallographic dimensions, $a \neq b \neq c$, $\alpha = \gamma = 90$ and $\beta \neq 90$ is
 - (1) Triclinic

(2) Monoclinic

(3) Orthorhombic

- (4) Tetragonal
- 2. While charging the lead storage battery,
 - (1) PbSO₄ on anode is reduced to Pb
 - (2) PbSO₄ on cathode is reduced to Pb
 - (3) PbSO₄ on cathode is oxidized to Pb
 - (4) PbSO₄ on anode is oxidized to PbO₂
- 3. Adenosine is an example of
 - (1) Nucleotide

(2) Purine base

(3) Pyrimidine base

(4) Nucleoside

- 4. Orlon has monomeric unit
 - (1) Acrolein

(2) Glycol

(3) Vinyl cyanide

- (4) Isoprene
- 5. The two electrons have the following set of quantum numbers:

$$P = 3, 2, -2, +\frac{1}{2}$$

$$Q = 3, 0, 0, +\frac{1}{2}$$

Which of the following statement is true?

- (1) P and Q have same energy
- (2) P has greater energy than Q
- (3) P has lesser energy than Q
- (4) P and Q represent same electron



- 6. H_2O_2 cannot oxidise
 - (1) PbS

(2) Na₂SO₃

(3) O

(4) KI

7. In the given set of reactions,

2-Bromopropane
$$\xrightarrow{\text{AgCN}} X \xrightarrow{\text{LiA/H}_4} Y$$

the IUPAC name of product 'Y' is

- (1) N-Methylpropanamine
- (2) N-Isopropylmethanamine

(3) Butan-2-amine

- (4) N-Methylpropan-2-amine
- 8. On heating with concentrated NaOH solution in an inert atmosphere of CO₂, white phosphorous gives a gas. Which of the following statement is <u>incorrect</u> about the gas?
 - (1) It is less basic than NH₃.
 - (2) It is more basic than NH₃.
 - (3) It is highly poisonous and has smell like rotten fish.
 - (4) It's solution in water decomposes in the presence of light.
- 9. Sodium metal crystallizes in B.C.C. lattice with edge length of 4.29 Å. The radius of sodium atom is
 - (1) 2.857 Å

(2) 1.601 Å

(3) 2.145 Å

(4) 1.857 Å

10.	0.06% (w/v) aqueous solution of urea is i	isotonic wit	th
	(1) 0.06% glucose solution	(2)	0.6% glucose solution
	(3) 0.01 M glucose solution	(4)	0.1 M glucose solution
11.	In a first order reaction, the concentration. When was it half completed?	n of the rea	actant is reduced to 12.5% in one hour.
	(1) 3 hr	(2)	20 min
	(3) 30 min	(4)	15 min

- 12. The electrolyte having maximum flocculation value for AgI/Ag⁺ sol. is
 - (1) NaCl

(2) Na₂S

(3) Na₂SO₄

- (4) Na₃PO₄
- 13. Copper is extracted from Copper pyrites by heating in a Bessemer converter. The method is based on the principle that
 - (1) Copper has more affinity for oxygen than Sulphur at high temperature.
 - (2) Iron has less affinity for oxygen than Sulphur at high temperature.
 - (3) Copper has less affinity for oxygen than Sulphur at high temperature.
 - (4) Sulphur has less affinity for oxygen at high temperature.
- 14. Which of the following will be able to show geometrical isomerism?
 - (1) MA₃B Square planar
- (2) MA_2B_2 Tetrahedral
- (3) MABCD Square planar
- (4) MABCD Tetrahedral



The electronic configuration of Gd²⁺ is (at. no. of Gd is 64)

[Xe] 4f⁸

(2) [Xe] $4f^7$

(3) [Xe] $4f^7 5d^1 6s^2$

(4) [Xe] $4f^7 5d^1$

16. $MSO_4 \xrightarrow{NH_4OH} \downarrow X \xrightarrow{NH_4OH} Y \xrightarrow{H_2S} \downarrow Z$ white

Here M and Z are

Cu, ZnS

Fe, FeS

(2) Zn, ZnS
(4) Al, Al₂S₃

The hydrolysis of optically active 2-bromobutane with aqueous NaOH result in the formation of

(1) (+) butan-2-ol

(2) (-) butan-2-ol

(3) (\pm) butan-1-ol

(±) butan-2-ol

The distinguishing test between methanoic acid and ethanoic acid is

Litmus test

Tollen's test

Esterification test (3)


Sodium bicarbonate test


19. In $H_2 - O_2$ fuel cell the reaction occurring at cathode is


- (1) $2H_{2(g)} + O_{2(g)} \longrightarrow 2H_2O_{(l)}$
- (2) $O_{2(g)} + 2H_2O_{(l)} + 4e^- \longrightarrow 4OH_{(aq)}$
- (3) $H^+ + e^- \longrightarrow \frac{1}{2} H_2$


(4) $H^{+}_{(aq)} + \overline{O}H_{(aq)} \longrightarrow H_{2}O_{(/)}$

20. Which of the following curve is in accordance with Freundlich adsorption isotherm?









21. How many ions per molecule are produced in the solution when Mohr salt is dissolved in excess of water?

(1) 4

(2)

(3) 6

(4) 10

22. Glycogen is

- (1) a polymer of β-D-glucose units
- (2) a structural polysaccharide
- (3) structurally very much similar to amylopectin
- (4) structurally similar to amylopectin but extensively branched

23. Number of possible alkynes with formula C₅H₈ is

(1) 2

(2)

(3) 4

(4) 5

24. Which of the following aqueous solution has the highest freezing point?

(1) 0.1 M Sucrose

(2) 0.01 M NaCl

(3) 0.1 M NaCl

(4) 0.01 M Na₂SO₄

25. Half life period of a first order reaction is 10 min. Starting with initial concentration 12 M, the rate after 20 min is

(1) 0.0693 M min⁻¹

- (2) $0.693 \times 3 \text{ M min}^{-1}$
- (3) $0.0693 \times 3 \text{ M min}^{-1}$
- (4) $0.0693 \times 4 \text{ M min}^{-1}$

26. The salt which responds to dilute and concentrated H₂SO₄ is

(1) CaF₂

(2) $Ba(NO_3)_2$

(3) Na₂SO₄

(4) Na₃PO₄

27. On heating potassium permanganate, one of the following compound is not obtained:

(1) O_2

(2) MnO

(3) MnO₂

(4) K_2MnO_4

28. \longrightarrow Br + Mg $\xrightarrow{\text{dry ether}}$ A $\xrightarrow{\text{H}_2\text{O}}$ B.

The product 'B' is

(1) OH

 $(2) \qquad \longrightarrow MgBr$

 $(3) \qquad \qquad \bigcirc$

 $(4) \qquad \bigcirc \longrightarrow OF$

29.	The formation of cyanohydrin from a ketone is an example of							
	(1)	Nucleophilic substitution	(2)	Nucleophilic addition				
	(3)	Electrophilic addition	(4)	Electrophilic substitution				

- onowing is an essential amino acid.
 - Tyrosine

Cysteine

Isoleucine (3)

- Serine
- The aqueous solution of following salt will have the lowest pH:
 - NaClO₃

NaClO

NaClO2

- NaClO₄
- 32. For one of the element various successive ionization enthalpies (in kJ mol⁻¹) are given below:

I.E.	1 st 2 nd		3 rd	4 th	5 th	
	577.5	1810	2750	11,580	14,820	

The element is

Si

(3.)

- Mg
- 0.30 g of an organic compound containing C, H and Oxygen on combustion yields 0.44 g CO₂ and 0.18 g H₂O. If one mol of compound weighs 60, then molecular formula of the compound is
 - CH₂O

 C_3H_8O

 C_4H_6O (3)

 $C_2H_4O_2$

- 34. One of the following amide will not undergo Hoffmann bromamide reaction:
 - (1) CH₃CONH₂
 - (2) CH₃CONHCH₃
 - (3) $C_6H_5CONH_2$
 - (4) CH₃CH₂CONH₂
- 35. Cheilosis and digestive disorders are due to the deficiency of
 - (1) Thiamine

(2) Ascorbic acid

(3) Riboflavin

- (4) Pyridoxine
- 36. How many Coulombs of electricity are required for the oxidation of one mol of water to dioxygen?
 - (1) $9.65 \times 10^4 \text{ C}$

(2) $1.93 \times 10^4 \text{ C}$

(3) $1.93 \times 10^5 \text{ C}$

- (4) $19.3 \times 10^5 \text{ C}$
- 37. 100 cm³ of 1 M CH₃COOH was mixed with 100 cm³ of 2 M CH₃OH to form an ester. The change in the initial rate if each solution is diluted with equal volume of water would be
 - (1) 2 times

(2) 4 times

(3) 0.5 times

(4) 0.25 times

Space For Rough Work

A-1

									-
38.	Which	of the	following	colloids	cannot	he	easily	coagulated	?
20.	AATITOTI	OI the	TOTTO WY TITE	Comoras	Culling		-uoii	COURTURE	

- (1) Lyophobic colloids
- (2) Multimolecular colloids
- (3) Macromolecular colloids
- (4) Irreversible colloids

39. The complex ion having minimum magnitude of Δ_0 (CFSE) is

- (1) $[Cr(CN)_6]^{3-}$
- (2) $[Co(NH_3)_6]^{3+}$
- (3) $[Co(Cl)_6]^{3-}$
- (4) $[Cr(H_2O)_6]^{3+}$

40. The arrangement of following compounds:

- i. bromomethane
- ii. bromoform
- iii. chloromethane
- iv. dibromomethane

In the increasing order of their boiling point is

(1) $iii \le i \le iv \le ii$

(2) iv < iii < i < ii

(3) ii < iii < i < iv

 $(4) \quad i < ii < iii < iv$

41. Iodoform can be prepared from all, except


(1) propan-2-ol

(2) butan-2-one

(3) propan-1-ol

(4) acetophenone

42. Identify 'Q' in the following sequence of reactions:



(1)

(2)

(3)

(4)

43. Cryolite is

- (1) Na₃A/F₆ and is used in the electrolysis of alumina for decreasing electrical conductivity.
- (2) Na₃A/F₆ and is used in the electrolysis of alumina for lowering the melting point of alumina only.
- (3) Na₃A/F₆ and is used in the electrolysis of alumina for lowering the melting point and increasing the conductivity of alumina.
- (4) Na₃A/F₆ and is used in the electrolytic refining of alumina.

44. Which of the following compound of Xenon has pyramidal geometry?

(1) XeOF₄

(2) XeF₂

(3) XeO₃

(4) XeF₄

45. After adding non-volatile solute freezing point of water decreases to -0.186 °C. Calculate ΔT_b if $K_f = 1.86$ K kg mol⁻¹ and $K_b = 0.521$ K kg mol⁻¹


(1) 0.521

(2) 0.0521

(3) 1.86

(4) 0.0186

46. Plot of Maxwell's distribution of velocities is given below:



Which of the following is correct about this plot?

(1) $T_1 < T_2$

(2) $f_1 > f_2$

 $(3) T_1 > T_2$

 $(4) \quad V_1 < V_2$

47. The pair of compound which cannot exist together in solution is

- (1) NaHCO₃ and NaOH
- (2) NaHCO₃ and H₂O
- (3) NaHCO₃ and Na₂CO₃
- (4) Na₂CO₃ and NaOH

48. What amount of dioxygen (in gram) contains 1.8×10^{22} molecules?

(1) 0.0960

(2) 0.960

(3) 9.60

(4) 96.0

- 49. Using MOT, compare O_2^+ and O_2^- species and choose the incorrect option.
 - (1) O_2^+ have higher bond order than O_2^- .
 - (2) O_2 is less stable.
 - (3) O_2^+ is diamagnetic while O_2^- is paramagnetic.
 - (4) Both O_2^+ and O_2^- are paramagnetic.
- 50. Which of the following is not true?
 - (1) Erythromycin is a bacteriostatic antibiotic.
 - (2) Ampicillin is not a natural antibiotic.
 - (3) Prontosil is not converted into sulphanilamide in the body.
 - (4) Vancomycin is a broad spectrum antibiotic.
- 51. In the reaction

$$S + \frac{3}{2}O_2 \longrightarrow SO_3 + 2x \text{ kJ and } SO_2 + \frac{1}{2}O_2 \longrightarrow SO_3 + y \text{ kJ}$$

heat of formation of SO₂ is

(1)
$$x + y$$

$$(2)$$
 $x-y$

$$(3) \quad 2x - y$$

(4)
$$2x + y$$

- 52. Arrange the following compounds in the increasing order of their acidic strength:
 - i. m-nitrophenol
- ii. m-cresol

iii. phenol

- iv. m-chlorophenol
- $(1) \quad iii < ii < i < iv$

(2) $ii \le iv \le iii \le i$

 $(3) \quad ii < iii < iv < i$

 $(4) \quad ii < iii < i < iv$

53. In the sequence of following reactions:

$$P \xrightarrow{(1) Br_{2}} Q \xrightarrow{(2) Sn/HCl} Q \xrightarrow{(2) H_{2}O/H_{3}PO_{2}} R \xrightarrow{KMnO_{4}} \overline{OH} \xrightarrow{COOH} Br$$

the starting compound 'P' is

(1) o-nitro toluene

(2) m-nitro toluene

(3) o-bromo toluene

- (4) p-nitro toluene
- 54. Acetic acid is treated with Ca(OH)₂ and the product so obtained is subjected to dry distillation. The final product is
 - (1) ethanal

(2) propanal

(3) propanone

(4) ethanol

- 55. The correct statement is
 - (1) BF₃ is the strongest Lewis acid among the other boron halides.
 - (2) Bl₃ is the weakest Lewis acid among the boron halides.
 - (3) There is maximum $p\pi p\pi$ back bonding in BF₃.
 - (4) There is minimum $p\pi p\pi$ back bonding in BF₃.
- 56. Which of the following compound possesses the "C H" bond with the lowest bond dissociation energy?
 - (1) Toluene

(2) Benzene

(3) n-pentane

(4) 2, 2-dimethyl propane

- 57. In presence of HCl, H₂S results the precipitation of Group-2 elements but not Gp-4 elements during qualitative analysis. It is due to
 - (1) higher concentration of S²⁻
- 2) higher concentration of H⁺
- (3) lower concentration of S²⁻
- (4) lower concentration of H⁺
- 58. One of the following conversion results in the change of hybridization and geometry:
 - (1) CH_4 to C_2H_6

(2) NH_3 to NH_4

(3) BF_3 to $B\overline{F}_4$

- (4) H_2O to H_3O
- 59. Water softening by Clark's process uses
 - (1) CaHCO₃

(2) NaHCO₃

(3) Na₂CO₃

- (4) Ca(OH)₂
- 60. An alkali metal hydride (NaH) reacts with diborane in 'A' to give a tetrahedral compound 'B' which is extensively used as reducing agent in organic synthesis. The compounds 'A' and 'B' respectively are
 - (1) C_2H_6 and C_2H_5Na

(2) CH_3COCH_3 and $B_3N_3H_6$

(3) C_6H_6 and NaBH₄

(4) $(C_2H_5)_2O$ and NaBH₄



