



**CHEMISTRY MARKING SCHEME**

**SET -56/2**




**Compt. July, 2015**

Qu es.	Value points	Marks
1	Emulsions are liquid – liquid colloidal systems. For example – milk, cream (or any other one correct example)	½ + ½
2	Formation of stable complex by polydentate ligand.	1
3	Propanal	1
4	p-Nitroaniline < Aniline < p-Toluidine	1
5	Frenkel defect	1
6	i) Due to high bond dissociation enthalpy of N ≡ N ii) Due to low bond dissociation enthalpy of F <sub>2</sub> than Cl <sub>2</sub> and strong bond formation between N and F	1 1
7	Potassium permanganate is prepared by fusion of MnO <sub>2</sub> with an alkali metal hydroxide and an oxidising agent like KNO <sub>3</sub> . This produces the dark green K <sub>2</sub> MnO <sub>4</sub> which disproportionates in a neutral or acidic solution to give permanganate. $2\text{MnO}_2 + 4\text{KOH} + \text{O}_2 \rightarrow 2\text{K}_2\text{MnO}_4 + 2\text{H}_2\text{O}$ $3\text{MnO}_4^{2-} + 4\text{H}^+ \rightarrow 2\text{MnO}_4^- + \text{MnO}_2 + 2\text{H}_2\text{O}$ Oxalate ion or oxalic acid is oxidised at 333 K: $5\text{C}_2\text{O}_4^{2-} + 2\text{MnO}_4^- + 16\text{H}^+ \rightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 10\text{CO}_2$ OR	1 1
7	i) <b>Iodine is liberated from potassium iodide :</b> $10\text{I}^- + 2\text{MnO}_4^- + 16\text{H}^+ \rightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 5\text{I}_2$ ii) <b>Hydrogen sulphide is oxidised, sulphur being precipitated:</b> $\text{H}_2\text{S} \rightarrow 2\text{H}^+ + \text{S}^{2-}$ $5\text{S}^{2-} + 2\text{MnO}_4^- + 16\text{H}^+ \rightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 5\text{S}$	1 1
8	 $\begin{array}{c} \text{H} \quad \text{H} \\   \quad   \\ \text{H}-\text{C}-\text{C}-\ddot{\text{O}}-\text{H} + \text{H}^+ \xrightleftharpoons{\text{Fast}} \text{H}-\text{C}-\text{C}-\overset{\text{H}}{\underset{ }{\text{O}^+}}-\text{H} \\   \quad   \\ \text{H} \quad \text{H} \end{array}$ $\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\   \quad   \quad   \\ \text{H}-\text{C}-\text{C}-\overset{\text{H}}{\underset{ }{\text{O}^+}}-\text{H} \xrightleftharpoons{\text{Slow}} \text{H}-\text{C}-\text{C}^+ + \text{H}_2\text{O} \\   \quad   \\ \text{H} \quad \text{H} \end{array}$	½ ½ 1



						
9	<p>i) <b>Mole fraction of a component =</b></p> $\frac{\text{Number of moles of the component}}{\text{Total number of moles of all the components}}$ <p>ii) Molality (<math>m</math>) is defined as the number of moles of the solute per kilogram (kg) of the solvent.</p> <p>Or</p> $\text{Molality (m)} = \frac{\text{Moles of solute}}{\text{Mass of solvent in kg}}$	1 1				
10	<p>Zero order : <math>\text{mol L}^{-1}\text{s}^{-1}</math>  Second order : <math>\text{L mol}^{-1}\text{s}^{-1}</math></p>	1 1				
11	<p>i) It lowers the melting point of alumina / acts as a solvent.</p> <p>ii)</p> <table border="1" data-bbox="368 1240 1665 1386"> <tbody> <tr> <td>Roasting</td> <td>Calcination</td> </tr> <tr> <td>Ore is heated in a regular supply of air</td> <td>Heating in a limited supply or absence of air.</td> </tr> </tbody> </table> <p>(Or with equation)</p> <p>iii) It is a process of separation of different components of a mixture which are differently adsorbed on a suitable adsorbent.</p> <p>OR</p>	Roasting	Calcination	Ore is heated in a regular supply of air	Heating in a limited supply or absence of air.	1 1 1
Roasting	Calcination					
Ore is heated in a regular supply of air	Heating in a limited supply or absence of air.					
11	<p><b><math>3\text{Fe}_2\text{O}_3 + \text{CO} \rightarrow 2\text{Fe}_3\text{O}_4 + \text{CO}_2</math></b>  <b>(Iron ore)</b></p> <p><b><math>\text{Fe}_3\text{O}_4 + \text{CO} \rightarrow 3\text{FeO} + \text{CO}_2</math></b></p> <p><b><math>\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2</math></b>  <b>(Limestone)</b></p> <p><b><math>\text{CaO} + \text{SiO}_2 \rightarrow \text{CaSiO}_3</math></b>  <b>(Slag)</b></p> <p><b><math>\text{FeO} + \text{CO} \rightarrow \text{Fe} + \text{CO}_2</math></b></p> <p><b><math>\text{C} + \text{CO}_2 \rightarrow 2\text{CO}</math></b>  <b>Coke</b></p> <p><b><math>\text{C} + \text{O}_2 \rightarrow \text{CO}_2</math></b></p> <p><b><math>\text{FeO} + \text{C} \rightarrow \text{Fe} + \text{CO}</math></b></p> <p>(any 6 correct equations)</p>	$6 \times \frac{1}{2}$ = 3				
12	<p>Disproportionation : The reaction in which an element undergoes self-oxidation and self-reduction simultaneously. For example –</p> $2\text{Cu}^+ (\text{aq}) \longrightarrow \text{Cu}^{2+} (\text{aq}) + \text{Cu}(\text{s})$ <p>(Or any other correct equation)</p>	$1 \frac{1}{2}$  $1 \frac{1}{2}$				
13	<p>i) Hexaamminecobalt(III) chloride</p> <p>ii) Tetrachlorido nickelate(II)</p> <p>iii) Potassium hexacyanoferrate(III)</p>	1 1 1				







14	i) 2-bromobutane ii) 1, 3-dibromobenzene iii) 3-choloropropene	1 1 1
15	i)  ii) $\text{CH}_3\text{CH}_2\text{MgCl} \xrightarrow[\text{H}_2\text{O}]{\text{HCHO}} \text{CH}_3\text{-CH}_2\text{-CH}_2\text{-OH}$ $\text{CH}_3\text{CH}=\text{CH}_2 + \text{H}_2\text{O} \xrightleftharpoons{\text{H}^+} \text{CH}_3\text{-}\underset{\text{OH}}{\text{CH}}\text{-CH}_3$	1 1 1
16	i) $\text{CH}_3\text{-CH}_2\text{OH} \xrightarrow{\text{PCl}_5} \text{CH}_3\text{CH}_2\text{Cl}$ ii)  iii) $\text{CH}_3\text{Cl} + \text{CH}_3\text{CH}_2\text{-ONa} \longrightarrow \text{CH}_3\text{CH}_2\text{-O-CH}_3$	1 1 1
17	i) Peptide linkage – in proteins, $\alpha$ -amino acids are connected to each other by <b>peptide bond</b> or <b>peptide linkage (-CONH- bond)</b> . ii) Primary structure - each polypeptide in a protein molecule having amino acids which are linked with each other in a specific sequence. iii) Denaturation - When a protein is subjected to physical change like change in temperature or chemical change like change in pH, protein loses its biological activity.	1 1 1
18	Copolymerisation is a polymerisation reaction in which a mixture of more than one monomeric species is allowed to polymerise and form a copolymer.   $n \text{ CH}_2=\text{CH}-\text{CH}=\text{CH}_2 + \text{Styrene} \longrightarrow \left[ \text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-\underset{\text{C}_6\text{H}_5}{\text{CH}}-\text{CH}_2 \right]_n$ <b>1, 3-Butadiene</b> <b>Styrene</b> <b>Butadiene - styrene copolymer</b>	1 1



	$n \text{ CH}_2=\text{CH}-\text{CH}=\text{CH}_2 + n \text{ CH}_2=\overset{\text{CN}}{\text{C}}\text{H} \xrightarrow{\text{Copolymerisation}} \left[ \text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-\text{CH}_2-\overset{\text{CN}}{\text{C}}\text{H} \right]_n$ <p><b>1,3-Butadiene</b>                      <b>Acrylonitrile</b>    <b>Buna-N</b> (or any other correct example)</p>	1
19	$r = \frac{\sqrt{2}a}{4}$ $r = \frac{1.414 \times 4.077 \times 10^{-8} \text{ cm}}{4}$ $r = 1.44 \times 10^{-8} \text{ cm}$	1 1 1
20	$\pi_{\text{cane sugar}} = \pi_X$ <p>Therefore, <math>c_{\text{cane sugar}} = c_X</math> (where c is molar concentration)</p> $\frac{W_{\text{cane sugar}}}{M_{\text{cane sugar}}} = \frac{W_X}{M_X}$ $\frac{5 \text{ g}}{342 \text{ g mol}^{-1}} = \frac{0.877}{M_X}$ $M_X = \frac{0.877 \times 342}{5} \text{ gmol}^{-1}$ $M_X = 59.9 \text{ or } 60 \text{ gmol}^{-1}$	1 1 1
21	$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$ $60 \text{ s}^{-1} = \frac{2.303}{t} \log \frac{[R]_0}{\frac{[R]_0}{10}}$ $t = \frac{2.303}{60 \text{ s}^{-1}} \log 10$ $t = \frac{2.303}{60} \text{ s}$ $t = 0.0384 \text{ s}$	1 1 1
22	<p>i) It is a process of removing the dissolved substance from a colloidal solution by means of diffusion through a semi - permeable membrane.</p> <p>ii) The movement of colloidal particles under an applied electric potential towards oppositely charged electrode is called electrophoresis.</p> <p>iii) Colloidal particles scatter light in all directions in space. This scattering of light illuminates the path of beam in the colloidal dispersion.</p>	1 1 1
23	<p>i) Aspartame, Saccharin (any one)</p> <p>ii) No</p> <p>iii) Social concern, empathy, concern, social awareness (any 2 )</p>	1 1 2
24	<p>a) Due to relatively stable half – filled p-orbitals of group 15 elements</p> <p>b) i) <math>\text{CaF}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{CaSO}_4 + 2\text{HF}</math></p> <p>ii) <math>\text{SO}_2(\text{g}) + \text{Cl}_2(\text{g}) \rightarrow \text{SO}_2\text{Cl}_2(\text{l})</math></p> <p>iii) <math>2\text{NH}_4\text{Cl} + \text{Ca}(\text{OH})_2 \rightarrow 2\text{NH}_3 + 2\text{H}_2\text{O} + \text{CaCl}_2</math></p> <p style="text-align: center;">OR</p>	2 1 1 1



24	<p>a) i)</p>  <p>ii)</p>  <p>b) i) Due to small size of nitrogen, the lone pair of electron on nitrogen is localized/ easily available for donation.  ii) Because they need only one electron to attain stable/noble gas configuration.</p>  <p>iii)</p>	1  1  1  1  1
25	<p>a) i)</p>  <p>ii)</p> $(\text{CH}_3)_2\text{C}=\text{CHCOCH}_3$ <p>b) i) Add <math>\text{NaHCO}_3</math>, benzoic acid will give brisk effervescence of <math>\text{CO}_2</math> whereas ethylbenzoate will not.  ii) Add <math>\text{NaOH}</math> and <math>\text{I}_2</math>, acetophenone forms yellow ppt of iodoform on heating whereas benzaldehyde will not.  iii) Add neutral <math>\text{FeCl}_3</math>, phenol gives violet colouration whereas benzoic acid does not.</p>	1  1  1  1

	(or any other correct test)	1
	OR	
25	<p>a) i)</p> $\begin{array}{c} \text{CH}_3 \\ \text{CH}_3 \end{array} \text{C}=\text{N}-\text{OH}$ <p>ii)</p> $\begin{array}{c} \text{CH}_3 \\ \text{H} \end{array} \text{C}=\text{N}-\text{NH}-\overset{\text{O}}{\parallel}{\text{C}}-\text{NH}_2$ <p>b) i)</p> $\text{CH}_3\text{CHO} \xrightarrow[\text{conc HCl}]{\text{Zn-Hg}} \text{CH}_3-\text{CH}_3$ <p>ii)</p> $2 \text{CH}_3-\text{CHO} \xrightleftharpoons{\text{dil. NaOH}} \text{CH}_3-\underset{\text{OH}}{\text{CH}}-\text{CH}_2-\text{CHO}$ <p>iii)</p> $\text{CH}_3\text{CHO} \xrightarrow{\text{LiAlH}_4} \text{CH}_3\text{CH}_2\text{OH}$	1 1 1 1 1
26	$E^0_{\text{cell}} = E^0_{\text{Sn}^{2+}/\text{Sn}} - E^0_{\text{Zn}^{2+}/\text{Zn}}$ $= -0.14\text{V} - (-0.76\text{V})$ $= 0.62\text{V}$ $\Delta_r G^0 = -n F E^0_{\text{cell}}$ $= -2 \times 96500 \text{ C mol}^{-1} \times 0.62 \text{ V}$ $= -119660 \text{ J mol}^{-1}$ $E_{\text{cell}} = E^0_{\text{cell}} - \frac{0.059}{n} \log \frac{[\text{Zn}^{2+}]}{[\text{Sn}^{2+}]}$ $E_{\text{cell}} = 0.62 - \frac{0.059}{2} \log \frac{[\text{Zn}^{2+}]}{[\text{Sn}^{2+}]}$ <p style="text-align: center;">OR</p>	1 1 1 1 1
26	<p>a) The conductivity of a solution at any given concentration is the conductance of one unit volume of solution kept between two platinum electrodes with unit area of cross section and at a distance of unit length.</p> <p>Molar conductivity of a solution at a given concentration is the conductance of the volume <math>V</math> of solution containing one mole of electrolyte kept between two electrodes with area of cross section <math>A</math> and distance of unit length.</p> <p>Molar conductivity increases with decrease in concentration.</p>	1/2 1/2 1

$\begin{aligned} \text{b) } E^{\circ}_{\text{cell}} &= E^{\circ}_{\text{C}} - E^{\circ}_{\text{A}} \\ &= 0.80\text{V} - 0.77\text{V} \\ &= 0.03\text{V} \end{aligned}$	1/2
$\begin{aligned} \Delta_r G^{\circ} &= -n F E^{\circ}_{\text{cell}} \\ &= -1 \times 96500 \text{ C mol}^{-1} \times 0.03 \text{ V} \\ &= -2895 \text{ J mol}^{-1} \end{aligned}$	1
$\text{Log } K_c = \frac{n E^{\circ}_{\text{cell}}}{0.059}$	1/2
$\text{Log } K_c = \frac{1 \times 0.03}{0.059}$	1/2
$\text{Log } K_c = 0.508$	

Dr. Sangeeta Bhatia

Sh. S.K. Munjal

Sh. D.A. Mishra

Ms. Garima Bhutani



collegedunia.com  
India's largest Student Review Platform

