

JEE-Main-29-07-2022-Shift-1 (Memory Based)

Physics

Question: Position of a particle x at time t are related as $t = \sqrt{2x+4}$. The velocity of the particle at $t = 4s$ is equal to (in S.I. units)

Options:

- (a) 4
- (b) 2
- (c) 1
- (d) 5

Answer: (a)

Solution:

$$t = \sqrt{2x+4} \Rightarrow x = \frac{1}{2}(t^2 - 4)$$

$$\Rightarrow \frac{dx}{dt} = v = t$$

$$\text{At } t = 4s, v = 4m/s$$

Question: Two rods of identical lengths and cross-sectional area are connected in series. If σ_1 and σ_2 is the thermal conductivity of material of two rods then equivalent conductivity of combination is equal to

Options:

- (a) $\frac{2\sigma_1\sigma_2}{\sigma_1 + \sigma_2}$
- (b) $\frac{\sigma_1\sigma_2}{\sigma_1 + \sigma_2}$
- (c) $\frac{\sigma_1\sigma_2}{\sigma_1 - \sigma_2}$
- (d) $\frac{2\sigma_1\sigma_2}{\sigma_1 - \sigma_2}$

Answer: (a)

Solution:



$$R_1 = \frac{L}{\sigma_1 A}, R_2 = \frac{L}{\sigma_2 A}$$

$$R_{net} = R_1 + R_2 = \frac{L}{A} \left(\frac{1}{\sigma_1} + \frac{1}{\sigma_2} \right)$$

Must be equivalent to $R_{net} = R_1 + R_2 = \frac{2L}{\sigma A}$

So, $\frac{2L}{\sigma A} = \frac{L}{A} \left(\frac{1}{\sigma_1} + \frac{1}{\sigma_2} \right)$

$$\sigma = \frac{2\sigma_1\sigma_2}{\sigma_1 + \sigma_2}$$

Question: A travelling microscope has vernier scale with 9MSD = 10 VSD. If one main scale division (MSD) is equal to 1 mm, then least count of travelling microscope is

Options:

- (a) 0.005 m
- (b) 0.002 m
- (c) 0.0001 m
- (d) 0.0005 m

Answer: (c)

Solution:

Least count, $LC = 1MSD - 1VSD$

$$\Rightarrow LC = 1MSD - \frac{9}{10}MSD$$

$$\Rightarrow LC = \frac{1}{10}MSD = \frac{1}{10} \times 0.001m$$

$$\Rightarrow LC = 0.0001m$$

Question: Find the ratio of energy of electron when it transitions from second to first energy state in comparison to highest state to first energy state of hydrogen atom

Options:

- (a) $\frac{1}{4}$
- (b) $\frac{5}{36}$
- (c) $\frac{8}{9}$
- (d) $\frac{3}{4}$

Answer: (d)


Solution:

Energy of photon is given as $E = h \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$

$$\text{So, } \frac{(hv)_{2 \rightarrow 1}}{(hv)_{\infty \rightarrow 1}} = \frac{\left(\frac{1}{1^2} - \frac{1}{2^2}\right)}{\left(\frac{1}{1^2} - \frac{1}{\infty^2}\right)} = \frac{\left(\frac{3}{4}\right)}{1}$$

Ratio = 3 : 4

Question: The value of current (in A) as shown is _____.



Options:

- (a) 2A
- (b) 3A
- (c) 4A
- (d) 5A

Answer: (a)

Solution:

All the resistance are in parallel.

$$\Rightarrow R_{net} = 3\Omega$$

$$\Rightarrow i = \frac{V}{R_{net}} = \frac{6}{3} = 2A$$

Question: Find the value of electric field at depletion layer in p-n junction if width is $6 \times 10^{-6} m$ and potential difference is 0.6 V, is _____ $\times 10^5 V / m$

Options:

- (a) $2 \times 10^{-5} V/m$
- (b) $6 \times 10^{-6} V/m$
- (c) $1 \times 10^5 V/m$
- (d) $3 \times 10^6 V/m$

Answer: (c)

Solution:

$$AV = E.D$$

$$E = \frac{(0.6)}{6 \times 10^{-6}}$$

$$E = 1 \times 10^5 V / m$$

Question: A projectile with kinetic energy E at point of projection is projected at angle 45° . Its kinetic energy at top most point is equal to

Options:

- (a) $\frac{E}{2}$
- (b) $\frac{3E}{2}$
- (c) $\frac{E}{4}$
- (d) $\frac{E}{3}$

Answer: (a)

Solution:

$$\Rightarrow K.E_i = \frac{1}{2}mv^2 = E$$

$$\text{Speed at highest point } v'; v \cos 45^\circ = \frac{v}{\sqrt{2}}$$

$$\Rightarrow K.E_f = \frac{1}{2}mv'^2 = \frac{1}{4}mv^2$$

$$K.E_f = \frac{E}{2}$$

Question: A particle thrown at angle 45° with horizontal with speed u has its range equal to R . At what angle should it be thrown with same speed for its range to be half of its initial value.

Options:

- (a) 60°
- (b) 30°
- (c) 15°
- (d) 70°

Answer: (c)

Solution:

$$\Rightarrow R = \frac{u^2 \sin(2 \times 45^\circ)}{g} = \frac{u^2}{g}$$

For range $\frac{R}{2}$

$$\Rightarrow \frac{u^2}{2g} = \frac{u^2 \sin 2\theta}{g}$$

$$\sin 2\theta = \frac{1}{2}$$

$$\Rightarrow \theta = 15^\circ$$

Question: A cart is moving down a smooth incline of inclination α . What is the time period of a bob hanging from the roof of the cart with a light string?

Options:

(a) $2\pi\sqrt{\frac{l}{g\cos\alpha}}$

(b) $2\pi\sqrt{\frac{l}{g}}$

(c) $2\pi\sqrt{\frac{l}{g\sin\alpha}}$

(d) $2\pi\sqrt{\frac{l}{g\cot\alpha}}$

Answer: (a)

Solution:

$$g_{\text{eff}} = g \cos \alpha$$

$$T = 2\pi\sqrt{\frac{l}{g \cos \alpha}}$$

Question: If one mole of monoatomic gas and three moles of diatomic gas are mixed, then the molar heat at constant volume is $\alpha^2 R/4$. The value of α is -----

Options:

(a) 2

(b) 3

(c) 5

(d) 1

Answer: (b)

Solution:

$$C_{V_{\text{mix}}} = \frac{(n_1 C_{V_1} + n_2 C_{V_2})}{n_1 + n_2}$$

$$C_{V_{\text{mix}}} = \frac{\left(1 \times \frac{3}{2}R + 3 \times \frac{5}{2}R\right)}{1+3}$$

$$C_{V_{\text{mix}}} = \frac{9}{4}R \quad \text{So, } \alpha = 3$$

Question: A wire of length 314 cm is made into a circular coil. Find its magnetic moment (in Am^2 if $I = 14 \text{ A}$. ($\pi = 3.14$))

Options:

(a) 10 Am^2

(b) 8 Am^2

(c) 6 Am^2

(d) 11 Am^2

Answer: (d)

Solution:

$$\mu = i\pi r^2$$

$$\mu = i\pi \left(\frac{l}{2\pi} \right)^2$$

$$\mu = 14 \times \pi \left(\frac{3.14}{2 \times 3.14} \right)^2$$

$$\mu = 11 \text{ Am}^2$$

Question: Assertion: Potential is constant on surface & inside of conductor.

Reason: E is perpendicular to surface of conductor.

Options:

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
- (b) If both assertion and reason are true, but the reason is not the correct explanation of the assertion.
- (c) If assertion is true, but reason is false.
- (d) If both the assertion and reason are false.

Answer: (a)

Solution:

Since $E=0$, therefore the potential V inside the surface is constant. Because there is no potential difference between any two points inside the conductor, the electrostatic potential is constant throughout the volume of the conductor.