



collegedunia.com
India's largest Student Review Platform

Marking scheme – 2020

CHEMISTRY (043) / CLASS XII



56/C/3

| Q.No | Expected Answer / Value Points | Marks |
|------------------|---|-------|
| SECTION A | | |
| 1 | PHBV / Nylon-2-nylon-6 / any natural polymer (or any other suitable example). | 1 |
| 2 | Homopolymer | 1 |
| 3 | Starch/ cellulose/ proteins / nucleic acids / natural rubber (or any other suitable example). | 1 |




*These answers are meant to be used by evaluators







collegedunia.com
India's largest Student Review Platform

| | | |
|------------------|--|---|
| 4 |  <p style="text-align: center;">Neoprene</p> | 1 |
| 5 | Teflon / PTFE | 1 |
| 6 | Molar conductivity increases. | 1 |
| 7 | Nucleoside | 1 |
| 8 | Kraft Temperature | 1 |
| 9 | Alitame | 1 |
| 10 | Cheese/Jellies (or any other suitable example) | 1 |
| 11 | (C) or (D) | 1 |
| 12 | (D) | 1 |
| 13 | (D) | 1 |
| 14 | (A) | 1 |
| 15 | (B) | 1 |
| 16 | (i) | 1 |
| 17 | (iii) | 1 |
| 18 | (i) | 1 |
| 19 | (i) | 1 |
| 20 | (iii) | 1 |
| SECTION B | | |
| 21 | <p>Vitamins are certain organic compounds, required in small amounts in our diet but their deficiency causes specific diseases / organic compounds required in the diet in small amounts to perform specific biological functions for normal maintenance of optimum growth and health of the organism.</p> <p>Vitamins are classified into two groups depending upon their solubility in water or fat.</p> <p>(i) Fat soluble vitamins (ii) Water soluble vitamins.</p> <p style="text-align: center;">OR</p> <p>Proteins are polymers of α-amino acids. (or any other correct answer)</p> <p>They are classified as Fibrous and Globular proteins on the basis of their shape.</p> | <p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> |
| 22. |  <p style="text-align: center;">Cis Trans</p> | 1 + 1 |
| 23 | <p>It is the rate of reaction when concentration of each reactant is taken as unity. / It is the proportionality constant in the rate law expression or in differential rate equation or in the rate of reaction.</p> <p>$K = 0.693 / t_{1/2}$</p> | <p>1</p> <p>1</p> |
| 24. | <p>a) At Anode: $2\text{H}_2\text{O}_{(l)} \longrightarrow 4\text{H}^+_{(aq)} + 4\text{e}^- + \text{O}_2$</p> <p>At Cathode: $\text{Cu}^{2+}_{(aq)} + 2\text{e}^- \longrightarrow \text{Cu}_{(s)}$. / Copper is deposited at cathode and Oxygen gas is liberated at anode.</p> <p>b) At Anode: $2\text{H}_2\text{O}_{(l)} \longrightarrow 4\text{H}^+_{(aq)} + 4\text{e}^- + \text{O}_2$</p> <p>At Cathode: $\text{Ag}^+_{(aq)} + \text{e}^- \longrightarrow \text{Ag}_{(s)}$. / Silver is deposited at cathode and oxygen gas is liberated at anode.</p> <p style="text-align: center;">OR</p> <p>$\text{Fe}^{3+} + \text{e}^- \longrightarrow \text{Fe}^{2+}$, so 1 mol of Fe^{3+} requires 1 F</p> <p>3 moles of Fe^{3+} require 3 F</p> <p>$Q = I \times t$</p> <p>$t = 3 \times 96500 / 2$</p> <p>$t = 144750 \text{ sec}$</p> | <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> |



| | | |
|------------------|--|---|
| 25. | a) Electrolytic refining -The more basic metal remains in the solution and the less basic ones go to the anode mud. / Anode is impure metal and pure metal strip is cathode while aqueous solution of the metal salt acts as the electrolyte. b) Zone refining - Impurities are more soluble in the melt than in the solid state of the metal. | 1 1 |
| 26. | $\text{Cr}^{3+} = 3$ unpaired electrons $\text{V}^{3+} = 2$ unpaired electrons Cr^{3+} is more stable due to half filled t_{2g}^3 configuration | $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ |
| 27. | Antiseptics the chemicals which either kill or prevent the growth of microorganisms but are applied to the living tissues such as wounds, cuts, ulcers and diseased skin surfaces. Examples are furacine, soframycin, etc. Disinfectants are also the chemicals which either kill or prevent the growth of microorganisms but applied to inanimate objects such as floors, drainage system, instruments, etc. e.g. Concentrated acids, Phenol (above 1% conc.) (OR any other suitable point of difference and example) | 1 1 |
| SECTION C | | |
| 28 | $K = \frac{2.303}{t} \log \frac{[R_0]}{[R]}$ $4.9 \times 10^{-3} = \frac{2.303}{t} \log \frac{4}{3}$ $t = \frac{2.303}{4.9 \times 10^{-3}} \log (0.6020 - 0.4771)$ $= 58.7 \text{ sec}$ <p style="text-align: center;">OR</p> g) Molecularity = 2 or bimolecular. h) Order = 1 or pseudo first order. i) $\text{Rate} = \frac{-\Delta[\text{C}_{12}\text{H}_{22}\text{O}_{11}]}{\Delta t} = + \frac{\Delta[\text{C}_6\text{H}_{12}\text{O}_6]}{\Delta t} = + \frac{\Delta[\text{C}_6\text{H}_{12}\text{O}_6]}{\Delta t}$ | $\frac{1}{2}$ 1 1 $\frac{1}{2}$ 1 1 1 |
| 29 | g) Tyndall effect / scattering of light / path of the light gets illuminated. h) Coagulation / precipitation / artificial rain. i) Demulsification / separation of fat from the milk / coagulation. | 1 1 1 |
| 30 | $\Lambda_m = \frac{k}{c} \times 1000$ $= \frac{8 \times 10^{-5}}{0.002} \times 1000$ $= 40 \text{ Scm}^2 \text{ mol}^{-1}$ $\alpha = \frac{\Lambda_m^c}{\Lambda_m^o}$ $= 40/390.5 = 0.102$ | $\frac{1}{2}$ 1 $\frac{1}{2}$ 1 |
| 31 | a)  b) $\text{CH}_3\text{CH}_2\text{OH}$ c)  <p style="text-align: center;">OR</p> a) i) $\text{H}_3\text{C}-\text{CH}=\text{CH}_2 \xrightarrow{\text{H}_2\text{O}, \text{H}^+} \text{H}_3\text{C}-\underset{\text{OH}}{\text{CH}}-\text{CH}_3$ ii)  b) 4-Methylphenol < phenol < 4-Nitrophenol / 4-Methylphenol , phenol , 4-Nitrophenol | 1 x 3 1 1 1 |

| | | | |
|--|---|-----|-------|
| 37 | <p>a) i)</p>  <p>ii)</p>  | 1+1 | |
| 37 | <p>b) i)</p>  <p>ii)</p>  | 1+1 | |
| <p>c) Carboxylate ion is more stabilised than phenoxide ion / conjugate base of carboxylic acid is more stable than that of phenol / carboxylate ion has two equivalent resonating structures while the structures are non equivalent in phenoxide ion/ negative charge in carboxylate ion is delocalised over more electronegative two Oxygen atoms while in phenoxide ion negative charge is delocalised over one Oxygen atom and less electronegative Phenyl ring (or C atoms). /Carboxylic acid reacts with NaHCO₃ to give brisk effervescence of CO₂ while phenol doesn't or reaction given by the student.</p> | | | 1 |
| <p>a) $\text{CH}_3\text{COOCH}_2\text{CH}_3 \xrightarrow{\text{H}_2\text{SO}_4 \text{ dil}} \text{CH}_3\text{COOH} + \text{CH}_3\text{CH}_2\text{OH}$ A B C</p> <p style="text-align: center;">OR</p> <p>$\text{CH}_3\text{CH}_2\text{OH} \xrightarrow{\text{oxidation}} \text{CH}_3\text{COOH}$</p> <p>$\text{CH}_3\text{CH}_2\text{OH} \xrightarrow{\text{dehydration}} \text{CH}_2=\text{CH}_2$</p> | | | ½ x 3 |
| <p>A = Ethyl acetate (CH₃COOCH₂CH₃), B = Acetic or Ethanoic acid (CH₃COOH), C = Ethanol (CH₃CH₂OH)</p> | | | ½ x 3 |
| <p>b) i) $\text{CH}_3\text{CHO} \xrightarrow{\text{LiAlH}_4, \text{ dry ether}} \text{CH}_3\text{CH}_2\text{OH}$</p> | | | 1 |
| <p>ii) $\text{CH}_3\text{CHO} \xrightarrow{\text{Zn/Hg in HCl}} \text{CH}_3\text{CH}_3$ Or NH₂NH₂</p> | | | 1 |
| <p>(or any other suitable reagent for the above reactions or any other correct method for conversion)</p> | | | |