DU MSc Chemistry

Topic:- DU_J18_MSC_CHEM

1) The halogen having metallic character is [Question ID = 790]

- 1. Bromine [Option ID = 3159]
- 2. Chlorine [Option ID = 3158]
- 3. Iodine [Option ID = 3160]
- 4. Fluorine [Option ID = 3157]

Correct Answer :-

• Iodine [Option ID = 3160]

2) If the density of air is 1.2 g/lit, what is the volume occupied by 7.8g of air? [Question ID = 860]

- 1. 10.10lit [Option ID = 3440]
- 2. 10 lit [Option ID = 3437]
- 6 lit [Option ID = 3438]
- 4. 6.5 lit [Option ID = 3439]

Correct Answer :-

6.5 lit [Option ID = 3439]

3) Which of the following statement/s is/are true?

[Question ID = 855]

- 1. All of these [Option ID = 3420]
- 2. Adsorption increases with increase in pressure [Option ID = 3419]
- 3. Adsorption decreases with increase in temperature [Option ID = 3418]
- 4. Adsorption is an exothermic process [Option ID = 3417]

Correct Answer :-

All of these [Option ID = 3420]

4) Which of the following species represent the example of dsp² hybridization?

[Question ID = 53390]

- 1. $[FeF_6]^{3-}$ [Option ID = 93549]
- 2. $[Fe(CN)_6]^{3-}$ [Option ID = 93546]
- 3. $[Ni(CN)_4]^{2}$ [Option ID = 93547]
- 4. $[Zn(NH_3)_4]^{2+}$ [Option ID = 93548]

Correct Answer :-

• $[Ni(CN)_4]^{2}$ [Option ID = 93547]

5) Correct characteristics of the functional groups of adenine in DNA base pair are

[Question ID = 824]

- 1. Both N(3) and C(6)NH₂ are hydrogen bond acceptors [Option ID = 3295]
- 2. Both N(3) and C(6)NH₂ are hydrogen bond acceptors. [Option ID = 3296]
- 3. N(3) is a hydrogen bond acceptor and C(6)NH₂ is a hydrogen bond donor. [Option ID = 3293]
- 4. N(1) is a hydrogen bond acceptor and C(6)NH₂ is a hydrogen bond donor. [Option ID = 3294]

Correct Answer :-

• N(1) is a hydrogen bond acceptor and C(6)NH₂ is a hydrogen bond donor. [Option ID = 3294]

6) Chemical potential is also known as [Question ID = 858]



- 1. Partial molar entropy [Option ID = 3429]
- 2. Partial molar Gibbs free energy [Option ID = 3431]
- 3. None of these [Option ID = 3432]
- 4. Partial molar enthalpy [Option ID = 3430]

• Partial molar Gibbs free energy [Option ID = 3431]

7) From the following, which is more covalent? [Question ID = 781]

1.
$$Al_2S_3$$
 [Option ID = 3122]

$$\begin{array}{c} Al_2Cl_6 \\ 3. \end{array}$$
 [Option ID = 3124]

4.
$$Al_2O_3$$
 [Option ID = 3121]

Correct Answer :-

•
$$Al_2S_3$$
 [Option ID = 3122]

8) The most probable candidate to form an octahedral complex is [Question ID = 792]

$$d^{10}$$
 [Option ID = 3168]

3.
$$d^8$$
 (low spin) [Option ID = 3166]

4.
$$d^1$$
 (low spin) [Option ID = 3165]

Correct Answer :-

$$d^{1}$$
 (low spin) [Option ID = 3165]

9) Percentage of gold in 18 carat gold is [Question ID = 787]

- 1. 18 [Option ID = 3145]
- 2. 100 [Option ID = 3148]
- 3. 75 [Option ID = 3146]
- 4. 83.6 [Option ID = 3147]

Correct Answer :-

• 75 [Option ID = 3146]

10) Which pair from the following behaves as metalloid? [Question ID = 789]

- 1. Al and Zn [Option ID = 3155]
- 2. Rb and Cs [Option ID = 3156]
- 3. Br and I [Option ID = 3153]
- 4. Pt and I [Option ID = 3154]

Correct Answer :-

• Al and Zn [Option ID = 3155]

11) For a substitution reaction following a dissociative mechanism, the rate determining step is [Question ID = 800]

- 1. dependent on the solvent concentration [Option ID = 3199]
- 2. dependent on the leaving group [Option ID = 3198]
- 3. dependent on the entering group [Option ID = 3197]
- 4. dependent on the nature of the complex [Option ID = 3200]

Correct Answer :-

• dependent on the leaving group [Option ID = 3198]

12) The amino acid constituents of artificial sweetener given below are: [Question ID = 826]



- 1. L-Aspartic acid and L-tyrosine [Option ID = 3304]
- 2. D-Glutamic acid and L-phenylgylcine [Option ID = 3301]
- 3. L-Aspartic acid and L-phenylalanine [Option ID = 3303]
- 4. L-Glutamic acid and L-phenylgylcine [Option ID = 3302]

• L-Aspartic acid and L-phenylalanine [Option ID = 3303]

13) In the following statements, which one is incorrect? [Question ID = 778]

Atomic radius of Zr and Hf are same because of lanthanide contraction

[Option ID = 3112]

- La(OH)₃ is less basic than $Lu(OH)_3$ [Option ID = 3109]
- 3. La is actually an element of transition series rather than lanthanides [Option ID = 3111]
- 4. In lanthanide series, ionic radius of Lu³⁺ ion decreases [Option ID = 3110]

Correct Answer :-

. La is actually an element of transition series rather than lanthanides [Option ID = 3111]

14) In the dichromate dianion [Question ID = 791]

- 1. 3 Cr-O bonds are equivalent [Option ID = 3163]
- 2. 6 Cr-O bonds are equivalent [Option ID = 3162]
- 3. All the Cr-O bonds are non-equivalent [Option ID = 3164]
- 4. 4 Cr-O bonds are equivalent [Option ID = 3161]

Correct Answer :-

• 6 Cr-O bonds are equivalent [Option ID = 3162]

15) Vacuum is a measure of [Question ID = 804]

- 1. Leaking rate of air [Option ID = 3214]
- 2. Leaking rate of oil [Option ID = 3216]
- 3. Leaking rate of moisture [Option ID = 3215]
- 4. Emptiness [Option ID = 3213]

Correct Answer :-

· Leaking rate of air [Option ID = 3214]

16) The Pre-exponential factor 'A' in the Arrhenius Equation depends on which of the following? [Question ID = 852]

- 1. Collision Frequency [Option ID = 3407]
- 2. Gibb's free energy of reaction [Option ID = 3406]
- 3. None of these [Option ID = 3408]
- 4. Energy of activation of the reaction [Option ID = 3405]

Correct Answer :-

• Collision Frequency [Option ID = 3407]

17) The process of heating the concentrated ore in a limited supply of air or in the absence of air is known as: [Question ID = 869]

- 1. Roasting [Option ID = 3473]
- 2. Calcination [Option ID = 3475]
- 3. Cupellation [Option ID = 3476]
- 4. Leaching [Option ID = 3474]

Correct Answer :-

• Calcination [Option ID = 3475]

18) Spectroscopic transitions leading to rotation of molecules will appear at which region of the electromagnetic spectrum? [Question ID = 866]

- 1. Ultraviolet [Option ID = 3461]
- 2. Radiofrequency [Option ID = 3464]
- 3. Infra-red [Option ID = 3463]
- 4. Microwave [Option ID = 3462]



Correct Answer :-Microwave [Option ID = 3462] 19) The ground state of a harmonic oscillator has number of nodes: [Question ID = 846] 1. 2 [Option ID = 3382] 2. 0 [Option ID = 3384] 3. 1 [Option ID = 3383] 4. 3 [Option ID = 3381] Correct Answer :-• 0 [Option ID = 3384] 20) Tritium is a radioisotope of hydrogen, it undergoes disintegration to give [Question ID = 786] α-particles [Option ID = 3142] _{2.} β-particles [Option ID = 3143] Neutrons [Option ID = 3144] 4. X-rays [Option ID = 3141] Correct Answer :-Neutrons [Option ID = 3144] 21) Which transitions are studied by UV spectrometer? [Question ID = 870] 1. Electronic [Option ID = 3478] 2. Vibrational [Option ID = 3480] 3. Nuclear [Option ID = 3479] 4. Rotational [Option ID = 3477] Correct Answer :- Electronic [Option ID = 3478] 22) What happens during digestion of a precipitate? [Question ID = 801] 1. Coalescence of smaller crystallites [Option ID = 3203] 2. Recrystallization takes place [Option ID = 3202] 3. Completion of precipitation [Option ID = 3201] 4. rate of the reaction increases [Option ID = 3204] Correct Answer :- Coalescence of smaller crystallites [Option ID = 3203] CaO, K_2O [Option ID = 3151] ${}_{2}. \ Fe_2O_3, \ ZnO \ \ [Option ID = 3150]$ PbO, Fe_3O_4 [Option ID = 3152] Correct Answer :-CaO, K_2O

24) In which of the following reaction migration of alkyl group from carbon to oxygen is observed? [Question ID = 813]



- 1. Pinacol-pinacolone rearrangement [Option ID = 3249]
- 2. Prepration of phenol from cumene hydroperoxide [Option ID = 3251]
- 3. Baeyer-villiger oxidation [Option ID = 3250]
- 4. Both Baeyer-villiger oxidation and Prepration of phenol from cumene hydroperoxide [Option ID = 3252]

Baeyer-villiger oxidation [Option ID = 3250]

25) Alkali metals form highly stable complexes with [Question ID = 795]

- 1. diethyl ether [Option ID = 3178]
- 2. Butadiene [Option ID = 3180]
- 3. Cryptand-222 [Option ID = 3177]
- 4. Cyclopentadiene [Option ID = 3179]

Correct Answer :-

• Cryptand-222 [Option ID = 3177]

26) The unit of rate constant for a second order reaction is: [Question ID = 850]

4.
$$\text{mol}^{-1} \text{ dm}^3 \text{ s}^{-1}$$
 [Option ID = 3399]

Correct Answer :-

•
$$\text{mol}^{-1} \text{ dm}^3 \text{ s}^{-1}$$
 [Option ID = 3399]

27) What is the unit of specific resistance (or resistivity) of a conductor? [Question ID = 868]

Correct Answer :-

Siemens⁻¹ cm [Option ID = 3469]

28) When a nucleophile encounters a ketone the site of attack is: [Question ID = 811]

- both the carbon and oxygen atoms, with equal probability [Option ID = 3243]
- 2. the carbon atom of the carbonyl [Option ID = 3241]
- 3. the oxygen atom of the carbonyl [Option ID = 3242]
- 4. no attack occur as ketones do not react with nucleophiles [Option ID = 3244]

Correct Answer :-

the carbon atom of the carbonyl [Option ID = 3241]

29) In the cases of gases adsorbing on solid, which of the following statement/s is/are true?

[Question ID = 845]

- 1. Decrease in temperature of the system results in increase in adsorption [Option ID = 3378]
- 2. Decrease in pressure of the system results in decrease in adsorption [Option ID = 3379]
- 3. All of these [Option ID = 3380]
- 4. Adsorption is an exothermic process [Option ID = 3377]

Correct Answer :-

• All of these [Option ID = 3380]



30) During a disproportionation reaction, [Question ID = 802]

- Simultaneous oxidation and reduction of metal ion takes place [Option ID = 3207]
- 2. Metal ion goes to lower oxidation state [Option ID = 3205]
- 3. Metal ion goes to higher oxidation state [Option ID = 3206]
- 4. Metal ion remains unchanged in its oxidation state [Option ID = 3208]

Correct Answer :-

• Simultaneous oxidation and reduction of metal ion takes place [Option ID = 3207]

31) The number of independent modes of vibration in a linear molecule having N atoms is [Question ID = 851]

- 1. 3N 6 [Option ID = 3402]
- 2. 3N 3 [Option ID = 3404]
- 3. 3N [Option ID = 3403]
- 4. 3N 5 [Option ID = 3401]

Correct Answer :-

3N - 5 [Option ID = 3401]

32) A system that maintains a constant volume is known as [Question ID = 857]

- 1. None of these [Option ID = 3428]
- 2. Isochoric system [Option ID = 3425]
- Adiabatic system [Option ID = 3427]
- 4. Isotactic system [Option ID = 3426]

Correct Answer :-

• Isochoric system [Option ID = 3425]

33) Cobalt is present in [Question ID = 777]

- 1. Vitamin B₂ [Option ID = 3106]
- Vitamin B_1 [Option ID = 3105]
- 3. Vitamin B_6 [Option ID = 3107]
- Vitamin B₁₂

f. [Option ID = 3108]

Correct Answer :-

Vitamin B₁₂

[Option ID = 3108]

34) In collision theory of bimolecular gaseous reactions, the Collision Frequency does not depend on: [Question ID = 849]

- 1. Pressure of the system [Option ID = 3394]
- 2. Number of molecules of each gas [Option ID = 3396]
- 3. Temperature of the system [Option ID = 3393]
- 4. Reduced mass of the system [Option ID = 3395]

Correct Answer :-

• Pressure of the system [Option ID = 3394]

35) An inorganic mixture dissolves in hot conc. HCl giving a blue colored solution which on addition of water becomes pink. The mixture contains [Question ID = 793]

1.
$$Fe^{3+}$$
 [Option ID = 3172]

2.
$$\frac{\text{Cr}^{3+}}{\text{[Option ID = 3171]}}$$

$$Co^{2+}$$
 [Option ID = 3170]



Co²⁺ [Option ID = 3170]

36) The Bragg's equation for crystallography can be written as: [Question ID = 844]

$$n\lambda = (2d / \sin \theta)$$
 [Option ID = 3375]
 $2 \cdot n\lambda = (2d \sin \theta)$ [Option ID = 3373]
 $2 \cdot n\lambda = (2/d) \sin 2 \theta$ [Option ID = 3374]

 $n\lambda = 1 / (2d \sin \theta)$ [Option ID = 3376]

Correct Answer :-

 $n\lambda = (2d \sin \theta)$ [Option ID = 3373]

The product X in the flowing reaction $6LiH + 8BF_3 \rightarrow 6LiBF_4 + X$ is

[Question ID = 783]

$$\begin{array}{ccc} B_4H_{10} & \text{[Option ID = 3129]} \\ B_2H_6 & \text{[Option ID = 3130]} \\ B_3H_8 & \text{[Option ID = 3132]} \\ A. & BH_3 & \text{[Option ID = 3131]} \end{array}$$

Correct Answer :-

• B_2H_6 [Option ID = 3130]

38) The product obtained in the following conversion is:

[Question ID = 840]






[Option ID = 3358]

39)

A compound with molecular formula C₄H₆O₂ shows band at 1770 cm-1 in IR spectra and peaks at 178,68,28,22 ppm in ¹³ C NMR spectrum. The correct structure of the compound is



(ii) 0 = c

(iii) O = C



[Question ID = 823]

- 1. ii [Option ID = 3290]
- 2. iv [Option ID = 3292]
- 3. iii [Option ID = 3291]
- 4. i [Option ID = 3289]

Correct Answer :-

- ii [Option ID = 3290]
- iii [Option ID = 3291]

40)

The product in the given reaction is.

OH
$$OH$$

$$OH$$

$$H_2SO_4$$
The product obtained is:

[Question ID = 810]

1.

[Option ID = 3239]

2. OH

[Option ID = 3237]

None of these

3. [Option ID = 3240]

. [Option ID = 3238]



[Option ID = 3239]

[Option ID = 3237]

Rank the following alkenes on order of increasing maximum wavelength

[Question ID = 815]

$$ii < iii < i$$
 [Option ID = 3257]

Correct Answer :-

42) The correct relation between the following compounds is

[Question ID = 5490]

- 1. enantiomers [Option ID = 21957]
- 2. homomers (identical) [Option ID = 21959]
- 3. constitutional isomers [Option ID = 21960]
- 4. diastereomers [Option ID = 21958]

Correct Answer :-

• homomers (identical) [Option ID = 21959]

Tl⁺ compounds are poisonous because

[Question ID = 806]

- 1. Stop blood circulation [Option ID = 3223]
- 2. They attack liver [Option ID = 3224]
- 3. Cut-off breathing capability [Option ID = 3222]
- 4. They can cause blood infection [Option ID = 3221]



• Cut-off breathing capability [Option ID = 3222]

44)

The major product formed in the following reaction is.

[Question ID = 839]

1. [Option ID = 3354]

2. [Option ID = 3356]

3. [Option ID = 3353]

4. [Option ID = 3355]

Correct Answer :-

• [Option ID = 3353]

⁴⁵⁾ Following reaction goes through?

[Question ID = 822]

- 1. carbene intermediate [Option ID = 3288]
- 2. free reaical intermediate [Option ID = 3285]
- 3. carbocation intermediate [Option ID = 3287]
- 4. carbanion intermediate [Option ID = 3286]

Correct Answer :-

• free reaical intermediate [Option ID = 3285]

46)

Consider an electrochemical reaction: Oxidized form $+ ne^- =$ reduced form. If an ion forms a complex with the oxidized form, then the following happens:

[Question ID = 843]



- 1. The reduction potential of the system is increased [Option ID = 3370]
- 2. The reduction potential of the system remains the same [Option ID = 3369]
- 3. The effective concentration of the reduced form is increased [Option ID = 3372]
- The reduction potential of the system is lowered [Option ID = 3371]

- The reduction potential of the system is lowered [Option ID = 3371]
- Total orbital angular momentum of np⁶ electronic system is (a.u.):

[Question ID = 864]

- 1. Option ID = 3453]
 - 1/2
 - [Option ID = 3456]
- 2 [Option ID = 3455]
- 4. [Option ID = 3454]

Correct Answer :-

• [Option ID = 3453]

48)

Identify the enantiomers among the following compounds.

[Question ID = 827]

- 1. C and D [Option ID = 3308]
- 2. B and D [Option ID = 3307]
- 3. A and C [Option ID = 3306]
- 4. A and B [Option ID = 3305]

Correct Answer :-

• C and D [Option ID = 3308]

49)



Match the following

List – 1		List – 2	
A	Phosphorescence	1	A schematic representation of the various types of <u>radiative</u> and non- <u>radiative</u> transitions that can occur in molecules
В	Intersystem Crossing	2	Spontaneous emission of radiation arising from transitions between energy states of same multiplicity
C	Jablonski Diagram	3	Non-radiative transitions between energy states of different multiplicity
D	Fluorescence	4	Spontaneous emission of radiation arising from transitions between energy states of different multiplicities

[Question ID = 841]

- 1. A4, B3, C1, D2 [Option ID = 3363]
- 2. A4, B3, C2, D1 [Option ID = 3362]
- 3. A3, B1, C2, D4 [Option ID = 3364]
- 4. A1, B2, C3, D4 [Option ID = 3361]

Correct Answer :-

• A4, B3, C1, D2 [Option ID = 3363]

The oxidation state of oxygen in O_2F_2 is

[Question ID = 794]

- 1. +2 [Option ID = 3174]
- 2. +1 [Option ID = 3173]
- 3. +4 [Option ID = 3175]
- 4. -2 [Option ID = 3176]

Correct Answer :-

• +1 [Option ID = 3173]

51)

The following molecule has

[Question ID = 829]

- 1. R configuration [Option ID = 3314]
- 2. centre of symmetry [Option ID = 3316]
- 3. S configuration [Option ID = 3315]
- 4. plane of symmetry [Option ID = 3313]

Correct Answer :-

• R configuration [Option ID = 3314]



In low chloride ion concentration, the anticancer drug <u>cis-platin</u> hydrolysis to give a <u>diaqua</u> complex and this binds to DNA via <u>adajacent</u> guanine.

The coordinating atom of guanine to Pt(II) is

[Question ID = 825]


- 1. N9 [Option ID = 3300]
- 2. N7 [Option ID = 3299]
- 3. N1 [Option ID = 3297]
- 4. N3 [Option ID = 3298]

Correct Answer :-

• N7 [Option ID = 3299]

53)

Which of the following species is aromatic in nature?









[Question ID = 818]

- 1. iv [Option ID = 3272]
- 2. i [Option ID = 3269]
- 3. ii [Option ID = 3270]
- 4. iii [Option ID = 3271]

Correct Answer :-

- iv [Option ID = 3272]
- Arrange the following in decreasing order of their acidity

[Question ID = 807]



i>ji>jjj>jv

[Option ID = 3225]

55)

The major product formed in the following reaction sequence is:

1) Li, liq. NH₃, t-BuOH

[Question ID = 5543]

1. [Option ID = 22165]

2.

[Option ID = 22163]

3.

[Option ID = 22164]

[Option ID = 22166]

Correct Answer :-

• [Ontion ID =

56)

The compound that gives precipitate on warming with aqueous AgNO3 is.

[Question ID = 834]

.....

[Option ID = 3336]

2.

[Option ID = 3333]



Br

3. [Option ID = 3335]

Br

4. [Option ID = 3334]

Correct Answer :-

Br

[Option ID = 3335]

What is the specific resistance (or resistivity) of a conductor with cross-sectional area 4 cm², length 2cm and resistance 8 ohms?

[Question ID = 853]

64 Siemens⁻¹cm [Option ID = 3411]

2. 16 Siemens⁻¹cm [Option ID = 3412]

4 Siemens⁻¹cm [Option ID = 3409]

4. Siemens⁻¹cm [Option ID = 3410]

Correct Answer :-

. 16 Siemens⁻¹cm [Option ID = 3412]

Which pair of ions cannot be precipitated by H₂S in dilute HCl?

[Question ID = 782]

 ${}_{1}.\,Al^{3+},\,Ni^{2+}\,$ [Option ID = 3127]

Bi³⁺, Sn⁴⁺ [Option ID = 3125]

3. Ni^{2+} , Cu^{2+} [Option ID = 3128]

4. Zn^{2+} , Cu^{2+} [Option ID = 3126]

Correct Answer :-

• Al^{3+} , Ni^{2+} [Option ID = 3127]

59)



Which of the following bromides is the major product of the reaction shown below, assuming that there are no carbocation rearrangement?

[Question ID = 816]

Correct Answer :-

Methyl groups in the following compound are

[Question ID = 828]

- 1. homotopic [Option ID = 3309]
- 2. enantiotopic [Option ID = 3311]
- 3. constitutionally heterotopic [Option ID = 3312]
- 4. diasterotopic [Option ID = 3310]

Correct Answer :-

homotopic [Option ID = 3309]

61)



What is the principal product of the following reaction?

[Question ID = 809]

1. [Option ID = 3234]

Correct Answer :-

Provide the suitable reagents for this conversion:

[Question ID = 833]

m-CPBA, HNO₃/H₂SO₄/PCI₃

HNO₃/H₂SO₄/POCl₃

2. [Option ID = 3332]

NaNO₂ /H₂SO₄/PCl₃

3. [Option ID = 3329]

H₂O₂/OH", HNO₃/H₂SO₄/PCI₃

4. [Option ID = 3330]

Correct Answer :-

m-CPBA, HNO₃/H₂SO₄/PCI₃

• [Option ID = 3331]

Identify the major product of the reaction?



[Question ID = 820]

[Option ID = 3280]

2.

[Option ID = 3278]

3.

[Option ID = 3279]

4.

[Option ID = 3277]

Correct Answer :-

[Option ID = 3278]

64) In the reaction given below,

R-CI

(ii) LiAlH₄

Product A

(i) AgCN

R-CI

(ii) LiAlH₄

Product B

The compound A and B are:

[Question ID = 814]

- 1. Metamers [Option ID = 3256]
- 2. Functional isomers [Option ID = 3255]
- 3. Chain isomers [Option ID = 3253]
- 4. Position isomers [Option ID = 3254]

Correct Answer :-

• Functional isomers [Option ID = 3255]

Which is product of the reaction:



[Question ID = 831]

1.

[Option ID = 3324]

2.

[Option ID = 3323]

3.

[Option ID = 3321]

4.

[Option ID = 3322]

Correct Answer :-

[Option ID = 3323]

An ionic solution consists of 0.2 mol dm⁻³ each of A²⁺ and B³⁻ ions. What is the ionic strength of the solution?

[Question ID = 854]

0.5 mol dm⁻³

[Option ID = 3416]

2. 1.0 mol dm⁻³

[Option ID = 3415]

3. 1.3 mol dm⁻³

[Option ID = 3414]

4. 2.6 mol dm⁻³

[Ontion ID = 3413]

Correct Answer :-

. 1.3 mol dm⁻³

[Option ID = 3414]

The molar weight of MgCO₃ is 84. The volume in litres of CO₂ at STP on heating 8.4g of MgCO₃ would be

[Question ID = 863]

1. 2.24 [Option ID = 3452]

2. 11.2 [Option ID = 3450]

3. 22.4 [Option ID = 3449]

4. 1.12 [Option ID = 3451]




- 2.24 [Option ID = 3452]
- It takes 20 minutes for the concentration of a radioactive species to decay to its 1/4th value of its original concentration. What is the rate constant of this radioactive decay reaction?

[Question ID = 856]


- 865.8 s^{-1} 1. [Option ID = 3424]
- 600 s⁻¹ [Option ID = 3421]
- 3. 415.8 s^{-1} [Option ID = 3423]
- 4. 0.001155 s^{-1} [Option ID = 3422]

Correct Answer :-

- 0.001155 s^{-1} [Option ID = 3422]
- Which of the following having the maximum Dipole moment?







[Question ID = 817]

- 1. i [Option ID = 3265]
- 2. ii [Option ID = 3266]
- 3. iv [Option ID = 3268]
- 4. iii [Option ID = 3267]

Correct Answer :-

- i [Option ID = 3265]
- 70)

. What is the likely product of the reaction shown?

[Question ID = 808]

2. [Option ID = 3232]



[Option ID = 3232]

The major product formed in the following reaction.

[Question ID = 835]

1. [Option ID = 3339]

2. [Option ID = 3340]

[Option ID = 3338]



[Option ID = 3339]

72) The major product formed in the following reaction:

[Question ID = 838]

1. [Option ID = 3351]

3. [Option ID = 3352]

5

Correct Answer :-

• [Option ID = 3349]

An optically active compound 'X' has molecular formula C₄H₈O₃. It evolves CO₂ with NaHCO₃. X reacts with LiAlH₄ to give achiral compounds. 'X' is:

[Question ID = 812]

CH₃CH₂CHCOOH

, oh

[Option ID = 3245]

CH₃CHCOOH

2. Me

3.

[Option ID = 3246]

CH₃CHCH₂COOH

ÓН

[Option ID = 3248]

СН₃СНСООН

ĊH₂OH

[Option ID = 3247]



CH₃CHCOOH

CH₂OH

[Option ID = 3247]

74)

Which is product of the reaction:

[Question ID = 832]

2. [Option ID = 3328]

3. [Option ID = 3326]

Correct Answer :-

The IUPAC name of the compound given below is

[Question ID = 821]

1. (2Z, 4Z)-3-chlorohexa-2, 4-diene-1,6-diol. [Option ID = 3283]

2. (2E, 4E)-3-chlorohexa-2, 4-diene-1,6-diol. [Option ID = 3281]

3. (2Z, 4Z)-3-chlorohexa-2, 4-diene-1,6-diol. [Option ID = 3284]

4. (2Z, 4E)-3-chlorohexa-2, 4-diene-1,6-diol. [Option ID = 3282]



(2Z, 4E)-3-chlorohexa-2, 4-diene-1,6-diol. [Option ID = 3282]

Arrange the following in decreasing order of O-O Bond length?

- (i) O₂
- (ii) O₂ + (iii) O₂ 2-
- (iv) O₃

[Question ID = 819]

- 1. iv>i>iii>ii [Option ID = 3274]
- 2. ii>i>iii>iv [Option ID = 3275]
- 3. i>iv>ii>iii [Option ID = 3276]
- 4. iii>iv>i>ii [Option ID = 3273]

Correct Answer :-

iii>iv>i>ii [Option ID = 3273]

PCl₅ does not react with

[Question ID = 779]

- CH₃COOH [Option ID = 3113]
- 2. C_6H_5OH [Option ID = 3115]
- 3. C_2H_5OH [Option ID = 3114]
- H_2SO_4 [Option ID = 3116]

Correct Answer :-

 C_6H_5OH [Option ID = 3115]

Partial pressure of CO2 in a mixture of CO2 and N2 is 1 atm while the total pressure of mixture is 5 atm. Mole fraction of nitrogen in the mixture is:

[Question ID = 873]

- 1. 0.65 [Option ID = 3492]
- 2. 0.8 [Option ID = 3491]
- 3. 0.75 [Option ID = 3490]
- 4. 0.82 [Option ID = 3489]

Correct Answer :-

0.8 [Option ID = 3491]

pH of the solution produced by mixing equal volumes of 2.0×10^{-3} M HClO₄ and 1.0×10^{-2} MKClO₄ is

[Question ID = 798]

- 1. 2.3 [Option ID = 3190]
- 2. 1 [Option ID = 3192]
- 3. 3 [Option ID = 3191]
- 4. 2.7 [Option ID = 3189]

Correct Answer :-

• 3 [Option ID = 3191]

80) . For a simple paramagnetic compound, which one of the following is true? [Question ID = 805]

- 1. Magnetic susceptibility decreases initially and then increases with decrease in temperature [Option ID = 3220]
- Magnetic susceptibility decreases with decrease in temperature [Option ID = 3218]



- 3. Magnetic susceptibility increases with decrease in temperature [Option ID = 3217]
- 4. Magnetic susceptibility increases initially and then decreases with decrease in temperature [Option ID = 3219]

Magnetic susceptibility increases with decrease in temperature [Option ID = 3217]

81) Two isotonic solutions will have same: [Question ID = 872]

- 1. Boiling point [Option ID = 3486]
- 2. Osmotic pressure [Option ID = 3485]
- 3. Vapour pressure [Option ID = 3488]
- 4. Freezing point [Option ID = 3487]

Correct Answer :-

• Osmotic pressure [Option ID = 3485]

82) Melting points of the chlorides of alkali metals decreases in the order [Question ID = 785]

- 1. LiCl > NaCl > KCl > RbCl > CsCl [Option ID = 3138]
- 2. LiCl > NaCl > KCl > RbCl > CsCl [Option ID = 3139]
- 3. LiCl > NaCl > KCl > RbCl > CsCl [Option ID = 3140]
- 4. LiCl > NaCl > KCl > RbCl > CsCl [Option ID = 3137]

Correct Answer :-

83) Residual entropy is the entropy of [Question ID = 862]

- 1. An isolated system [Option ID = 3445]
- 2. A system undergoing reversible reaction [Option ID = 3448]
- 3. A system at equilibrium [Option ID = 3446]
- 4. A system at absolute zero of temperature [Option ID = 3447]

Correct Answer :-

A system at absolute zero of temperature [Option ID = 3447]

84) Although carbon and oxygen are the constituents of carbonate and oxalate, the reason behind oxalate being an interfering anion [Question ID = 803]

- 1. Higher oxidizability of oxalate than carbonate [Option ID = 3210]
- 2. Higher reducibility of oxalate than carbonate [Option ID = 3209]
- 3. Higher chelating ability of oxalate than carbonate [Option ID = 3211]
- 4. Higher polarisability of oxalate than carbonate [Option ID = 3212]

Correct Answer :-

Higher chelating ability of oxalate than carbonate [Option ID = 3211]

85) The major product formed in the dinitration of 4-bromotoluene is. [Question ID = 837]

, B

[Option ID = 3348]

2.

[Option ID = 3345]

. Br [Option ID = 3347]



[Option ID = 3346]

Correct Answer :-

[Option ID = 3346]

86) Electronic spin a has eigen value [Question ID = 861]

2.
$$h/4\pi$$
 [Option ID = 3441]

3.
$$1/2h$$
 [Option ID = 3442]

Correct Answer :-

•
$$h/4\pi$$
 [Option ID = 3441]

87) Which of the following shows Jahn-Teller Distortion? [Question ID = 799]

$$Mn^{2+}$$

$$\frac{Mn^{2+}}{2}$$
 [Option ID = 3195]

4.
$$Fe^{2+}$$
 [Option ID = 3193]

Correct Answer :-

88) Which of the following is an incorrect representation of the order of a reaction: [Question ID = 848]

$$N_2O_5(g)$$

$$\rightarrow$$

$$2N_2(g) + \frac{1}{2} O$$

$$N_2O_5(g) \rightarrow 2N_2(g) + \frac{1}{2}O_2$$
 is a 1st order reaction

$$2CH_4 + 2CC$$

$$2CH_3CHO \rightarrow 2CH_4 + 2CO$$
 is a 2nd order reaction

$$S_2O_8^{2-} + 2I^- \rightarrow 2SO_4^{2-} + I_2$$

Correct Answer :-

$$S_2O_8^{2-} + 2I \rightarrow 2SO_4^{2-} + I_2$$
 is a 3rd order reaction

89) Which of the following pair has the same electronic structure? [Question ID = 775]



Ar, Cl⁻ [Option ID = 3100]

90) Which of the following is not a colligative property? [Question ID = 842]

- 1. Osmotic pressure [Option ID = 3368]
- 2. Relative increase in vapour pressure [Option ID = 3367]
- 3. Depression of freezing point [Option ID = 3366]
- 4. Elevation of boiling point [Option ID = 3365]

Correct Answer :-

• Relative increase in vapour pressure [Option ID = 3367]

91) Which of the following statement is false? [Question ID = 859]

- 1. Oxidation reaction takes place at the cathode of a galvanic cell [Option ID = 3435]
- 2. The potential of normal hydrogen electrode (NHE) is assigned a value of zero volts [Option ID = 3436]
- 3. The EMF of a galvanic cell can be measured with a voltmeter [Option ID = 3433]
- 4. Oxidation reaction takes place at the anode of a galvanic cell [Option ID = 3434]

Correct Answer :-

Oxidation reaction takes place at the cathode of a galvanic cell [Option ID = 3435]

92) Which one of the following is a superconductor? [Question ID = 796]

$$YB_{2}Cu_{3}O_{7} \\ YBe_{2}Cu_{3}O_{7} \\ 2. \qquad \qquad [\text{Option ID = 3184}] \\ YBi_{2}Cu_{3}O_{7} \\ 3. \qquad YBi_{2}Cu_{3}O_{7} \\ 4. \qquad YBa_{2}Cu_{3}O_{7} \\ [\text{Option ID = 3181}] \\ 4. \qquad (\text{Option ID = 3182}]$$

Correct Answer :-

 $YBa_2Cu_3O_7$ [Option ID = 3182]

93) Which one of the following plays a major role in EDTA complexometric titrations? [Question ID = 797]

- 1. Concentration of ligand [Option ID = 3186]
- 2. Concentration of metal ion [Option ID = 3185]
- 3. Temperature of the reaction [Option ID = 3188]
- 4. Nature of buffer [Option ID = 3187]

Correct Answer :-

• Nature of buffer [Option ID = 3187]

94) Pyrosilicates are the silicates in which the two tetrahedral units are linked at [Question ID = 780]

- 1. Three points [Option ID = 3119]
- 2. One point [Option ID = 3117]
- 3. Four points [Option ID = 3120]
- 4. Two points [Option ID = 3118]

Correct Answer :-

• One point [Option ID = 3117]

95) In a face-centre cubic (FCC) type of crystal lattice, the number of atoms belonging exclusively to each unit cell within the lattice is/are: [Question ID = 865]

- 1. 4 [Option ID = 3460]
- 2. 2 [Option ID = 3458]
- 3. 3 [Option ID = 3459]
- 1 [Option ID = 3457]

Correct Answer :-

4 [Option ID = 3460]

96) Langmuir adsorption isotherm equation shows the variation of extent of adsorption as a function of: [Question ID = 847]



- 1. pH of medium [Option ID = 3387]
- 2. Pressure [Option ID = 3385]
- 3. Temperature [Option ID = 3386]
- 4. All of these [Option ID = 3388]

Pressure [Option ID = 3385]

97) According to Lambert-Beer's law, for a solution the transmittance is independent of which following factor? [Question ID = 867]

- 1. Molar extinction coefficient of the solute in solution. [Option ID = 3468]
- 2. Path length of the sample holder [Option ID = 3466]
- 3. Concentration of the solution [Option ID = 3465]
- 4. Temperature of the system [Option ID = 3467]

Correct Answer :-

• Temperature of the system [Option ID = 3467]

98) The compressibility factor for ideal gas is:

[Question ID = 871]

- 1. 1 [Option ID = 3483]
- 2. >1 [Option ID = 3482]
- 3. Zero [Option ID = 3481]
- 4. <1 [Option ID = 3484]

Correct Answer :-

1 [Option ID = 3483]

99) The following compounds have been arranged in the order of increasing thermal stabilities. Identify the correct order K₂CO₃ (I), MgCO₃ (II), CaCO₃ (III), BeCO₃ (IV)

[Question ID = 784]

- 1. II<IV<III<I [Option ID = 3136]
- 2. IV<II<I<III [Option ID = 3135]
- 3. IV < II < III < I [Option ID = 3134]
- 4. I<II<III<IV [Option ID = 3133]

Correct Answer :-

IV<II<I<III [Option ID = 3135]

100) The covalent radius of Li is 123 pm. The crystal radius of Li will be [Question ID = 774]

$$> 123 \text{ pm}$$
 [Option ID = 3093]

