

Magnetic Effects Of Current And Magnetism

JEE Main PYQ – 3

Total Time: 25 Minute

Total Marks: 40

Instructions

Instructions

1. Test will auto submit when the Time is up.
2. The Test comprises of multiple choice questions (MCQ) with one or more correct answers.
3. The clock in the top right corner will display the remaining time available for you to complete the examination.

Navigating & Answering a Question

1. The answer will be saved automatically upon clicking on an option amongst the given choices of answer.
2. To deselect your chosen answer, click on the clear response button.
3. The marking scheme will be displayed for each question on the top right corner of the test window.

Magnetic Effects Of Current And Magnetism

1. In a circuit for finding the resistance of a galvanometer by half deflection method, a 6 V battery and a high resistance of $11\text{ k}\Omega$ are used. The figure of merit of the galvanometer is $60\text{ }\mu\text{A}$ division. In the absence of shunt resistance, the galvanometer produces a deflection of $\theta = 9$ divisions when current flows in the circuit. The value of the shunt resistance that can cause the deflection of $\theta/2$, is closest to :

(+4, -1)

[Online April 16, 2018]

- a. $550\ \Omega$
- b. $220\ \Omega$
- c. $55\ \Omega$
- d. $110\ \Omega$

2. Two wires A & B are carrying currents I_1 & I_2 as shown in the figure. The separation between them is d . A third wire C carrying a current I is to be kept parallel to them at a distance x from A such that the net force acting on it is zero. The possible values of x are :

(+4, -1)

[10 April 2019, I]

- a. $x = \left(\frac{I_1}{I_1 - I_2}\right)d$ and $x = \frac{I_2}{(I_1 + I_2)}d$
- b. $x = \pm \frac{I_1 d}{(I_1 + I_2)}$
- c. $x = \left(\frac{I_1}{I_1 + I_2}\right)d$ and $x = \frac{I_2}{(I_1 - I_2)}d$
- d. $x = \left(\frac{I_1}{I_1 + I_2}\right)d$ and $x = \left(\frac{I_2}{I_1 + I_2}\right)d$

3. When a current of 5 mA is passed through a galvanometer having a coil of resistance $15\ \Omega$, it shows full scale deflection. The value of the resistance to be put in series with the galvanometer to convert it into a voltmeter of range $0 - 10\text{ V}$ is :

(+4, -1)

[2017]


- a. $1.985 \times 10^3\ \Omega$
- b. $2.045 \times 10^3\ \Omega$

c. $2.535 \times 10^3 \Omega$

d. $4.005 \times 10^3 \Omega$

-
4. The magnetic moment of an octahedral homoleptic Mn(II) complex is $5.9 BM$ (+4, -1). The suitable ligand for this complex is :

[Online April 16, 2018]



-
5. This question has Statement I and Statement II. Of the four choices given (+4, -1) after the Statements, choose the one that best describes the two Statements. Higher the range, greater is the resistance of ammeter. To increase the range of ammeter, additional shunt needs to be used across it.

a. Statement - I is true, Statement - II is true, Statement - II is the correct explanation of Statement-I

b. Statement - I is true, Statement - II is true, Statement - II is not the correct explanation of Statement-I

c. Statement - I is true, Statement - II is false

d. Statement - I is false, Statement - II is true

-
6. To know the resistance G of a galvanometer by half deflection method, a (+4, -1) battery of emf V_E and resistance R is used to deflect the galvanometer by angle θ . If a shunt of resistance S is needed to get half deflection then G , R and S are related by the equation :

[Online April 9, 2016]

a. $2S(R + G) = RG$

b. $S(R + G) = RG$

c. $2S = G$

d. $2G = S$

7. Two coaxial solenoids of different radii carry current I in the same direction. (+4, -1)
 Let \vec{F}_1 be the magnetic force on the inner solenoid due to the outer one and \vec{F}_2 be the magnetic force on the outer solenoid due to the inner one. Then,

a. \vec{F}_1 is radially outwards and $\vec{F}_2 = 0$ [2015]

b. \vec{F}_1 is radially inwards and \vec{F}_2 is radially outwards

c. \vec{F}_1 is radially inwards and $\vec{F}_2 = 0$

d. $\vec{F}_1 = \vec{F}_2 = 0$

8. A current i is flowing through the loop. The direction of the current and the shape of the loop are as shown in the figure. The magnetic field at the centre of the loop is, $\frac{0}{16}$ times (+4, -1)

[$MA = R$, $MB = 2R$, $\angle MA = 90^\circ$]

a. $\frac{5}{16}$, but out of the plane of the paper

b. $\frac{5}{16}$, but into the plane of the paper

c. $\frac{7}{16}$, but out of the plane of the paper

d. $\frac{7}{16}$, but into the plane of the paper

9. The magnetic moments associated with two closely wound circular coils A and B of radius $r_A = 10 \text{ cm}$ and $r_B = 20 \text{ cm}$ respectively are equal if : (Where N_A, I_A and N_B, I_B are number of turn and current of A and B respectively) (+4, -1)

a. $2N_A I_A = N_B I_B$ [30-Jan-2023 Shift 1]

b. $N_A = 2N_B$

c. $N_A I_A = 4N_B I_B$

d. $4N_A I_A = N_B I_B$

10. Two long straight wires P and Q carrying equal current 10 A each were kept parallel to each other at 5 cm distance. Magnitude of magnetic force experienced by 10 cm length of wire P is F_1 . If distance between wires is halved and currents on them are doubled, force F_2 on 10 cm length of wire P will be: (+4, -1)

[24-Jan-2023 Shift 1]

a. $10F_1$

b. $8F_1$

c. $\frac{F_1}{8}$

d. $\frac{F_1}{10}$



Answers

1. Answer: d

Explanation:

Current required by unit deflection is $60 \mu A$

For, $\theta = 9$ current is $I = 9 \times 60 \mu A$

$$\Rightarrow I = 540 \mu A = 540 \times 10^{-6} A$$

Let G is resistance of galvanometer. Then,

$$540 \times 10^{-6} = \frac{6}{(11000+G)}$$

$$[11000 + G]90 \times 10^{-6} = 1$$

$$99000 + 9G = 10^5$$

$$9G = 100000 - 99000$$

$$9G = 1000$$

$$G = \frac{1000}{9} \Omega$$

Also in half deflection method,

$$G = \frac{RS}{R-S}$$

$$\Rightarrow \frac{1000}{9} = \frac{11000S}{11000-S}$$

$$\frac{1}{9} = \frac{11S}{11000-S}$$

$$\Rightarrow 11000 - S = 99S$$

$$100S = 11000 \Rightarrow S = 110 \Omega$$

Concepts:

1. Moving Charges and Magnetism:

Moving charges generate an electric field and the rate of flow of charge is known as **current**. This is the basic concept in **Electrostatics**. Another important concept related to moving **electric charges** is the magnetic effect of current. Magnetism is caused by the current.

Magnetism:

- The relationship between a [Moving Charge and Magnetism](#) is that Magnetism is produced by the movement of charges.
- And Magnetism is a property that is displayed by Magnets and produced by moving charges, which results in objects being attracted or pushed away.

Magnetic Field:

Region in space around a magnet where the magnet has its magnetic effect is called the magnetic field of the magnet. Let us suppose that there is a point charge q (moving with a velocity v and, located at r at a given time t) in presence of both the electric field $E(r)$ and the magnetic field $B(r)$. The force on an electric charge q due to both of them can be written as,

$$F = q [E(r) + v \times B(r)] \equiv F_{\text{Electric}} + F_{\text{magnetic}}$$

This force was based on the extensive experiments of Ampere and others. It is called the Lorentz force.

2. Answer: b

Explanation:

Net force on wire carrying current I per unit length is

$$\begin{aligned} \frac{\mu_0 I_1 I}{2\pi x} + \frac{\mu_0 I_2 I}{2\pi(d-x)} &= 0 \\ \frac{I_1}{x} &= \frac{I_2}{x-d} \\ \Rightarrow x &= \frac{I_1 d}{I_1 - I_2} \end{aligned}$$

Concepts:

1. Moving Charges and Magnetism:

Moving charges generate an electric field and the rate of flow of charge is known as **current**. This is the basic concept in **Electrostatics**. Another important concept related to moving **electric charges** is the magnetic effect of current. Magnetism is caused by the current.

Magnetism:

- The relationship between a [Moving Charge and Magnetism](#) is that Magnetism is produced by the movement of charges.
- And Magnetism is a property that is displayed by Magnets and produced by moving charges, which results in objects being attracted or pushed away.

Magnetic Field:

Region in space around a magnet where the Magnet has its Magnetic effect is called the Magnetic field of the Magnet. Let us suppose that there is a point charge q (moving with a velocity v and, located at r at a given time t) in presence of both the electric field $E(r)$ and the magnetic field $B(r)$. The force on an electric charge q due to both of them can be written as,

$$F = q [E(r) + v \times B(r)] \equiv F_{\text{Electric}} + F_{\text{magnetic}}$$

This force was based on the extensive experiments of Ampere and others. It is called the Lorentz force.

3. Answer: a

Explanation:

$$i_g = 5 \times 10^{-3} A$$

$$G = 15 \Omega$$

Let series resistance be R .

$$V = i_g(R + G)$$

$$10 = 5 \times 10^{-3}(R + 15)$$

$$R = 2000 - 15 = 1985 = 1.985 \times 10^3 \Omega$$

Concepts:

1. Moving Charges and Magnetism:

Moving charges generate an electric field and the rate of flow of charge is known as **current**. This is the basic concept in **Electrostatics**. Another important concept related to moving **electric charges** is the magnetic effect of current. Magnetism is caused by the current.

Magnetism:

- The relationship between a [Moving Charge and Magnetism](#) is that Magnetism is produced by the movement of charges.

- And Magnetism is a property that is displayed by Magnets and produced by moving charges, which results in objects being attracted or pushed away.

Magnetic Field:

Region in space around a magnet where the Magnet has its Magnetic effect is called the Magnetic field of the Magnet. Let us suppose that there is a point charge q (moving with a velocity v and, located at r at a given time t) in presence of both the electric field $E(r)$ and the magnetic field $B(r)$. The force on an electric charge q due to both of them can be written as,

$$F = q [E(r) + v \times B(r)] \equiv F_{\text{Electric}} + F_{\text{magnetic}}$$

This force was based on the extensive experiments of Ampere and others. It is called the Lorentz force.

4. Answer: b

Explanation:

$\mu = 5.9BM \therefore n$ (no of unpaired.e-) = 5 Cation $Mn^{II} - 3d^5$ confn only possible for relatively weak ligand. $\therefore NCS^-$

Concepts:

1. Moving Charges and Magnetism:

Moving charges generate an electric field and the rate of flow of charge is known as **current**. This is the basic concept in **Electrostatics**. Another important concept related to moving **electric charges** is the magnetic effect of current. Magnetism is caused by the current.

Magnetism:

- The relationship between a [Moving Charge and Magnetism](#) is that Magnetism is produced by the movement of charges.
- And Magnetism is a property that is displayed by Magnets and produced by moving charges, which results in objects being attracted or pushed away.

Magnetic Field:

Region in space around a magnet where the Magnet has its Magnetic effect is called the Magnetic field of the Magnet. Let us suppose that there is a point charge q (moving with a velocity v and, located at r at a given time t) in presence of both the electric field $E(r)$ and the magnetic field $B(r)$. The force on an electric charge q due to both of them can be written as,

$$F = q [E(r) + v \times B(r)] \equiv F_{\text{Electric}} + F_{\text{magnetic}}$$

This force was based on the extensive experiments of Ampere and others. It is called the Lorentz force.

5. Answer: d

Explanation:

Answer (d) Statement - I is false, Statement - II is true

Concepts:

1. Moving Charges and Magnetism:

Moving charges generate an electric field and the rate of flow of charge is known as **current**. This is the basic concept in **Electrostatics**. Another important concept related to moving **electric charges** is the magnetic effect of current. Magnetism is caused by the current.

Magnetism:

- The relationship between a [Moving Charge and Magnetism](#) is that Magnetism is produced by the movement of charges.
- And Magnetism is a property that is displayed by Magnets and produced by moving charges, which results in objects being attracted or pushed away.

Magnetic Field:

Region in space around a magnet where the Magnet has its Magnetic effect is called the Magnetic field of the Magnet. Let us suppose that there is a point charge q (moving with a velocity v and, located at r at a given time t) in presence of both the electric field $E(r)$ and the magnetic field $B(r)$. The force on an electric charge q due to both of them can be written as,

$$F = q [E(r) + v \times B(r)] \equiv F_{\text{Electric}} + F_{\text{magnetic}}$$

This force was based on the extensive experiments of Ampere and others. It is called the Lorentz force.

6. Answer: b

Explanation:

$$I_g = \frac{V}{R+G} \quad R_c = R + \frac{GS}{G+S} \quad I = \frac{V}{R+\frac{GS}{G+S}} \quad I'_g G = (I - I'_g) S \quad I'_g (G + S) = IS \quad \frac{I_g}{2} = \frac{IS}{G+S} \quad \frac{V}{2(R+G)} = \frac{V}{R+\frac{GS}{G+S}} \times \frac{S}{G+S} \quad \frac{1}{2(R+G)} = \frac{S}{R(G+S)+GS} \quad R(G+S) + GS = 2S(R+G) \quad RG + RS + GS = 2S(R+G) \quad RG = 2S(R+G) - S(R+G) \quad RG = S(R+G)$$

Concepts:

1. Moving Charges and Magnetism:

Moving charges generate an electric field and the rate of flow of charge is known as **current**. This is the basic concept in **Electrostatics**. Another important concept related to moving **electric charges** is the magnetic effect of current. Magnetism is caused by the current.

Magnetism:

- The relationship between a [Moving Charge and Magnetism](#) is that Magnetism is produced by the movement of charges.
- And Magnetism is a property that is displayed by Magnets and produced by moving charges, which results in objects being attracted or pushed away.

Magnetic Field:

Region in space around a magnet where the Magnet has its Magnetic effect is called the Magnetic field of the Magnet. Let us suppose that there is a point charge q (moving with a velocity v and, located at r at a given time t) in presence of both the electric field $E(r)$ and the magnetic field $B(r)$. The force on an electric charge q due to both of them can be written as,


$$F = q [E(r) + v \times B(r)] \equiv F_{\text{Electric}} + F_{\text{magnetic}}$$

This force was based on the extensive experiments of Ampere and others. It is called the Lorentz force.


7. Answer: d

Explanation:

Cross-sectional view



(Both solenoids are taken to be ideal in nature.) Both wires will attract each other, but net force on each wire will be zero. Concept: Two current carrying elements attract each other if direction of current is same. *F.B.D*



$$\vec{F}_1 = 0 \quad \vec{F}_2 = 0$$

Concepts:

1. Moving Charges and Magnetism:

Moving charges generate an electric field and the rate of flow of charge is known as **current**. This is the basic concept in **Electrostatics**. Another important concept related to moving **electric charges** is the magnetic effect of current. Magnetism is caused by the current.

Magnetism:

- The relationship between a [Moving Charge and Magnetism](#) is that Magnetism is produced by the movement of charges.
- And Magnetism is a property that is displayed by Magnets and produced by moving charges, which results in objects being attracted or pushed away.

Magnetic Field:

Region in space around a magnet where the Magnet has its Magnetic effect is called the Magnetic field of the Magnet. Let us suppose that there is a point charge q (moving with a velocity v and, located at r at a given time t) in presence of both the electric field $E(r)$ and the magnetic field $B(r)$. The force on an electric charge q due to both of them can be written as,

$$F = q [E (r) + v \times B (r)] \equiv E_{\text{Electric}} + F_{\text{magnetic}}$$

This force was based on the extensive experiments of Ampere and others. It is called the Lorentz force.

8. Answer: d

Explanation:

Explanation:

Given:

$MA = R, MB = 2R, \angle DMA = 90^\circ$ We have to find the magnetic field at the center of the loop. The magnitude of magnetic field due to an arc of radius R and angle θ , carrying

current at the center of the loop is $B = \frac{\mu_0 I}{4\pi R} \theta$ where, θ is the angle subtended by the arc. Given figure is redrawn below:



The magnetic field due to curve DA is $B_{DA} = \frac{\mu_0 I}{4\pi R} (3\pi)$

Similarly, magnetic field due to the arc BC is $B_{BC} = \frac{\mu_0 I}{4\pi R} (\pi)$ [given $MB=2R$]

Also, the magnetic field due to the curve DC and AB is $B_{DC} = B_{AB} = 0$ [since the angle between the current element and the position vector is zero, i.e. $\sin 0 = 0$]

Now, the resultant magnetic field at the centre M is $B_{net} = B_{DA} + B_{AB} + B_{BC} + B_{CD}$

$$\Rightarrow B_{net} = \frac{\mu_0 I}{4\pi R} (3\pi) + 0 + \frac{\mu_0 I}{4\pi R} (\pi) + 0$$

$$\Rightarrow B_{net} = \frac{\mu_0 I}{4\pi R} [3\pi + \pi] \Rightarrow B_{net} = \frac{\mu_0 I}{4\pi R} [4\pi] \Rightarrow B_{net} = \mu_0 I / R$$

Using the right-hand thumb rule for the magnetic field, the direction of the magnetic field is into the plane of the paper, as the direction of the current is clockwise.

Hence, the correct option is (D).

9. Answer: c

Explanation:

Sol. $M = NIA$

$$M_A = M_B$$

$$\therefore N_A I_A A_A = N_B I_B A_B$$

$$\therefore N_A I_A \pi (0.1)^2 = N_B I_B \pi (0.2)^2$$

$$\therefore N_A I_A = 4 N_B I_B$$

Concepts:

1. Moving Charges and Magnetism:

Moving charges generate an electric field and the rate of flow of charge is known as **current**. This is the basic concept in **Electrostatics**. Another important concept related to moving **electric charges** is the magnetic effect of current. Magnetism is caused by the current.

Magnetism:

- The relationship between a [Moving Charge and Magnetism](#) is that Magnetism is produced by the movement of charges.

- And Magnetism is a property that is displayed by Magnets and produced by moving charges, which results in objects being attracted or pushed away.

Magnetic Field:

Region in space around a magnet where the Magnet has its Magnetic effect is called the Magnetic field of the Magnet. Let us suppose that there is a point charge q (moving with a velocity v and, located at r at a given time t) in presence of both the electric field $E(r)$ and the magnetic field $B(r)$. The force on an electric charge q due to both of them can be written as,

$$F = q [E(r) + v \times B(r)] \equiv F_{\text{Electric}} + F_{\text{magnetic}}$$

This force was based on the extensive experiments of Ampere and others. It is called the Lorentz force.

10. Answer: b

Explanation:

Force per unit length between two parallel straight

$$\text{wires} = \frac{\mu_0 i_1 i_2}{2\pi d}$$

$$\frac{F_1}{F_2} = \frac{\frac{\mu_0 (10)^2}{2\pi(5cm)}}{\frac{\mu_0 (20)^2}{2\pi(\frac{5cm}{2})}} = \frac{1}{8}$$

$$\Rightarrow F_2 = 8F_1$$

Concepts:

1. Moving Charges and Magnetism:

Moving charges generate an electric field and the rate of flow of charge is known as [current](#). This is the basic concept in [Electrostatics](#). Another important concept related to moving [electric charges](#) is the magnetic effect of current. Magnetism is caused by the current.

Magnetism:

- The relationship between a [Moving Charge and Magnetism](#) is that Magnetism is produced by the movement of charges.
- And Magnetism is a property that is displayed by Magnets and produced by moving charges, which results in objects being attracted or pushed away.

Magnetic Field:

Region in space around a magnet where the Magnet has its Magnetic effect is called the Magnetic field of the Magnet. Let us suppose that there is a point charge q (moving with a velocity v and, located at r at a given time t) in presence of both the electric field $E (r)$ and the magnetic field $B (r)$. The force on an electric charge q due to both of them can be written as,

$$F = q [E (r) + v \times B (r)] \equiv E_{\text{Electric}} + F_{\text{magnetic}}$$

This force was based on the extensive experiments of Ampere and others. It is called the Lorentz force.

