## **MH CET 2018**

## (QUESTION WITH ANSWER)

## **BIOLOGY**

- Cambium is essential for grafting in plants because
  - (1) Cambia of both stock and scion fuse together
  - (2) Cambium produces new leaves
  - (3) Cambium produces new roots
  - (4) Cambium helps in the production of flowers

Ans: (1)

- Sol: Cambium is essential for grafting in plants because cambia which consists of meristematic cells of stock and scion fuse together.
- **2.** Which one of the following carbohydrates is a heteropolysaccharide?
  - Cellulose
  - (2) Starch
  - (3) Glycogen
  - (4) Hyaluronic acid

Ans: (4)

- Sol: Hyaluronic acid is a hetropolysaccharide. It is made up of glucuronic acid and N-acetyl-D-glucosamine disaccharide units.
- "Pusa Sadabahar" is resistant to disease.
  - (1) Leaf curl and chilli mosaic virus
  - (2) Leaf and stripe rust
  - (3) Black rot

(4) Curl blight black rot

Ans: (A)

Sol: 'Pusa Sadabhar' is resistance to disease leaf curl and chilli mosaic virus of chilli. Himgiri of wheat is resistance to leaf and stripe rust.

- 4. The adsorption of water by hydrophilic compounds like cellulose and pectin in root hair cell wall is called
  - (1) Diffusion

(2) Imbibition

(3) Guttation

(4) Osmosis

Ans: (2)

Sol: The adsorption of water by hydrophilic compounds like cellulose and pectine in root hair cell wall is called imbibition.

- 5. In Kreb's cycle, the conversion of oxalosuccinate into -ketoglutarate involves
  - (1) Oxidation

(2) Reduction

(3) Hydration

(4) Decarboxylation

Ans: (4)

- Sol: During Kreb's cycle, oxalosuccinate is converted into  $\alpha$ -ketoglutarate by oxidative decarboxylation.
- 6. Double fertilization in angiosperms was first discovered by S.G. Nawaschin in plants.
  - (1) Lillium

(2) Sunflower

3) Wheat

(4) Mango



Ans:(A)

Sol: Double fertilization in angiosperms was first discovered by S.G. Nawaschin in Lillium plants.

- Oxidative phosphorylation in eukaryotes occurs during \_\_\_\_\_\_.
  - (1) Photosynthesis
  - (2) Protein synthesis
  - (3) Lipid synthesis
  - (4) Respiration

Ans: (4)

Sol: Oxidative phosphorylation in eukaryotes occurs during respiration (Glycolysis). In oxidative pay off phase of glycolysis in which PGAL undergoes oxidation and phosphorylation.

- **8.** Snapdragon is an example of gene interaction.
  - (1) Co-dominance
  - (2) Multiple alleles
  - (3) Incomplete dominance
  - (4) Polygenic inheritance

Ans: (3)

Sol: Snapdragon is an example of incomplete dominance.

- 9. In one nucleosome, which one of the following histone molecule is NOT double?
  - (1)  $H_4$
- (2)  $H_3$
- (3)  $H_2$
- (4) H

Ans: (4)

Sol: H<sub>1</sub> histone is present in the linker region. In the nucleosome, two molecules of each of H<sub>2</sub>A, H<sub>2</sub>B, H<sub>3</sub> and H<sub>4</sub> are present to form octamer.

- 10. Which one of the following is NOT true about Salvia flower?
  - (1) Flowers are bisexual
  - (2) Flowers are protandrous
  - (3) The upper branch of bifurcated connective bears sterile anther lobe and lower branch bears fertile anther lobe
  - (4) The upper branch of bifurcated connective bears fertile anther lobe and lower branch bears sterile anther lobe

Ans: (4)

- Identify the INCORRECT statement regarding Bt Cotton.
  - Dried spores of Bacillus thuringiensis are sprayed on its vulnerable leaves
  - (2) It releases toxin that causes swelling in the gut of insect that ingest its leaves
  - (3) It is a transgenic plant
  - (4) It can produce toxin due to gene introduced in it by transgenesis

Ans: (1)

- 12. In angiosperms, the fusion of male gamete with the secondary nucleus is considered as "Second fertilization" because
  - (1) It is fusion of two nuclei
  - (2) Secondary nucleus is a sister nucleus of the egg
  - (3) It takes place in embryo sac
  - (4) It takes place after pollination

Ans: (2)

Sol: In angiosperms, secondary nucleus and egg
cell is formed from haploid nucleus of
functional megaspore which undergoes three
successive free-nuclear mitotic division out
of which one nucleus from either of the pole
come to the centre and then functions as



	pola	ar nuclei. So, sec	ondary	nucleus is a		(2)	Prevention of al	oscission	of leaves and
	sist	er of nucleus of	the egg	<b>5.</b>		flow	ers by auxin ap	plication	n
						(3)	Increasing com	mercial v	alue of leafy
13.	In A	Airabilis jalapa, v	vhen re	ed and white		vege	etables by induc	ing leaf	expansion
	vari	eties are crossed	l, the h	ybrid obtained will		thro	ugh gibberellin	applicat	tion
	be_	pink.		70		(4)	Breaking dorr	nancy of	seeds and buds
	(1) 2	25%	(2)	50%		thro	ugh application	of ethyl	ene
	\$ 8	75%	(4)	100%	Ans	: (1)			
Ans	50 50		3 3		Sol	: Cyto	kinins delays tl	ne onset	of senescence by
	10 10	irabilis jalapa, ir	compl	ete dominance is		app	lication of cytol	kinins.	
		n. So all hybrid o							
	pinl	en e			17.	In a	person sufferin	ng from s	ickle cell
	•					ana	emia, the	be	comes half moon
14.	Dur	ing r-DNA techn	ology v	which one of the		sha	ped.		
			· · · · · · · · · · · · · · · · · · ·	for cleaving DNA		(1)	RBC	(2)	Thrombocyte
		ecule ?	asca .	ior creaving Dim		(3)	Monocyte	(4)	Lymphocyte
	(1)	Exonuclease			Ans	: (A)			
	100 TO	Endonuclease			Sol	: In a	person suerein	g from s	ickle cell anaemia
	(2) (3)	Restriction end	lonuole	222		the	R.B.C's become	e half mo	oon shaped.
	(4)	Helicase	onacie	asc					
Ans	8 3	Trenease			18.	Whi	ch one of the fo	llowing i	s true about
	2 8	technology restr	riction	endonuclease		res	piration in gree	n plants	? Respiration
3.8				g DNA molecule.		tak	es place		
		,		9		(1)	Only when ste	omata ai	re open
15.	Chir	oko Movement to	ok plac	ce in		(2)	Only when pl	notosynt	hesis stops
		Montreal				(3)	Only when ph	otosyntl	nesis is in
	(1)					pro	gress		
	(2)	Jodhpur				(4)	Throughout li	fe	
	(3)	Bangalore			Ans	: (4)			
	(4)	Garhwal Himal	layas		Sol	: Resp	piration is very i	importar	nt process for
Ans	: (4)					sur	vival of the livin	ig organi	sms as it oxidizes
Sol:	The	Chipka moveme	nt start	ted in the early		res	piratory subtrat	e and er	nergy is released
	197	0s in the Garhw	al of U	ttarakhand with		in t	he form of ATP	which is	required for all
	grov	wing awareness	toward	s rapid		met	tabolic activities	3.	
	defo	orestation.							
					19.		eukaryotes, the		
16.	In p	lants, Richmond	-Lang	effect is		5. <b>5</b> 12.25563		t codon o	of mRNA during
	(1)	Delaying the ons	et of se	enescence by		trai	nslation is		
	appl	ication of cytoki	nin			(1) U	JAA	(2) U	ICA



(3) UAC

(4) UUU

Ans: (3)

- Sol: In eukaryotes, initial codon on m-RNA is AUG. During translation, t-RNA, anticodons are exactly complementary to m-RNA. Hence anticodon will be UAC.  $[A = U, U = A, C \equiv G]$
- **20.** Biopatent is granted for biological entities by authority of a country.
  - (1) Government
  - (2) Educational institute
  - (3) Scientist
  - (4) Business

Ans: (1)

Sol: Biopatent is granted for biological entities by government authority of a country.

- 21. The exine of pollen grain is made up of
  - (1) Chitin
  - (2) Cellulose
  - (3) Sporopollenin
  - (4) Hemicellulose

Ans: (3)

Sol: The exine of pollen grain is made up of sporopollenin.

- **22.** Continuous upward flow of water stream in tall trees is maintained due to .
  - (1) Guttation and transpiration
  - (2) Transpiration pull only
  - (3) Cohesive force between H<sub>2</sub>O molecules only
  - (4) Cohesive force between H<sub>2</sub>O molecules and transpiration pull

Ans: (4)

Sol: Continuous upward flow of water stream in tall tree is maintained due to cohesive force between H<sub>2</sub>O molecules and transpiration pull.

- **23.** Which of the following is **NOT** correct regarding plasmids?
  - (1) They are small, extra chromosomal circular form of DNA
  - (2) They are naturally found in bacteria only
  - (3) They are 'replicons' capable of autonomous replication within a suitable host
  - (4) They carry genes related to survival and reproduction of carrier bacterium under unfavourable conditions

Ans: (4)

- 24. Algae used for the discovery of path of carbon in dark reaction were \_\_\_\_\_\_.
  - (1) Spirogyra and Chlorella
  - (2) Spirulina and Scenedesmus
  - (3) Chlorella and Scenedesmus
  - (4) Spirogyra and Spirulina

Ans: (3)

- Sol: Algae used for the discovery of path of carbon in dark reaction were chlorella and scenedesmus.
- **25.** Which one of the following is **NOT** a plant like protest?
  - (1) Desmid
  - (2) Dinoflagellate
  - (3) Diatom
  - (4) Slime mould

Ans. (4)

- Sol. Desmid, Dinoflagellate and Diatom are the plant - like protist. Slime mould is fungi like protist.
- 26. In case of anaerobic respiration, the RQ is

(1) Always less than one



	(2) Always more than one		(1) z
	(3) Always infinity		(2) y
	(4) Variable on the basis of substrate		(3) a
Ans.	(3)		(4) i
Sol.	In case of anaerobic respiration, the RQ is	Ans.	(4)
	always infinity.	Sol.	'i' gene (Regulatory gene) transcribes a
			repressor mRNA when glucose is the normal
27.	Which one of the following is NOT true about		energy source in bacterial cell.
	self pollination?.		
	(1) A sure method	31.	Which character of angiosperms helped in
	(2) Most economic		their dominance on earth?
	(3) Maintains genetic purity		(1) Formation of seeds
	(4) Favours evolution		(2) Formation of endosperm
Ans.	(4)		(3) Double fertilization
Sol.			(4) Presence of xylem vessels
		Ans.	(3)
28.	In biogas plant, the role of acidogenic	Sol.	Double fertilization is the characteristic
	bacteria is to.		feature of angiosperms.
	(1) Convert monomers to organic acids		
	(2) Convert monomers to complex polymers	32.	Identify the WRONG statement regarding the
	(3) Transform acetic acid to biogas		process in sewage treatment
	(4) Digest fungi in the sludge		(1) Primary treatment is a physical process
Ans.	(1)		(2) Secondary treatment is a biological
Sol.	Monomers are converted into organic acids		process
	by acidogenic bacteria.		(3) The BOD of effluent increases after
			secondary treatment
29.	A part of phgotosynthetically fixed CO2 goes		(4) The tanks used for tertiary treatment are
	back to the atmosphere due to		anaerobic sludge digesters
	(1) Cyclic photophosphorylation	Ans.	(3)
	(2) Non-cyclic photophosphorylation	Sol.	The BOD of eluent decreases aer secondary
	(3) Dark reaction		treatment as bacteria decompose most of the
	(4) Photoresrpiration		organic matter in the eluents.
Ans.	(4)		
Sol.	A part of phgotosynthetically fixed CO2 goes	33.	A type of seed germination in which epicotyl
	back to the atmosphere due to		region of embryo elongates rapidly to pull
	photoresrpiration.		plumule out of the soil surface is called
			germination
30.	When glucose is the normal energy source in		(1) Hypogeal
	bacterial cell gene transcribes a		(2) Epigeal
6	repressor mRNA.		(3) Viviparous



(4) Invitro

Ans. (1)

- Sol. Hypogeal germination is a type of seed germination in which epicotyl region of embryo elongates rapidly and brings the plumule above the soil surface.
- 34. In a dihybrid cross with contrasting characters, the number of different genotypes obtained in F<sub>2</sub> generation are
  - (1) 16
  - (2)9
  - (3)6
  - (4) 4

Ans. (2)

- Sol. Dierent genotypes obtained

  The genotypic ratio is 1:2:2:4:1:2:1:2:1
- **35.** Which one of the following is **NOT** caused by global warming?
  - (1) Melting of glacier
  - (2) Change in rainfall pattern
  - (3) Increase in sea level
  - (4) Increase in crop production

Ans. (4)

- Sol. Global warming causes melting of glacier, change in rainfall pattern and increase in sea level.
- **36.** During cyclic photophosphorylation, formation of ATP occurs between which of the following two compounds?
  - (1) FRS → Ferredoxin
  - (2) Cytochrome b<sub>6</sub> → Cytochrome f
  - (3) Cytochrome  $f \rightarrow Plastocyanin$
  - (4) Plastocyanin → Ionised Chl a

Ans. (2)

Sol. In cyclic photophosphorylation, formation of ATP occurs between

Cytochrome  $b_6 \rightarrow$  Cytochrome fFerredoxin  $\rightarrow$  Cytochrome  $b_6$ 

- **37.** The development of male gametes in the pollen grains in angiosperms involves \_\_\_\_
  - (1) Only one mitotic division
  - (2) Two mitotic divisions
  - (3) Both mitotic and meiotic divisions
  - (4) Only one meiotic divisons

Ans. (1)

- Sol. After pollination, 2 celled pollen grain is deposited on stigma surface and it undergoes germination. When generative cell undergoes mitosis only once for formation of 2 male gametes.
- **38.** In the system of classification, which one of the following is **NOT** a category?
  - (1) Kingdom
  - (2) Series
  - (3) Angiospermae
  - (4) Genus

Ans. (3)

- Sol. Angiospermae is not a category. Angiospermae is a division of Angiosperms.
- 39. Mendel obtained pure lines by selfing yellow round seeds as female parent and green wrinkled as male parent for \_\_\_\_\_ generations.
  - (1) 2
  - (2) 3
  - (3) 4
  - (4) 5

Ans. (2)

Sol. Mendel obtained pure lines by selfing yellow round seeds as female parent and green wrinkled as male parent for 3 generations.



40.	The enzyme Taq polymerase used in PCR		reaction.
	technique is obtained from		
	(1) Salmonella typhimurium		
	(2) Thermus acquaticus	44.	The sequential events that occur during
	(3) Rhizobium phaseoli		protein synthesis are _
	(4) Bacillus thuringiensis		(1) Protein $\rightarrow$ Translation $\rightarrow$ Transcription
Ans.	(2)		DNA
Sol.	The enzyme Taq polymerase used in PCR		(2) DNA $\rightarrow$ Transcription $\rightarrow$ Translation
	technique is obtained		Protein
	from Thermus acquaticus.		(3) Translation → Transcription → DNA
			Protein
41.	Which one of the following is <b>NOT</b> true about		(4) DNA → Protein → Translation
	vegetative propagation		Transcription
	(1) Easy and cheaper method	Ans.	(2)
	(2) Rapid propagation	Sol.	This is central Dogma for protein synthesis.
	(3) Production of genetically similar plants		
	(4) Production of gentically dissimilar plants	45.	The net gain of ATP molecules during
Ans.	(4)		aerobic breakdown of one glucose molecule
Sol.	Vegetative propagation results on production		is
	of gentically similar plants as this is a type		(1) 40
	of sexual reproduction.		(2) 38
			(3) 36
42.	Which one of the following characfteristic is		(4) 30
	NOT shown by a virus?	Ans.	(2)
	(1) They are acellular	Sol.	38 ATP molecules are produced during
	(2) They can be crystallised		breakdown of glucose molecule.
	(3) Active outside the hosts body		
	(4) Have genetic material	46.	The 'eyes' of potato are located at the
Ans.	(3)		(1) Root apex
Sol.	Virus is NOT active outside the host body.		(2) Leaf apex
			(3) Nodes
43.	The co-enzyme which acts as hydrogen		(4) Inter-nodes
	acceoptor during light reaction is	Ans.	(3)
	(1) PQ	Sol.	A stem tuber has many notches on its
	(2) FAD		surface called 'eyes'. Each 'eye' is actually at
	(3) CO-Q		a node and constant of one or more small
	(4) NADP		axillary buds and reduced scale leaves.
Ans.	(4)		

Sol. NADP is the hydrogen acceptor during light



- **47.** Ozone layer in the trophosphere is for plants and animals
  - (1) Good
  - (2) Harmful
  - (3) Providing protection from UV rays
  - (4) Releasing more oxygen

Ans. (2)

- Sol. Ozone layer in stratosphere is provides protection from UV rays while ozone layer in trophosphere is also known as bad ozone because it is harmful to breathe and it damages crops trees and other vegetation. And it also harmful for animals.
- **48.** Steroids are characterised by \_\_\_\_ in their structure
  - (1) Carbon atom arranged in four inter-locking rings
  - (2) Magnesium atom arranged in the centre of tetrapyrole
  - (3) Two, six carbon rings
  - (4) Skeletal heterocyclic ring of hydrocarbons

Ans. (1)

- Sol. Steroids are characterised by carbon atom arranged in four inter-locking rings.
- **49.** Chl-a and Chl-b shown maximum absorption in \_\_\_\_\_ regions of visible light
  - (1) Blue, violet and red
  - (2) Red, indigo and green
  - (3) Yellow, blue and red
  - (4) Blue, violet and green

Ans. (1)

- Sol. Chl-a and Chl-b shown maximum absorption in Blue, violet and red regions of visible light.
- **50.** Explant used for the production of disease free plants is \_\_\_\_\_.

- (1) Vascular tissue
- (2) Apical meristem
- (3) Intrafascicular cambium
- (4) Cork cambium

Ans. (2)

- Sol. Explant used for the production of disease free plants is apical meristem.
- **51.** Explant used for the production of disease free plants is \_\_\_\_\_
  - (1) Vascular tissue
  - (2) Apical meristem
  - (3) Intrafascicular cambium
  - (4) Cork cambium

Ans. (4)

- Sol. Sacculina and crab show parasitism. It is an intra specific interaction in which one of the species (parasite) is benefited while the other one (host) is harmed.
- **52.** Match the Column I and II and select the a propriate option with the given code.

	Column - I		Column - II
(P)	Coronary sinus	(i)	Mitral valve
(Q)	Base of systemic aorta	(ii)	Eustachian valve
(R)	Left atrioventricular valve	(iii)	Semilunar valves
(S)	Opening of inferior vena cava	(iv)	The besian valve

- (1) P iv, Q iii, R i, S ii
- (2) P iii, Q iv, R ii, S i
- (3) P ii, Q i, R iv, S iii
- (4) P i, Q ii, R iii, S iv

Ans. (1)

Sol. Coronary sinus - (iv)

Base of systemic aorta - (iii)

Le atrioventricular valve - (i)

	Opening of inferior vena cava - (ii)	(2) Cytochromes, catalases, peroxidases
		(3) Hyaluronidases, dehydrogenases,
53.	Density of population increases when	hydrolases
		(4) Flavoproteins, anhydrases, phosphatases
	(1) Mortality and emigration increases	Ans. (1)
	(2) Natality and immigration increases	Sol. Co-enzymes of Electron Transport Chain in
	(3) Mortality increases and immigration	mitochondria are cytochromes,
	decreases	dehydrogenases, flavoproteins
	(4) Natality decreases and emigration	
	increases	57. In the second trimester of pregnancy,
Ans.	(2)	there is decline in the level of
Sol.	Density increases when birth rate (Natality)	(1) ACTH
	and Immigration (inflow of population)	(2) ICSH
	increases.	(3) HCG
		(4) MSH
54.	Total number of linkage groups in honeybee	Ans. (3)
	is	Sol. In second trimester, the hormone levels
	(1) 16	stabilizes as HCG declines. Corpus luteum
	(2) 23	deteriorates and placenta completely takes
	(3) 32	over the production of progesterone which
	(4) 46	maintains the pregnancy.
Ans.	(1)	
Sol.	Linkage groups in a species correspond to	
	its haploid number. In honeybees, total chromosomes = 32	58. Heavily built, short prehistoric man
	2	having cranial capacity of about 1450 C.C is
	$\therefore \text{ Linkage groups } = \frac{3}{2}$	
	Haploid no. of chromosome in honeybee = 16	(1) Homo sapiens fossilis
55.	Hormone erythropoietin is produced by	(2) Homo erectus
	(1) Heart	(3) Homo heidelbergensis
	(2) Kidney	(4) Homo neanderthalensis
	(3) Ovary	Ans. (4)
	(4) Pancreas	Sol. Its fossil was invented by J.K. Fuhlort on
Ans.		Neanderthal valley, Germany.
Sol.	Erythropoietin is produced in the kidneys	C.C = 1450  mL
	and plays a key role in production of RBC's.	These were heavily built & short.
56.	Co-enzymes of Electron Transport Chain	<b>59.</b> Ventrolateral walls of cerebrum are called
	in mitochondria are	
	(1) Cytochromes, dehydrogenases, flavoproteins	(1) Crura cerebrii
	es as 1500 to 500 to	(2) Corpora lutea



- (3) Corpora quadrigemina
- (4) Corpora striata


Ans. (4)

Sol. Ventrolateral walls of cerebrum is corpora striata.

- 60.Carbon monoxide combines with haemoglobin t
  - o form \_\_\_\_\_
  - (1) Carbaminohaemoglobin
  - (2) Carbonic acid
  - (3) Carboxyhaemoglobin
  - (4) Cytochrome oxidase

Ans. (3)

- Sol. Carbon monoxide combines with haemoglobin and forms
  Carboxyhaemoglobin. Hb. has more ainity for CO than for oxygen.
- 61. In the given diagram of antibody, 'X' indicates \_



- (1) Constant region of heavy chain
- (2) Antigen binding site
- (3) Disulphide bond
- (4) Light chain


Ans. (2)

- Sol. X is antigen binding site. Antigen combines with the antibody at this site in a lock and key manner.
- **62**. The termination of process of oogenesis

is indicated by formation of \_\_\_\_\_

- (1) Secondary oocyte
- (2) First polar body
- (3) Second polar body
- (4) Perivitelline space

Ans. (3)



**63.** Dr. Lalji Singh obtained radioactive DNA probe from Y chromosome of \_\_\_\_\_

- (1) Female drosophila
- (2) Female banded krait
- (3) Male drosophila
- (4) Male banded krait

Ans. (2)

- Sol. Dr. Lalji Singh obtained radioactive DNA probe from Y chromosome of female banded krait.
- **64.** Which one of the following is NOT the in-situ conservation of biodiversity?
  - (1) Zoological parks
  - (2) Wildlife sanctuaries
  - (3) National parks
  - (4) Biosphere reserves

Ans. (1)

- Sol. Zoological parks are ex-situ conservation of biodiversity.
  - **65.** Which of the following is a unicellular gland?
    - (1) Goblet cell
    - (2) Kupffer's cell
    - (3) Pedicel
    - (4) Neuroglial cell

Ans. (1)

Sol. Unicellular glands are single celled secretory cells. Goblet & mucous cells are major example of unicellular glands in humans.



	In the second se	
66.	In ovary, estrogen is secreted by	flow decreases or stops to a part of the heart
	(1) Follicular cells	causing damage to heart muscle.
	(2) Corona radiata cells	CO M 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	(3) Sertoli cells	<b>69.</b> Molecular evolution and origin of life
200mm20m	(4) Primary oocyte	occurred in Era.
Ans.	(1)	(1) Archeozoic
Sol.		(2) Cenozoic
Ovar	y shows cyclic changes during menstrual	(3) Palaeozoic
	cycle. Cortical region shows dierent stages	(4) Proterozoic
	of development of ovarian follicles or	Ans. (1)
	Graafian follicles. Each follicle has many	Sol. Molecular evolution and origin of Earth was
	layers of follicle cells. The follicle cells of a	observed in Archeozoic Era.
	maturing follicle secrete estrogen.	
		70. The part labelled 'X' in given diagram of
67.	Select the CORRECT statement.	T.S. of adrenal gland secretes
	(1) Sex-linked traits appear more	
	frequently in females	X - HA CONTRACTOR
	(1) Colour blindness is caused due	
	to proper formation of colour	(1) Calcitonin
	sensitive cells	(2) Catecholamines
	(2) Human females have double	(3) Cortisol
	dose of X chromosome	(4) Mineralocorticoids
	(4) Drone bees produce sperms by meiosis	Ans. (2)
Ans.	(3)	Sol. X is adrenal Medulla. It secretes two
Sol.	Human females are XX and hence they show	hormones known as adrenaline
	double dose of X chromosome.	(epinephrine) and noradrenaline (nor
	Sex linked traits appear more frequently in	epinephrine). These are together known as
	males and females are carriers.	Catecholamines.
	Colour blindness is due to improper	
	formation of cones (colour sensitive cells).	71. Given below are cell organelles and their
	Drone bees are infertile males.	functions. Select the INCORRECT match.
68.	Heart attack is also called	(1) Lysosome — Phagocytosis
	(1) Arteriosclerosis	(2) Centriole — Spindle formation
	(2) Angina pectoris	(3)Sphaerosomes —
	(3) Bradycardia	Storage and synthesis of fats
	(4) Myocardial infarction	(4) Leucoplast — Photosynthesis
Ans.	(4)	Ans. (4)
Sol.	Myocardial infarction is commonly known as	Sol. Chloroplast is plastid for Photosynthesis.
		concreptant to place for i notosynthesis.

heart attack. It occurs when blood



Leucoplast is plastid for storage of starch, lipids & proteins.

72.An extra	sex	chromosome	is	present i	in	
-------------	-----	------------	----	-----------	----	--

(1) Turner's syndro


- (1) Turner's syndrome
- (2) Thalassemia
- (3) Klinefelter's syndrome
- (4) Down's syndrome

Ans. (3)

Sol. Klinefelter's syndrome has 44A + XXY one extra X-chromosome is present.

**73.** Given below is the diagram of Animal cell.

Select option giving correct identification of labels X and Y respectively



- (1)X Smooth endoplasmic reticulum, Y Ro ugh endoplasmic reticulum
- (2) X Microvilli, T Golgi apparatus
- (3)X Endoplasmic reticulum, Y Mitochond ria
- (4) X Centrioles, Y Microvilli

Ans. (2)

Sol.  $X \Rightarrow$  Microvilli

Y ⇒ Golgi apparatus

- 74. Parasympathetic ganglia are located \_\_\_\_\_
  - (1) In cortex of cerebrum
  - (2) In medulla of cerebellum
  - (3) On the side of visceral organs
  - (4) On either side of vertebral column

Ans. (3)

Sol. Parasympathetic ganglia are located on the side of visceral organs like heart, kidney etc.

- **75.**One of the following groups of enzymes forms c ontents of succus entericus
  - (1) Maltase, Enterokinase, Trypsin
  - (2) Trypsin, Pepsin, Lactase
  - (3) Nuclease, Amylase, Chymotrypsin
  - (4) Sucrase, Maltase, Dipeptidase

Ans. (4)


- Sol. Succus entericus contains Sucrase, Maltase, Dipeptidase
- **76.** Adam's apple is also called \_\_\_\_
  - (1) Vestibule
  - (2) Pharynx
  - (3) Osteon
  - (4) Larynx

Ans. (4)

Sol. Adam's apple is Larynx (Sound Box)

More prominent in Males

**77.** In the following diagram of T.S. of Artery, the region marked 'X' is



- (1) Tunica media
- (2) Tunica externa
- (3) Tunica interna
- (4) Tunica albuginea

Ans. (3)


Sol. Tunica interna is the innermost layer

- **78.** Which of the following processes is initiated by FSH?
  - (1) Degeneration of endometrium
  - (2) Growth of Graafian follicle
  - (3) Maturation of spermatozoa
  - (4) Secretion progesterone



Ans.	(2)	82.	Which of the following is NOT a connecting
Sol.	FSH is Folicle Stimulating Hormone brings		link?
	about maturation of graafian follicle.		(1) Archeopteryx
			(2) Dryopithecus
79.	The immunity developed by a child after		(3) Ichthyostegia
	attack of measles is immunity		(4) Seymouria
	(1) Natural acquired active	Ans.	(2)
	(2) Artificial acquired active	Sol.	Dryopitheus is not a connecting link
	(3) Natural acquired passive		Archeopteryx is connecting link between
	(4) Artificial acquired passive		birds & Reptils
Ans.	(1)		Ichthyostegia is connecting link between
Sol.	A child aer attack of measles develops		Fish & Amphibian
	Natural Acquired active Immunity. Natural		Seymouria is connecting link between
	Acquired active Immunity is the immunity		Amphobians & Reptils
	acquired due to infection		
		83.	Oogonia arise form the endoderm of the
80.	As an example of adaptive radiation, Darwin		
	observed a variety of Finches in		(1) Allantois
	(1) South Africa		(2) Chorion
	(2) Madagascar		(3) Trophoblast
	(3) Galapagos Islands		(4) Yolk-sac
	(4) East Indies	Ans.	(4)
Ans.	(3)	Sol.	Oogonia arise form the endoderm of the yolk
Sol.	Darwin observed a variety of Finches in		sac and migrate to ovaries during embryonic
	Galapagos Islands		development
81.	Following are various symptoms of	84.	Lacteals absorb
	Marasmus EXCEPT		(1) Amino acids
	(1) Oedema of lower legs and face		(2) Fatty acids and glycerol
	(2) Dry, wrinkled skin		(3) Glucose and fructose
	(3) Extreme leanness		(4) Amylose and maltose
	(4) Atrophy of digestive glands	Ans.	(2)
Ans.		Sol.	Lacteals absorb Fatty acids and glycerol
Sol.	Marasmus is prolonged protein energy		
	malnutrition. Atrophy of digestive glands &	85.	In the given diagram of Malpighian body,
	intestinal mucosa is seen due to which		blood is filtered from part labelled
	absorption of food stops. There is no oedema		
	seen in Marasmus, oedema is a		
	characteristic of kwashiorkor		





- (1) L
- (2) M
- (3) N
- (4) O

Ans. (3)

- Sol. Blood is filtered from Glomerulus. In the given diagram 'N' is the Glomerulus. L-Aerent arteriole, M-eerent arteriole. O-Neck of Bowman's capsule.
- 86. Majority of kidney stones consist crystals of
  - (1) Calcium oxalate, sodium bicarbonate
  - (2) Calcium oxalate, calcium phosphate
  - (3) Calcium pohsphate, sodium chloride
  - (4) Calcium carbonate, copper sulphate

Ans. (2)

- Sol. There are several types of kidney stones based on type of crystals of which they consist. The majority are calcium oxalete stones, followed by calcium phsophate stones
- 87. Each testicular lobule contains \_\_\_\_\_
  - (1) 1 to 4
  - (2) 5 to 7
  - (3) 8 to 10
  - (4) 11 to 14

Ans. (1)

Sol. In each testis, there are 200 - 300 lobules.

In each lobule there are 1-4 convoluted loops called seminiferous tubules.

- **88.** Select the **INCORRECT** statement with reference to DNA fingerprinting. It is \_\_\_\_
  - (1) PCR based technique
  - (2) Based on fingerprints of individual
  - (3) Used in forensic medicine technique
  - (4) Used to test paternity

Ans. (2)

- Sol. DNA fingerprinting is not based on fingerprints of individual
- **89.** Lac is used in the manufacture of following products EXCEPT
  - (1) Toys
  - (2) Polish
  - (3) Isinglass
  - (4) Ink

Ans. (3)

- Sol. Lac is used in the manufacture of Toys, Polish, Ink etc.
- **90.** Which of the following group of animals is guanotelic?
  - (1) Labeo, Turtle, Camel
  - (2) Lizard, Snake, Scorpion
  - (3) Penguin, Spider, Scorpion
  - (4) Spider, Scorpion, Snake

Ans. (3)

- Sol. Spiders, Scorpions, Penguins excrete mostly guanine and hence are called guanotelic
- **91.** Which of the following is an avascular tissue?
  - (1) Connective
  - (2) Epithelial
  - (3) Muscular
  - (4) Nervous

Ans. (2)

Sol. Epithelial tissue lacks blood vessels and hence is an avascular tissue



(1) Arboreal

(3) Cursorial

(2) Aerial

trypsin is used to tre se	eat Ans. Sol.	<ul><li>(4) Fossorial</li><li>(2)</li><li>Pneumatic bones and le sided ovary is characteristic feature of Aerial adaptation</li></ul>
		Pneumatic bones and le sided ovary is
		· ·
ritie		
ritie		characteristic reature of fierial adaptation
itis	96.	Regeneration as a method of asexual
would be useful		reproduction is observed in
		(1) Ascaris
24 0000b W 10 50		(2) Planaria
trypsin. This is used	to	(3) Prawn
		(4) Salmonella
1 41 1 1		(2)
	501.	Regeneration as a type of Asexual
es its maximum in t	ine	Reproduction observed in Planaria
_	97.	Branching tree like processes of white matter
e		extending into grey cortex of cerebellum is
		called
		(1) Arbor vitae
		(2) Gyri
	ım	(3) Crura cerebrii
se		(4) Thalami
C 11 1		(1)
•	sed Sol.	Arbor vitae is seen in grey cortex of
177		cerebellum
	98.	In the given diagram of mammalian bone, X
		indicates
ume		
successfully produc	and	
due to short generati		x
nd left sided ovary	is	(1) Bone marrow
feature	of	(2) Haversian canal
		(3) Inner circumferential lamella
	cycle, the level es its maximum in the exercises are reaches its maximum in the exercises are successfully product the following:  I estrous cycle plantation young ones time  successfully product the exercise are successfully product the exercises are	es its maximum in the  97.  e Ans. successfully produced Sol. due to following: estrous cycle colantation young ones time  successfully produced due to short generation  and left sided ovary is

(4) Volkmann's canal

Ans. (4)



- Sol. Two adjacent Haversian systems are connected by Volkmann's canal
- 99. All of the following are marine Fish EXCEPT
  - (1) Cirrhina
  - (2) Herpedon
  - (3) Sardinella
  - (4) Rastrelliger
- Ans. (1)
- Sol. Cirrhina is fresh water fish
- 100. Heroin is extracted from the latex of plant
  - (1) Atropa belladona
  - (2) Claviceps purpurea
  - (3) Erythroxylum coca
  - (4) Papaver somniferum
- Ans. (4)
- Sol. Heroin is extracted from Papaver Somniferum

