

**CHEMISTRY MARKING SCHEME**

**DELHI - 2013**

**SET 56/1/1**

Q no.	Answers	Marks
1	4	1
2	Mond Process/ Vapour phase refining method	1
3	4	1
4	4-chloropent-1-ene	1
5	CH <sub>3</sub> CN is for n <sub>e</sub> d or ethanenitrile is for n <sub>e</sub> d.	1
6	H <sub>3</sub> C-CH(CH <sub>3</sub> )-CH <sub>2</sub> -CHO	1
7	(CH <sub>3</sub> ) <sub>3</sub> N < CH <sub>3</sub> NH <sub>2</sub> < (CH <sub>3</sub> ) <sub>2</sub> NH	1
8	mRNA, rRNA, tRNA	1
9	$\Delta T_b = K_b \cdot m$ $T_b - T_b^0 = 0.52 \text{ K kg mol}^{-1} \times \frac{18 \text{ g}}{180 \text{ g mol}^{-1}} \times \frac{1}{1 \text{ kg}}$ $T_b - 373.15 \text{ K} = 0.052 \text{ K}$ $T_b = 373.202 \text{ K}$	1/2  1/2  1/2  1/2
10	$\Lambda_m = \kappa / C$ $\Lambda_m = \frac{0.025 \text{ S cm}^{-1}}{0.20 \text{ mol L}^{-1}}$ $\Lambda_m = 125 \text{ S cm}^2 \text{ mol}^{-1}$ <p align="center">(deduct 1/2 mark for wrong or no unit)</p>	1/2  1/2  1






11	<table border="0"> <thead> <tr> <th></th> <th>Dispersed phase</th> <th>Dispersion Medium</th> </tr> </thead> <tbody> <tr> <td>(i)</td> <td>Smoke</td> <td>Gas</td> </tr> <tr> <td>(ii)</td> <td>Milk</td> <td>Liquid</td> </tr> </tbody> </table>		Dispersed phase	Dispersion Medium	(i)	Smoke	Gas	(ii)	Milk	Liquid	1 1	
	Dispersed phase	Dispersion Medium										
(i)	Smoke	Gas										
(ii)	Milk	Liquid										
11	<p style="text-align: center;"><b>OR</b></p> <p><b>Lyophilic sols are solvent attracting sols whereas Lyophobic sols are Solvent repelling sols</b>  <b>Lyophobic sols can be easily coagulated</b></p>	1/2 + 1/2 1										
12	<table border="1"> <thead> <tr> <th>Physisorption</th> <th>Chemisorption</th> </tr> </thead> <tbody> <tr> <td>It is not very specific.</td> <td>It is highly specific.</td> </tr> <tr> <td>It usually takes place at low temperature and decreases with increasing temperature.</td> <td>It takes place at high temperature.</td> </tr> <tr> <td>It is reversible.</td> <td>It is irreversible.</td> </tr> <tr> <td>Low enthalpy of adsorption.</td> <td>High enthalpy of adsorption.</td> </tr> </tbody> </table>	Physisorption	Chemisorption	It is not very specific.	It is highly specific.	It usually takes place at low temperature and decreases with increasing temperature.	It takes place at high temperature.	It is reversible.	It is irreversible.	Low enthalpy of adsorption.	High enthalpy of adsorption.	1/2 x 4 = 2
Physisorption	Chemisorption											
It is not very specific.	It is highly specific.											
It usually takes place at low temperature and decreases with increasing temperature.	It takes place at high temperature.											
It is reversible.	It is irreversible.											
Low enthalpy of adsorption.	High enthalpy of adsorption.											
13	<p>(a) NaCN solution</p> <p>(b) CO</p>	1+1										
14	<p>(i)</p> $\text{PCl}_5 \xrightarrow{\text{heat}} \text{PCl}_3 + \text{Cl}_2$ <p>(ii)</p> $4\text{H}_3\text{PO}_3 \xrightarrow{\text{heat}} 3\text{H}_3\text{PO}_4 + \text{PH}_3$ <p style="text-align: center;">(Full marks may be given if equation is not balanced)</p>	1 1										





15	<p>(a) Cu, because in +1 oxidation state it has stable <math>3d^{10}</math> configuration.</p> <p>(b) <math>Mn^{2+}</math>, <math>V^{3+}</math>: because of the presence of unpaired electrons.</p> <p>(if only one ion is mentioned deduct <math>\frac{1}{2}</math> mark)</p>	<p><math>\frac{1}{2} + \frac{1}{2}</math> <math>\frac{1}{2} + \frac{1}{2}</math></p>
16	<p>(i) Due to resonance / diagrammatic representation, C-Cl bond acquires a partial double bond character which is difficult to cleave.</p> <p>(ii) Due to <math>sp^2</math> hybridisation of 'C' of C-Cl bond.</p> <p>(iii) Due to unstable phenyl cation</p> <p>(iv) Due to repulsion between nucleophile and electron rich arenes.</p> <p>(any two)</p>	1+1
17	<p>(i) <math>CH_3-CH_2-\ddot{O}-H + H^+ \rightarrow CH_3-CH_2-\overset{+}{O}-H</math></p> <p>(ii) <math>CH_3CH_2-\ddot{O}: + CH_3-CH_2-\overset{+}{O}-H \rightarrow CH_3CH_2-\overset{+}{O}-CH_2CH_3 + H_2O</math></p> <p>(iii) <math>CH_3CH_2-\overset{+}{O}-CH_2CH_3 \rightarrow CH_3CH_2-O-CH_2CH_3 + H^+</math></p>	<p><math>\frac{1}{2}</math> <math>\frac{1}{2}</math> 1</p>
18	<p>(i)</p> $CH_3-CH=CH_2 \xrightarrow{H_2O/H^+} CH_3-\underset{\substack{  \\ OH}}{CH}-CH_3$ <p>(ii)</p>  <p>(or by any other correct suitable method)</p>	1+1






19	<p>(a) p-type semiconductor          (b) Ferromagnetism          (c) Impurity defect / Cation vacancy defect</p>	1x3=3
20	<p>When <math>K_2SO_4</math> is dissolved in water, ions are produced          Total number of ions produced = 3</p> $i = 3$ $\pi = i CRT = i \times \frac{n}{V} \times R \times T$ $\pi = 3 \times \frac{2.5 \times 10^2 \text{ g}}{174 \text{ g mol}^{-1}} \times \frac{1}{2L} \times 0.0821 \text{ Lat mK}^1 \text{ mol}^{-1} \times 298 \text{ K}$ $\pi = 5.27 \times 10^3 \text{ atm}$ <p style="text-align: right;">(deduct 1/2 mark for wrong or no unit)</p>	<p>1/2          1/2          1          1</p>
21	<p>The cell reaction: <math>Fe(s) + 2H^+(aq) \rightarrow Fe^{2+}(aq) + H_2(g)</math></p> <p><math>E_{cell} = 0.44 \text{ V}</math></p> <p>Nernst equation</p> $E_{cell} = E_{cell}^{\circ} - \frac{0.059}{2} \log \frac{[Fe^{2+}]}{[H^+]^2}$ $E_{cell} = 0.44 \text{ V} - \frac{0.059}{2} \log \left( \frac{0.001 \text{ M}}{(1 \text{ M})^2} \right)$ $= 0.44 \text{ V} - \frac{0.059}{2} \log (10^{-3})$ $= 0.44 \text{ V} + 0.0885 \text{ V}$ $= 0.5285 \text{ V}$ <p style="text-align: right;">(deduct 1/2 mark for wrong or no unit)</p>	<p>1          1/2          1/2          1</p>





22	<p>(i) Due to incomplete filling of d-orbitals, transition metals show variable oxidation states.</p> <p>(ii) Because of Lanthanoid Contraction.</p> <p>(iii) Because of their ability to show multiple / variable oxidation states.</p> <p style="text-align: center;">OR</p>	1 x 3=3
22	<p>(i) <math>\text{Cr}_2\text{O}_7^{2-} + 6\text{Fe}^{2+} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 6\text{Fe}^{3+} + 7\text{H}_2\text{O}</math></p> <p>(ii) <math>2\text{CrO}_4^{2-} + 2\text{H}^+ \rightarrow \text{Cr}_2\text{O}_7^{2-} + \text{H}_2\text{O}</math></p> <p>(iii) <math>2\text{MnO}_4^- + 5\text{C}_2\text{O}_4^{2-} + 16\text{H}^+ \rightarrow 2\text{Mn}^{2+} + 10\text{CO}_2 + 8\text{H}_2\text{O}</math></p> <p style="text-align: right;">(Accept only balanced equation)</p>	1 x 3=3
23	<p>(i) Triamminetri chloridochromium(III)</p> <p>(ii) Potassium hexacyanoferrate(III)</p> <p>(iii) Dibromodibis-(ethane-1,2-diamine)cobalt(III) / Dibromodibis-(ethylenediamine)cobalt(III)</p>	1 1 1
24	<p>(i) A=C<sub>6</sub>H<sub>5</sub>CN                      B=C<sub>6</sub>H<sub>5</sub>COOH                      C=C<sub>6</sub>H<sub>5</sub>CONH<sub>2</sub></p> <p>(ii) A=C<sub>6</sub>H<sub>5</sub>NH<sub>2</sub>                      B=C<sub>6</sub>H<sub>5</sub>N<sub>2</sub><sup>+</sup>Cl<sup>-</sup>                      C=C<sub>6</sub>H<sub>5</sub>-OH</p>	$\frac{1}{2} \times 3 = 1 \frac{1}{2}$ $\frac{1}{2} \times 3 = 1 \frac{1}{2}$
25	<p>(i) Buna-S: 1,3-Butadiene and Styrene</p> <p style="text-align: center;"><math>\text{CH}_2 = \text{CH} - \text{CH} = \text{CH}_2</math> and </p> <p>(ii) Neoprene: Chloroprene</p> <p style="text-align: center;"><math>\text{CH}_2 = \overset{\text{Cl}}{\underset{ }{\text{C}}} - \text{CH} = \text{CH}_2</math></p>	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$











28	<p>(a)</p> <p>(i) rate = <math>k[A]^2[B]</math></p> <p>(ii) Rate will increase 9 times of the actual rate of reaction</p> <p>(iii) Rate will increase 8 times of the actual rate of reaction</p> <p>(b)</p> $k = \frac{2.303}{t} \log \frac{[A]_0}{[A]}$ $k = \frac{2.303}{40 \text{ min}} \log \frac{100}{70}$ $k = \frac{2.303}{40} \times 0.155 = 0.00892 \text{ min}^{-1}$ $t_{1/2} = \frac{0.693}{k}$ $t_{1/2} = \frac{0.693}{0.00892} \text{ min}$ $t_{1/2} = 77.7 \text{ min}$	<p>1x3=3</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>
28	<p style="text-align: center;">OR</p> <p>(a)</p> $t_{99\%} = \frac{2.303}{k} \log \frac{100}{1}$ $t_{90\%} = \frac{2.303}{k} \log \frac{100}{10}$ <p>on comparison</p> $\frac{t_{99\%}}{t_{90\%}} = \frac{\log 100}{\log 10}$ <p>Hence <math>t_{99\%} = 2 t_{90\%}</math></p> <p style="text-align: center;">(or solved by any other correct suitable method)</p>	<p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>






	<p>(b)</p> $\text{Slope} = -\frac{E_a}{2303R}$ $-4250 \text{ K} = -\frac{E_a}{2303 \times 8314 \text{ J K}^{-1} \text{ mol}^{-1}}$ $E_a = 81375 \text{ J mol}^{-1} \text{ or } 81.375 \text{ kJ mol}^{-1}$	<p>1</p> <p>1</p> <p>1</p>
29.	<p>(i) Because of smaller size of F atom/ shorter bond length, the electron-electron repulsion among the lone pairs is greater in <math>\text{F}_2</math> than <math>\text{Cl}_2</math></p> <p>(ii) Due to hydrogen bonding in <math>\text{NH}_3</math>.</p> <p>(b)</p> <p>(i)</p>  <p>(ii)</p> 	1+1





(iii)




OR


- (a) (i) Because of its low solubility in blood.  
 (ii) Because of its highest electronegativity.  
 (iii) Because O-O single bond is weaker than S-S single bond.

(b)

(i)



(ii)




1x3=  
3

1x3=  
3

1+1

29





collegedunia.com  
 India's largest Student Review Platform









	<p>(iii)</p>  <p>(b)</p> <p>(i) <b>Ethanal and Propanal</b> : Ethanal gives yellow ppt of Iodoform (<math>\text{CHI}_3</math>) on addition of <math>\text{NaOH}/\text{I}_2</math> whereas Propanal does not give this test. ( or any other suitable test)</p> <p>(ii) <b>Benzoic acid and Phenol</b> : Add neutral <math>\text{FeCl}_3</math> to both, phenol gives purple / violet colouration whereas Benzoic acid does not give this test or / Add <math>\text{NaHCO}_3</math> to both, Benzoic acid will give brisk effervescence whereas phenol does not give this test. ( or any other suitable test)</p>	<p>1 x3=3</p> <p>1+1</p>
	<p>Sh. S K Murj al Pr of. R D Shuk l a Dr. K N Uppadhya Mr. Rakesh Dhawan Ms. Neer u Sof at Mr. Vrendra S ngh</p>	<p>Dr ( Ms.) Sangeet a Bhati a M. K M Abdul Raheem M. D A Mishra M. Deshbir S ngh M. Akhileshwar Mishra</p>

