

1. If $f(x) = xe^{x(1-x)}$, $x \in \mathbb{R}$, then $f(x)$ is
 - (a) Decreasing on $(-\frac{1}{2}, 1]$
 - (b) Decreasing on $(\frac{1}{2}, 1)$
 - (c) Decreasing on $(\frac{1}{2}, 1]$
 - (d) None of the above
2. $\int (\frac{1}{x} \tan(\frac{1}{x}))^2 dx$ is equal to
 - (a) $x - \tan(x) + C$
 - (b) $\frac{1}{x} + \tan(\frac{1}{x}) + C$
 - (c) $\frac{1}{x} - \tan(\frac{1}{x}) + C$
 - (d) None of the above
3. If $\log_{10}(x^3 + y^3) - \log_{10}(x^2 + y^2 - xy) \leq 2$, then the maximum value of xy for all $x \geq 0, y > 0$, is
 - (a) 2500
 - (b) 3000
 - (c) 1200
 - (d) None of the above
4. If $f(x) = ax + b$ and $f^{-1}(x) = bx + a$, with $a, b, x \in \mathbb{R}$, then what is the value of $a + b$?
 - (a) -2
 - (b) -1
 - (c) 0
 - (d) 1
5. If, $\ln(a + c), \ln(c - a), \ln(a - 2b + c)$ are in Arithmetic Progression, then
 - (a) a, b, c are in Arithmetic Progression
 - (b) a, b, c are in Geometric Progression
 - (c) a, b, c are in Harmonic Progression
 - (d) None of the above

6. Let $x > 0$ and $\log_2 x + \log_2 \sqrt{x} + \log_2 \sqrt[4]{x} + \log_2 \sqrt[8]{x} + \dots = 4$. Then x is equal to
- 2
 - 3
 - 4
 - None of the above
7. A fair coin is tossed n times. If the probability that head occurs 6 times is equal to the probability that head occurs 8 times, then the value of n is
- 14
 - 16
 - 24
 - None of the above
8. If $A = \begin{bmatrix} 2 & 1 \\ -4 & -2 \end{bmatrix}$ and $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then $I + 2A + 3A^2 + \dots$ is equal to
- $\begin{bmatrix} 4 & 1 \\ -4 & 0 \end{bmatrix}$
 - $\begin{bmatrix} 3 & 1 \\ -4 & 1 \end{bmatrix}$
 - $\begin{bmatrix} 5 & 2 \\ -8 & -3 \end{bmatrix}$
 - $\begin{bmatrix} 5 & 2 \\ -3 & -8 \end{bmatrix}$
9. $\lim_{x \rightarrow \frac{\pi}{4}} [\tan(x)]^{\tan(2x)}$ is equal to
- 1
 - e
 - 1
 - None of the above

10. The sum of all squared numbers between 50 and 500 is
- (a) 3704
 - (b) 3655
 - (c) 4233
 - (d) None of the above
11. Coefficient of x^{99} in the expansion of $(x + 1)(x + 3)(x + 5)\dots(x + 199)$ is equal to
- (a) 10250
 - (b) 10000
 - (c) 10500
 - (d) None of the above
12. If $N = n!$, where n is a natural number with $n > 2$, then
- $$\lim_{N \rightarrow \infty} [\log_2 N]^{-1} + [\log_3 N]^{-1} + [\log_4 N]^{-1} + \dots + [\log_n N]^{-1}$$
- is,
- (a) 1
 - (b) 2
 - (c) 3
 - (d) None of the above
13. The final score in a recreational soccer game between Team A and Team B, is 6 goals for A to 3 goals for B. How many possibilities exist for the score at the end of first half?
- (a) 20
 - (b) 24
 - (c) 28
 - (d) None of the above

14. Integers a, b, c and d , not necessarily distinct, are chosen independently and at random from 0 to 2007 (both inclusive). What is the probability that $ad - bc$ is even?

- (a) $\frac{3}{8}$
- (b) $\frac{7}{16}$
- (c) $\frac{9}{16}$
- (d) $\frac{5}{8}$

15. If the function f satisfies the relation $f(x+y) = f(x)f(y)$ for all $x, y \in \mathbb{N}$. Further if $f(1) = 2$ and $\sum_{k=1}^n f(a+k) = 16(2^n - 1)$, then value of a , (where $a \in \mathbb{N}$), is equal to

- (a) 3
- (b) 1
- (c) 2
- (d) 4

16. If for any real number y , $[y]$ is the greatest integer less than or equal to y , then the value of the integral $\int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} [2 \sin x] dx$, is

- (a) $-\pi$
- (b) 0
- (c) $-\frac{\pi}{2}$
- (d) None of the above

17. The value of real number x for which the matrix $\begin{bmatrix} 6 & 4 & 2x \\ 8 & 2 & -2 \\ 0 & 6 & 8 \end{bmatrix}$ has no inverse is

- (a) $\frac{12}{11}$
- (b) $\frac{11}{12}$
- (c) 1
- (d) 0

18. The sum of the infinite series

$$\frac{5}{13} + \frac{55}{13^2} + \frac{555}{13^3} + \dots$$

is equal to

- (a) $\frac{31}{18}$
 - (b) $\frac{65}{32}$
 - (c) $\frac{65}{36}$
 - (d) $\frac{75}{36}$
19. Water is being poured at the rate of 2 cubic metres per second into a cone which has semi vertical angle of 45° . The rate at which perimeter of water surface changes when the height of water in the cone is 2 metres is
- (a) 2 metres per second
 - (b) 1 metre per second
 - (c) 3 metres per second
 - (d) 4 metres per second
20. The number of real solutions for $x^2 + 5|x| + 6 = 0$ is
- (a) 0
 - (b) 2
 - (c) 3
 - (d) 4
21. $\lim_{x \rightarrow \infty} \left(1 - \frac{4}{x-1}\right)^{3x-1}$ is
- (a) e^{-12}
 - (b) e^{12}
 - (c) e^{-4}
 - (d) e^{-3}

22. Let $I = \int_0^1 \frac{\sin x}{\sqrt{x}} dx$ and $J = \int_0^1 \frac{\cos x}{\sqrt{x}} dx$, then which of the following is true?
- (a) $I < \frac{2}{3}$ and $J > 2$
 (b) $I > \frac{2}{3}$ and $J < 2$
 (c) $I > \frac{2}{3}$ and $J > 2$
 (d) $I < \frac{2}{3}$ and $J < 2$

23. The sum of the infinite series

$$1 + \frac{1}{4 \times 2!} + \frac{1}{16 \times 4!} + \frac{1}{64 \times 6!} + \dots$$

is

- (a) $\frac{e-1}{2\sqrt{e}}$
 (b) $\frac{e+1}{2\sqrt{e}}$
 (c) $\frac{e-1}{\sqrt{e}}$
 (d) $\frac{e+1}{\sqrt{e}}$
24. A traffic light runs repeatedly through the following cycle: green for 30 seconds, then yellow for 3 seconds, and then red for 30 seconds. Jack picks a random three - second time interval to watch the light. What is the probability that the color changes while he is watching?
- (a) $\frac{1}{3}$
 (b) $\frac{1}{7}$
 (c) $\frac{1}{10}$
 (d) None of the above

25. The term independent of x in the binomial expansion of

$$\left(\frac{x+1}{x^{\frac{2}{3}} - x^{\frac{1}{3}} + 1} - \frac{x-1}{x - x^{\frac{1}{2}}} \right)^{10}$$

is

- (a) 4
 (b) 120
 (c) 210
 (d) 310

26. $\lim_{n \rightarrow \infty} \left(\frac{n!}{n^n} \right)^{\frac{1}{n}}$ is equal to
- e
 - $\frac{1}{e}$
 - $\frac{\pi}{4}$
 - $\frac{4}{\pi}$
27. Largest possible area of a right angled triangle having hypotenuse of length 4cm is equal to
- 3 sq cm
 - 4 sq cm
 - 5 sq cm
 - None of the above
28. Let f be a one-to-one function with domain $\{x, y, z\}$ and range $\{1, 2, 3\}$. It is given that exactly one of the following statements is true and the remaining two are false: $f(x) = 1$, $f(y) \neq 1$ and $f(z) \neq 2$. Then $f^{-1}(1)$ is equal to
- x
 - y
 - z
 - None of the above
29. If $f(x) = ae^{2x} + be^x + cx$ satisfies the conditions $f(0) = 1$, $f'(\log 2) = 31$ and $\int_0^{\log 4} (f(x) - cx) dx = \frac{39}{2}$, then
- $a = 5, b = 6, c = 3$
 - $a = 5, b = -6, c = 3$
 - $a = -5, b = 6, c = 3$
 - None of the above
30. If x_1, x_2, x_3 and x_4 are the roots of the equation
- $$x^4 - x^3 \sin 2\beta + x^2 \cos 2\beta - x \cos \beta - \sin \beta = 0$$
- then $\tan^{-1}(x_1) + \tan^{-1}(x_2) - \tan^{-1}(x_3) - \tan^{-1}(x_4)$ is equal to
- β
 - $\frac{\pi}{2} - \beta$
 - $\pi - \beta$
 - None of the above