2013 - MA

Test Paper Code: MA

Time: 3 Hours Maximum Marks: 100

INSTRUCTIONS

- 1. This question-cum-answer booklet has 32 pages and has 30 questions. Please ensure that the copy of the question-cum-answer booklet you have received contains all the questions.
- 2. Write your **Registration Number**, Name and the name of the Test Centre in the appropriate space provided on the right side.
- 3. Write the answers to the objective questions against each Question No. in the **Answer Table for Objective Questions**, provided on Page No. 4. Do not write anything else on this page.
- 4. Each objective question has 4 choices for its answer: (A), (B), (C) and (D). Only ONE of them is the correct answer. There will be negative marking for wrong answers to objective questions. The following marking scheme for objective questions shall be used:
 - (a) For each correct answer, you will be awarded 2 (Two) marks.
 - (b) For each wrong answer, you will be awarded **-0.5** (Negative 0.5) mark.
 - (c) Multiple answers to a question will be treated as a wrong answer.
 - (d) For each un-attempted question, you will be awarded **0** (**Zero**) mark.
 - (e) Negative marks for objective part will be carried over to total mark.
- Answer the fill in the blank type and descriptive type questions only in the space provided after each question. No negative marks for fill in the blank type questions.
- 6. Do not write more than one answer for the same question. In case you attempt a fill in the blank or a descriptive question more than once, please cancel the answer(s) you consider wrong. Otherwise, the answer appearing last only will be evaluated.
- All answers must be written in blue/black/ blue-black ink only. Sketch pen, pencil or ink of any other colour should not be used.
- 8. All rough work should be done in the space provided and scored out finally.
- 9. No supplementary sheets will be provided to the candidates.
- 10. Clip board, log tables, slide rule, cellular phone and electronic gadgets in any form are NOT allowed. Non Programmable Calculator is allowed.
- 11. The question-cum-answer booklet must be returned in its entirety to the Invigilator before leaving the examination hall. Do not remove any page from this booklet.
- 12.Refer to special instructions/useful data on the reverse.



2013 - MA

READ INSTRUCTIONS ON THE LEFT SIDE OF THIS PAGE CAREFULLY

	REG	ISTRA	TION	NUM	BER	
Nam	<u></u> е:				<u></u>	1
Test	Centre	•				
1 001	Contro	•				
			1 - V - WAS MAN	**	115 (T. J.)	

Do not write your Registration Number or Name anywhere else in this question-cum-answer booklet.

I	have	read	all	the	instructions	and	shall
al	bide b	y then	n.				

Signature of the Candidate

I have verified the information filled by the Candidate above.

Signature of the Invigilator





	Special Instructions/ Useful Data				
$\mathbb R$: The set of all real numbers				
\mathbb{N}	: The set of all positive integers				
f'	: First derivative of a real function f of single variable				
\mathbb{Z}_p	: $\{0,1,,p-1\}$ with addition and multiplication modulo p				
S^o	: Interior of a set $S \subseteq \mathbb{R}$				
$\overline{\overline{S}}$: Closure of a set $S \subseteq \mathbb{R}$				



IMPORTANT NOTE FOR CANDIDATES

- Questions 1-10 (objective questions) carry two marks each, questions 11-20 (fill in the blank questions) carry three marks each and questions 21-30 (descriptive questions) carry five marks each.
- The marking scheme for the objective type question, is as follows:
 - (a) For each correct answer, you will be awarded 2 (Two) marks.
 - (b) For each wrong answer, you will be awarded **-0.5** (Negative **0.5**) mark.
 - (c) Multiple answers to a question will be treated as a wrong answer.
 - (d) For each un-attempted question, you will be awarded **0** (Zero) mark.
 - (e) Negative marks for objective part will be carried over to total marks.
- There is no negative marking for fill in the blank questions.
- Write the answers to the objective questions in the Answer Table for Objective Questions provided on page 4 only.

Objective Questions

Q.1		1 1 1 -1 1 5 3		vector space of all	$X \in \mathbb{R}^3$ such that	AX = 0.7	Then $\dim(V)$
	is						
	$(\Delta) = 0$		(B) 1	(C)	2	(D) 3	

The value of n for which the divergence of the function $\vec{F} = \frac{\vec{r}}{|\vec{r}|^n}$, $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$, $|\vec{r}| \neq 0$, Q.2 vanishes is

(B)
$$-1$$

(D)
$$-3$$

Q.3 Let A and B be subsets of \mathbb{R} . Which of the following is **NOT** necessarily true?

(A)
$$(A \cap B)^o \subseteq A^o \cap B^o$$

(B)
$$A^{o} \cup B^{o} \subseteq (A \cup B)^{o}$$

(D) $\overline{A} \cap \overline{B} \subseteq \overline{A \cap B}$

(C)
$$\overline{A} \cup \overline{B} \subseteq \overline{A \cup B}$$

(D)
$$\overline{A} \cap \overline{B} \subseteq \overline{A \cap B}$$

Let [x] denote the greatest integer function of x. The value of α for which the function Q.4

$$f(x) = \begin{cases} \frac{\sin[-x^2]}{[-x^2]}, & x \neq 0\\ \alpha, & x = 0 \end{cases}$$

is continuous at x=0 is

(B)
$$\sin(-1)$$

MA-1/32



Q.5 Let the function f(x) be defined by

$$f(x) = \begin{cases} e^x, & x \text{ is rational} \\ e^{1-x}, & x \text{ is irrational} \end{cases}$$

for x in (0, 1). Then

- (A) f is continuous at every point in (0, 1)
- (B) f is discontinuous at every point in (0, 1)
- (C) f is discontinuous only at one point in (0, 1)
- (D) f is continuous only at one point in (0, 1)
- Q.6 The value of the integral

$$\iint_{D} \sqrt{x^{2} + y^{2}} \, dx \, dy \,, \qquad D = \left\{ (x, y) \in \mathbb{R}^{2} : \ x \le x^{2} + y^{2} \le 2x \right\}$$

is

- (A) 0
- (B) $\frac{7}{9}$

- (C) $\frac{14}{9}$
- (D) $\frac{28}{9}$

Q.7 Let

$$x_n = \left(1 - \frac{1}{3}\right)^2 \left(1 - \frac{1}{6}\right)^2 \left(1 - \frac{1}{10}\right)^2 \dots \left(1 - \frac{1}{\frac{n(n+1)}{2}}\right)^2, \quad n \ge 2.$$

Then $\lim_{n\to\infty} x_n$ is

- (A) $\frac{1}{3}$
- (B) $\frac{1}{9}$

- (C) $\frac{1}{81}$
- (D) 0
- Q.8 Let p be a prime number. Let G be the group of all 2×2 matrices over \mathbb{Z}_p with determinant 1 under matrix multiplication. Then the order of G is
 - (A) (p-1)p(p+1)

(B) $p^2(p-1)$

(C) p^3

(D) $p^2(p-1)+p$

Q.9 Let V be the vector space of all 2×2 matrices over \mathbb{R} . Consider the subspaces $W_1 = \left\{ \begin{pmatrix} a & -a \\ c & d \end{pmatrix} : a, c, d \in \mathbb{R} \right\}$ and $W_2 = \left\{ \begin{pmatrix} a & b \\ -a & d \end{pmatrix} : a, b, d \in \mathbb{R} \right\}$.

If $m = \dim(W_1 \cap W_2)$ and $n = \dim(W_1 + W_2)$, then the pair (m, n) is

- (A) (2,3)
- (B) (2, 4)
- (C) (3, 4)
- (D) (1, 3)
- Q.10 Let \mathcal{O}_n be the real vector space of all polynomials of degree at most n. Let $D: \mathcal{O}_n \to \mathcal{O}_{n-1}$ and $T: \mathcal{O}_n \to \mathcal{O}_{n+1}$ be the linear transformations defined by

$$D(a_0 + a_1x + a_2x^2 + \dots + a_nx^n) = a_1 + 2a_2x + \dots + na_nx^{n-1},$$

$$T(a_0 + a_1x + a_2x^2 + \dots + a_nx^n) = a_0x + a_1x^2 + a_2x^3 + \dots + a_nx^{n+1},$$

respectively. If A is the matrix representation of the transformation $DT-TD: \mathcal{O}_n \to \mathcal{O}_n$ with respect to the standard basis of \mathcal{O}_n , then the trace of A is

- (A) -n
- (B) n

- (C) n+1
- (D) -(n+1)

Answer Table for Objective Questions

Write the Code of your chosen answer only in the 'Answer' column against each Question Number. Do not write anything else on this page.

Question Number	Answer	Do not write in this column
01	, , , , , , , , , , , , , , , , , , , ,	
02		
03		
04		
05		
06		
07		
08		
09		
10		

FOR EVALUATION ONLY

Number of Correct Answers	Marks (+)
Number of Incorrect Answers	Marks (-)
Total Marks in Question	n Nos. 1-10 ()

Fill in the blank questions

Q.11 The equation of the curve satisfying $\sin y \frac{dy}{dx} = \cos y (1 - x \cos y)$ and passing through the origin is

Ans:

Q.12 Let f be a continuously differentiable function such that $\int_{0}^{2x^2} f(t) dt = e^{\cos x^2}$ for all $x \in (0, \infty)$. The value of $f'(\pi)$ is

Ans:

Q.13 Let $u = \frac{y^2 - x^2}{x^2 y^2}$, $v = \frac{z^2 - y^2}{y^2 z^2}$ for $x \neq 0$, $y \neq 0$, $z \neq 0$. Let w = f(u, v), where f is a real valued function defined on \mathbb{R}^2 having continuous first order partial derivatives. The value of $x^3 \frac{\partial w}{\partial x} + y^3 \frac{\partial w}{\partial y} + z^3 \frac{\partial w}{\partial z}$ at the point (1, 2, 3) is

Ans:

Q.14	The set of points at which the function	$f(x, y) = x^4 + y^4 - x^2 - y^2 + 1, (x, y) \in \mathbb{R}^2$	attains local
	maximum is		

Ans:

Q.15 Let C be the boundary of the region in the first quadrant bounded by $y=1-x^2$, x=0 and y=0, oriented counter-clockwise. The value of $\oint_C (xy^2 dx - x^2 y dy)$ is

Ans:

Q.16 Let $f(x) = \begin{cases} 0, & -1 \le x \le 0 \\ x^4, & 0 < x \le 1 \end{cases}$. If $f(x) = \sum_{k=0}^{n} \frac{f^{(k)}(0)}{k!} x^k + \frac{f^{(n+1)}(\xi)}{(n+1)!} x^{n+1}$ is the Taylor's formula for f about x = 0 with maximum possible value of n, then the value of ξ for $0 < x \le 1$ is

Ans:

Q.17 Let $\vec{F} = 2z\hat{i} + 4x\hat{j} + 5y\hat{k}$, and let C be the curve of intersection of the plane z = x + 4 and the cylinder $x^2 + y^2 = 4$, oriented counter-clockwise. The value of $\oint_C \vec{F} \cdot d\vec{r}$ is

Ans:

Q.18 Let f and g be the functions from $\mathbb{R}\setminus\{0,1\}$ to \mathbb{R} defined by $f(x) = \frac{1}{x}$ and $g(x) = \frac{x-1}{x}$ for $x \in \mathbb{R}\setminus\{0,1\}$. The smallest group of functions from $\mathbb{R}\setminus\{0,1\}$ to \mathbb{R} containing f and g under composition of functions is isomorphic to

Ans:

Q.19 The orthogonal trajectory of the family of curves $\frac{x^2}{2} + y^2 = c$, which passes through (1, 1) is

Ans:

Q.20 The function to which the power series $\sum_{n=1}^{\infty} (-1)^{n+1} n x^{2n-2}$ converges is

Ans:



Descriptive questions

Q.21 Let $0 < a \le 1$, $s_1 = \frac{a}{2}$ and for $n \in \mathbb{N}$, let $s_{n+1} = \frac{1}{2}(s_n^2 + a)$. Show that the sequence $\{s_n\}$ is convergent, and find its limit.





Q.22 Evaluate

$$\int_{1/4}^{1} \int_{\sqrt{x-x^2}}^{\sqrt{x}} \frac{x^2 - y^2}{x^2} \, dy \, dx$$

by changing the order of integration.

answer
the
for
pace
$ \sqrt{2} $

		1 1 1 1
		1 1 1 1

MA-10/32		

Space for the answer

MA-11/32




Q.23 Find the general solution of the differential equation

$$x^{2} \frac{d^{3}y}{dx^{3}} + x \frac{d^{2}y}{dx^{2}} - 6 \frac{dy}{dx} + 6 \frac{y}{x} = \frac{x \ln x + 1}{x^{2}}, \quad x > 0.$$

answer
the
; for tl
Space
Ö

MA-12/32





Q.24 Let S_1 be the hemisphere $x^2 + y^2 + z^2 = 1$, z > 0 and S_2 be the closed disc $x^2 + y^2 \le 1$ in the xy plane. Using Gauss' divergence theorem, evaluate $\iint_S \vec{F} \cdot d\vec{S}$, where $\vec{F} = z^2 x \hat{i} + \left(\frac{y^3}{3} + \tan z\right) \hat{j} + (x^2 z + y^2) \hat{k}$ and $S = S_1 \cup S_2$.

Also evaluate $\iint_{S_1} \vec{F} \cdot d\vec{S}$.

1	N.
Ţ	7

1			•	
			•	
1				
1				
1				
1				
1				
1				
1				
1				
1				
1				
1				
•				
į				
1				
1				
•				
1				
}				
!				
1				
1				
		•		
1				
1				
Ì				
•				
İ				
•				
i				
i				·
•				
i				
<u> </u>				
i				
!				
•				
1				
!				
!				
!				
i !				
i !				
!				
•				

Space for the answer

MA-15/32



$$f(x,y) = \begin{cases} \frac{2(x^3 + y^3)}{x^2 + 2y}, & (x,y) \neq (0,0) \\ 0, & (x,y) = (0,0). \end{cases}$$

Show that the first order partial derivatives of f with respect to x and y exist at (0,0). Also show that f is not continuous at (0,0).

M	Δ	_1	G/	39

A

	r
	1 1 1 1
	1 1 1 1
}	
}	
	1 1 1 1
;	
· · · · · · · · · · · · · · · · · · ·	
)	
	1 1 1
	; ; ;



A	
\mathbf{A}	

; t				
; ; ;				
; ; ; ;				
i 1 1				
; t ; ;				
† ! !				
i ! ! !				
1 1 1 4 3				
! ! !				
i ! ! !				
: ! ! !				
; ; ; ; ;				
1 1 1 1				
• • • • • • • • • • • • • • • • • • •				
; ; ; ; ;				
1 1 1 1 1				
: 				
1 1 1				

MA-19/32



Q.27	Let A be an $n \times n$ real symmetric matrix with n distinct eigenvalues. Prove that there exists an orthogonal matrix P such that $AP = PD$, where D is a real diagonal matrix.
Space for the answer	









	A
Space for the answer	





Q.29 Let $f:[a,b]\to\mathbb{R}$ be a continuous function such that f is differentiable in (a,c) and (c,b), a < c < b. If $\lim_{x \to c} f'(x)$ exists, then prove that f is differentiable at c and $f'(c) = \lim_{x \to c} f'(x)$. Space for the answer







Γ	A
	\mathbf{A}

	į	
	i e e e e e e e e e e e e e e e e e e e	
	i e e e e e e e e e e e e e e e e e e e	
	i i	
	!	
	<u>;</u>	
	I and the second	
	: 1	
		- 1
	<u> </u>	- 1
		į
	i 	į
	;	
		- 1
		-
		-
		1
		i
D D		i
⋛	i 1	ij
23	1 1	÷
Space for the answer		:
5)		į
Ĕ		i
		:
≍		÷
≍		!
บ		į
2		į
		ì
<u> </u>		÷
-		į
		i
		İ
		1
		:
		į
		Ť
		;
	i de la companya de la companya de la companya de la companya de la companya de la companya de la companya de	į
		ì
		į
		;
		i
		i
		í
		:
	-	1
		į
		i
		;
- 1		
		į.
į		į
į		i
1	· · · · · · · · · · · · · · · · · · ·	1
		į
:		i
į	· · · · · · · · · · · · · · · · · · ·	i
ì	· · · · · · · · · · · · · · · · · · ·	1
ĺ	i i i i i i i i i i i i i i i i i i i	!
ļ	· · · · · · · · · · · · · · · · · · ·	4
ļ		i
•		1
į	i i i i i i i i i i i i i i i i i i i	1
Ì	i de la companya de la companya de la companya de la companya de la companya de la companya de la companya de	į
		į.







A





A



Object	3 - MA tive Part umber 1 – 10)
Total Marks	Signature

Question Number	Marks	Question Number	Marks	
11		21		
12		22	JAV.	
13		23		
14		24		
15		25		
16		26		
17		27		
18		28		
19		29		
20		30		

·	
Total (Objective Part)	:
Total (Fill in the blanks Part and Descriptive Part)	:
Grand Total	:
Total Marks (in words)	:
Signature of Examiner(s)	:
Signature of Head Examiner(s)	
Signature of Scrutinizer	
Signature of Chief Scrutinizer	:
Signature of Coordinating Head Examiner	:

