

DU MSc Physics

Topic:- PHY MSC S2

1) An atomic transition line with wavelength 350 nm is observed to be split into three components in a spectrum of light from a sunspot. Adjacent components are separated by 1.7 pm. Determine the strength of the magnetic field in the sunspot

[Question ID = 6662]

1. 3 T [Option ID = 26642]
2. 0.03 T [Option ID = 26643]
3. 3.3 T [Option ID = 26644]
4. 0.3 T [Option ID = 26645]

Correct Answer :-

- 0.3 T [Option ID = 26645]

2) which one of the following is correct in respect of an electron and a proton having a same de-Broglie wavelength of 2 \AA

[Question ID = 6663]

1. Both have same kinetic energy [Option ID = 26646]
2. Both have same velocity [Option ID = 26647]
3. Both have same momentum [Option ID = 26648]
4. The kinetic energy of proton is more than that of electron [Option ID = 26649]

Correct Answer :-

- Both have same momentum [Option ID = 26648]

3) If r_p & r_H are the radius and E_p & E_H are the energy of an electron in the n^{th} orbit of positronium atom and hydrogen atom respectively, then

[Question ID = 6664]

1. $r_p = 2 r_H$ and $E_p = E_H/2$
[Option ID = 26650]
2. $r_p = 2 r_H$ and $E_p = 2 E_H$
[Option ID = 26651]
3. $r_p = 2 r_H$ and $E_p = E_H/4$
[Option ID = 26652]
4. $r_p = r_H$ and $E_p = 2 E_H$
[Option ID = 26653]

Correct Answer :-

- $r_p = 2 r_H$ and $E_p = E_H/2$
[Option ID = 26650]

4) An X-ray beam of wavelength 0.16 nm is incident on a set of planes of a certain crystal. The first Bragg reflection is observed for an incidence angle of 30° . What is the corresponding inter planar spacing?

[Question ID = 6669]

1. 0.16 nm [Option ID = 26670]
2. 0.67 nm [Option ID = 26671]
3. 1.02 nm [Option ID = 26672]
4. 0.89 nm [Option ID = 26673]

Correct Answer :-

- 0.16 nm [Option ID = 26670]

5) What is the velocity of conduction electron of silver having Fermi energy 5.52 eV

[Question ID = 6670]

1. 1.39×10^6 m/s [Option ID = 26674]
2. 2.39×10^6 m/s [Option ID = 26675]
3. 0.89×10^6 m/s [Option ID = 26676]
4. 0 [Option ID = 26677]

Correct Answer :-

- 1.39×10^6 m/s [Option ID = 26674]

6) Given that a piece of n-type silicon contains $8 \times 10^{21} \text{ m}^{-3}$ phosphorus impurity atoms, calculate the carrier concentration of silicon at room temperature. Given that the intrinsic electron concentration of silicon at room temperature is $1.6 \times 10^{16} \text{ m}^{-3}$

[Question ID = 6671]

1. $3.2 \times 10^{10} \text{ m}^{-3}$ [Option ID = 26678]
2. $2.3 \times 10^{11} \text{ m}^{-3}$ [Option ID = 26679]
3. $1.5 \times 10^{10} \text{ m}^{-3}$ [Option ID = 26680]
4. $3.2 \times 10^{11} \text{ m}^{-3}$ [Option ID = 26681]

Correct Answer :-

- $3.2 \times 10^{10} \text{ m}^{-3}$ [Option ID = 26678]

7) The dispersion relation for a one-dimensional monoatomic lattice chain is given by the equation, $\omega = \frac{2}{a} v_s \left| \sin\left(\frac{Ka}{2}\right) \right|$, where, 'a' is the interatomic spacing, $K = \frac{2\pi}{\lambda}$ and v_s has the dimension of velocity. The relation between the phase velocity V_p and group velocity V_g in the long wavelength limit is given by

[Question ID = 6674]

1. $V_p = V_g$
[Option ID = 26690]
2. $V_p = 2V_g$
[Option ID = 26691]
3. $V_p = V_g/2$
[Option ID = 26692]
4. $V_p \neq V_g$
[Option ID = 26693]

Correct Answer :-

- $V_p = V_g$
[Option ID = 26690]

8) The largest wavelength of light falling on double slits separated by $1.5 \mu\text{m}$, for which there is a first order maximum is in the,

[Question ID = 6676]

1. ultraviolet range [Option ID = 26698]
2. visible range [Option ID = 26699]
3. infrared range [Option ID = 26700]
4. X-ray range [Option ID = 26701]

Correct Answer :-

- infrared range [Option ID = 26700]

9) In a multi-stage R-C coupled amplifier, the coupling capacitor

[Question ID = 6680]

1. limits the low frequency response [Option ID = 26714]
2. limits the high frequency response [Option ID = 26715]
3. reduces the amplitude of input signal [Option ID = 26716]
4. blocks d.c. component without affecting the frequency response [Option ID = 26717]

Correct Answer :-

- blocks d.c. component without affecting the frequency response [Option ID = 26717]

10) An AM transmitter is coupled to an aerial. The input current is found to be 5 A. With modulation the current value increases to 5.9 A. The depth of modulation is

[Question ID = 6681]

1. 83.4% [Option ID = 26718]
2. 88.6% [Option ID = 26719]
3. 78.2% [Option ID = 26720]
4. 74.3% [Option ID = 26721]

Correct Answer :-

- 88.6% [Option ID = 26719]

11) Hexadecimal equivalent of a digital number 10011101 is

[Question ID = 6683]

1. H913 [Option ID = 26726]
2. 9D [Option ID = 26727]
3. AE [Option ID = 26728]
4. 157 [Option ID = 26729]

Correct Answer :-

- 9D [Option ID = 26727]

12) If the doping concentration in a Si-Zener diode is increased, the Zener breakdown voltage

[Question ID = 6684]

1. Decreases [Option ID = 26730]
2. Increases [Option ID = 26731]
3. Remains unchanged [Option ID = 26732]
4. Becomes broader [Option ID = 26733]

Correct Answer :-

- Decreases [Option ID = 26730]

13) Which one of the following is an example of doubly magic nuclei

[Question ID = 6685]

1. ^{18}O
[Option ID = 26734]
2. ^{48}Ca
[Option ID = 26735]
3. ^{124}Sn
[Option ID = 26736]
4. ^{204}Pb
[Option ID = 26737]

Correct Answer :-

- ^{48}Ca
[Option ID = 26735]

14) Which radiation has maximum ionization power?

[Question ID = 6686]

1. Alpha [Option ID = 26738]
2. Beta [Option ID = 26739]
3. Neutron [Option ID = 26740]
4. Gamma [Option ID = 26741]

Correct Answer :-

- Alpha [Option ID = 26738]

15) For beta-minus decay, which statement is TRUE?

[Question ID = 6688]

1. Daughter nuclide atomic mass (A_D) is more than that of the parent nuclide atomic mass (A_P) [Option ID = 26746]
2. Daughter nuclide atomic number (Z_D) is same that of the parent nuclide atomic number (Z_P) [Option ID = 26747]
3. Daughter nuclide neutron number (N_D) is less than that of the parent nuclide neutron number (N_P) [Option ID = 26748]
4. Daughter nuclide neutron number (N_D) is same that of the parent nuclide neutron number (N_P) [Option ID = 26749]

Correct Answer :-

- Daughter nuclide neutron number (N_D) is less than that of the parent nuclide neutron number (N_P) [Option ID = 26748]

16) The probability that student A solves the problem is $1/2$, and that of B is $2/3$. What is the probability that the problem is solved?

[Question ID = 6689]

1. $4/6$
[Option ID = 26750]
2. $1/3$
[Option ID = 26751]
3. $5/6$
[Option ID = 26752]
4. none of these
[Option ID = 26753]

Correct Answer :-

- $5/6$
[Option ID = 26752]

17) Are the three points whose position vectors are $2i+3j-4k$, $i-2j+3k$ and $-7j+10k$ collinear?

[Question ID = 6690]

1. Yes
[Option ID = 26754]
2. No

[Option ID = 26755]

3. Cannot be determined

[Option ID = 26756]

4. None of these

[Option ID = 26757]

Correct Answer :-

- Yes

[Option ID = 26754]

18) The number of independent fundamental solutions in n-th order ordinary differential equation is

[Question ID = 6692]

1. n-1 [Option ID = 26762]

2. n [Option ID = 26763]

3. n+1 [Option ID = 26764]

4. 2n [Option ID = 26765]

Correct Answer :-

- n [Option ID = 26763]

19) If $z_1 = 2 - 3i$ and $z_2 = 4 + 6i$ then find $\frac{z_1}{z_2}$

[Question ID = 6693]

1. $-5/26-6i/13$

[Option ID = 26766]

2. $-5/26+6i/13$

[Option ID = 26767]

3. $8+18i$

[Option ID = 26768]

4. $8-18i$

[Option ID = 26769]

Correct Answer :-

- $-5/26-6i/13$

[Option ID = 26766]

20) The rank of the following matrix $\begin{pmatrix} 1 & 5 & 1 \\ 2 & 1 & 1 \\ 3 & 6 & 2 \end{pmatrix}$ is

[Question ID = 6699]

1. 1

[Option ID = 26790]

2. 2

[Option ID = 26791]

3. 3

[Option ID = 26792]

4. 4

[Option ID = 26793]

Correct Answer :-

- 2

[Option ID = 26791]

21) Two Carnot engines X and Y are operating in series. The engine X receives heat at 1200 K and rejects to a reservoir at a temperature T. The second engine Y receives the heat rejected by X and in turn rejects to a heat reservoir at 300 K. Calculate the temperature T (in Kelvin) for the situation when the efficiency of the engines is same

[Question ID = 6704]

1. 600 K [Option ID = 26810]

2. 750 K [Option ID = 26811]

3. 0 [Option ID = 26812]

4. 450 K [Option ID = 26813]

Correct Answer :-

- 600 K [Option ID = 26810]

22) A square conducting loop of mass m, side l and resistance R is dropped into a region with a uniform

field B whose direction is perpendicular to the plane of the falling loop. The loop will reach a terminal velocity v given by

[Question ID = 7330]

1. $v = \frac{mgR}{(Bl)^2}$

[Option ID = 29314]

2. $v = \frac{2mgR}{B^2l^2}$

[Option ID = 29315]

3. $v = \frac{mgR}{2B^2l^2}$

[Option ID = 29316]

4. None of these

[Option ID = 29317]

Correct Answer :-

• $v = \frac{mgR}{(Bl)^2}$

[Option ID = 29314]

23) An ideal inductor, a resistor of resistance R Ohms and a capacitor with adjustable capacitance are connected in series to an alternating voltage with an effective value of V Volts and with frequency of f Hz. The current flowing through the circuit when the capacitance of the capacitor is set to C_1 is the same as when the capacitance of the capacitor is set to C_2 , $C_2 > C_1$. The inductance of the inductor L is given by

[Question ID = 7331]

1. $\frac{1}{8\pi^2 f^2} \frac{C_1 + C_2}{C_1 C_2}$

[Option ID = 29318]

2. $\frac{1}{8\pi^2 f^2} \frac{C_1 C_2}{C_1 + C_2}$

[Option ID = 29319]

3. $\frac{1}{2\pi f} \frac{C_1 C_2}{C_1 - C_2}$

[Option ID = 29320]

4. $\frac{1}{2\pi^2 f^2 R(C_1 - C_2)} \frac{C_1 C_2}{C_1 + C_2}$

[Option ID = 29321]

Correct Answer :-

• $\frac{1}{8\pi^2 f^2} \frac{C_1 + C_2}{C_1 C_2}$

[Option ID = 29318]

24) A cylinder of length L is made up of an inner core of steel of radius r_1 and an outer sheath of copper of thickness r_1 . The resistivities of steel and copper are ρ_1 and ρ_2 respectively. The total resistance of the cylinder is

[Question ID = 7332]

1. $\frac{(\rho_1 \rho_2)L}{\pi r_1^2 (3\rho_1 + \rho_2)}$

[Option ID = 29322]

2. $\frac{(3\rho_1 + \rho_2)L}{\pi r_1^2}$

[Option ID = 29323]

3. $\frac{(\rho_1 + \rho_2)L}{4\pi r_1^2}$

[Option ID = 29324]

4. Cannot be determined from the information provided above

[Option ID = 29325]

Correct Answer :-

• $\frac{(\rho_1 \rho_2)L}{\pi r_1^2 (3\rho_1 + \rho_2)}$

[Option ID = 29322]

25) A meter stick is at an angle of 45° to the x' axis in its rest frame. The rod moves with a speed of $\frac{1}{\sqrt{2}}c$ along the $+x$ direction w.r.t. a frame S . The length of the rod in S is

[Question ID = 7333]

1. $\frac{\sqrt{3}}{2}$ meters

[Option ID = 29326]

2. $\frac{\sqrt{5}}{3}$ meters

[Option ID = 29327]

3. $\frac{\sqrt{2}}{3}$ meters

[Option ID = 29328]

4. $\frac{2}{3}$ meters

[Option ID = 29329]

Correct Answer :-

• $\frac{\sqrt{3}}{2}$ meters

[Option ID = 29326]

26) An AC generator with output and frequency f is connected to the plates of an air filled parallel plate capacitor of plate area A and plate separation d . The maximum value of the displacement current is

[Question ID = 7334]

1. $\frac{2\pi \epsilon_0 fVA}{d}$

[Option ID = 29330]

2. $\frac{\epsilon fV}{d}$

[Option ID = 29331]

3. $\frac{2\pi f \epsilon_0 A}{Vd}$

[Option ID = 29332]

4. Cannot be determined from the information provided

[Option ID = 29333]

Correct Answer :-

• $\frac{2\pi \epsilon_0 fVA}{d}$

[Option ID = 29330]

27) An electron enters a uniform magnetic field of flux density 1.2 Wb/m^2 . Find the energy difference in (eV), between electrons having spins parallel and anti-parallel to the field. (Given: $\mu_B = 9.3 \times 10^{-24} \text{ J/T}$)

[Question ID = 7335]

1. $3.95 \times 10^{-5} \text{ eV}$

[Option ID = 29334]

2. $13.95 \times 10^{-5} \text{ eV}$

[Option ID = 29335]

3. $23.95 \times 10^{-5} \text{ eV}$

[Option ID = 29336]

4. $33.95 \times 10^{-5} \text{ eV}$

[Option ID = 29337]

Correct Answer :-

- $13.95 \times 10^{-5} \text{ eV}$

[Option ID = 29335]

28) Using the vector atom model, determine the possible values of the angular momentum of an electron in f - shell

[Question ID = 7336]

1. $\frac{3\sqrt{7}}{2} \hbar, \frac{\sqrt{35}}{2} \hbar$

[Option ID = 29338]

2. $\frac{2\sqrt{7}}{2} \hbar, \frac{\sqrt{25}}{2} \hbar$

[Option ID = 29339]

3. $\frac{5\sqrt{7}}{2} \hbar, \frac{\sqrt{15}}{2} \hbar$

[Option ID = 29340]

4. $\frac{\sqrt{7}}{2} \hbar, \frac{\sqrt{5}}{2} \hbar$

[Option ID = 29341]

Correct Answer :-

- $\frac{3\sqrt{7}}{2} \hbar, \frac{\sqrt{35}}{2} \hbar$

[Option ID = 29338]

29) The two eigenvalues of the matrix $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ are

[Question ID = 7337]

1. 2,0

[Option ID = 29342]

2. 1,1

[Option ID = 29343]

3. 1,2

[Option ID = 29344]

4. 0,1

[Option ID = 29345]

Correct Answer :-

- 2,0

[Option ID = 29342]

30) The commutator, $\pi[x^2, p_x]$, is equal to

[Question ID = 7338]

1. ihx

[Option ID = 29346]

2. $2ihx$

[Option ID = 29347]

3. $2ihp_x$

[Option ID = 29348]

4. Zero

[Option ID = 29349]

Correct Answer :-

- ihx

[Option ID = 29346]

31) A particle of mass m is confined in the ground state of a one dimensional box extending from $x = -2L$ to $x = +2L$. The wave function of the particle in this state is

From the wave function of the particle in this state is
 $\Psi(x) = \psi_0 \cos\left(\frac{\pi x}{4L}\right)$, Where, ψ_0 is a constant. The energy eigenvalue corresponding to this state is

[Question ID = 7339]

1. $\frac{\hbar^2 \pi^2}{32mL^2}$

[Option ID = 29350]

2. $\frac{\hbar^2 \pi^2}{2 mL^2}$

[Option ID = 29351]

3. $\frac{\hbar^2 \pi^2}{4 mL^2}$

[Option ID = 29352]

4. $\frac{\hbar^2 \pi^2}{16mL^2}$

[Option ID = 29353]

Correct Answer :-

• $\frac{\hbar^2 \pi^2}{32mL^2}$

[Option ID = 29350]

32) The normalized wave functions ψ_1 and ψ_2 , correspond to the ground state and the first excited states of a particle in a potential. The operator \hat{A} acts on the wave functions as $\hat{A}\psi_1 = \psi_2$ and $\hat{A}\psi_2 = \psi_1$. The expectation value of the operator \hat{A} for the state $\psi = (3\psi_1 + 4\psi_2)/5$ is

[Question ID = 7340]

1. 0.96 [Option ID = 29354]

2. - 0.32 [Option ID = 29355]

3. 0 [Option ID = 29356]

4. 0.75 [Option ID = 29357]

Correct Answer :-

• 0.96 [Option ID = 29354]

33) The primitive translation vector of a two-dimensional lattice are $\mathbf{a} = 2\hat{i} + \hat{j}$, $\mathbf{b} = 2\hat{j}$. The primitive translation vector of its reciprocal lattice in x-direction is given by

[Question ID = 7341]

1. $\mathbf{a}^* = \pi\hat{i}$

[Option ID = 29358]

2. $\mathbf{a}^* = 2\pi\hat{i}$

[Option ID = 29359]

3. $\mathbf{a}^* = \hat{i}$

[Option ID = 29360]

4. $\mathbf{a}^* = \pi\hat{j}$

[Option ID = 29361]

Correct Answer :-

• $\mathbf{a}^* = \pi\hat{i}$

[Option ID = 29358]

34) The mean drift speed v_d of an electron in an applied electric field E with electron density 'n' can be expressed as

[Question ID = 7342]

1. $v_d = |\sigma E/ne|$

[Option ID = 29362]

2. $v_d = |\sigma E/e|$

[Option ID = 29363]

3. $v_d = |\sigma e/nE|$

[Option ID = 29364]

4. none of these

[Option ID = 29365]

Correct Answer :-

• $v_d = |\sigma E/ne|$

[Option ID = 29362]

35) An un-damped oscillator has time period $\tau_0 = 1.0$ sec. Now a little damping is added so that its time period changes to $\tau_1 = 1.001$ sec. By what factor will the amplitude of oscillation decrease after 10 cycles?

[Question ID = 7343]

1. ≈ 17

[Option ID = 29366]

2. ≈ 1

[Option ID = 29367]

3. $\approx 1/17$

[Option ID = 29368]

4. None of these

[Option ID = 29369]

Correct Answer :-

• ≈ 17

[Option ID = 29366]

36) A kilogram of water has a constant heat capacity of 4.2 kJ/K/kg over the temperature range 0°C to 100°C . The water was initially at 0°C and is brought into contact with a heat reservoir at 100°C . When the water is in thermal equilibrium with the heat reservoir calculate the change in entropy of the universe (Water + Reservoir).

[Question ID = 7344]

1. 184.8 J/K

[Option ID = 29370]

2. 2437.8 J/K

[Option ID = 29371]

3. 0 J/K

[Option ID = 29372]

4. 1310.8 J/K

[Option ID = 29373]

Correct Answer :-

• 184.8 J/K

[Option ID = 29370]

37) Two identical finite bodies of constant volume and of constant heat capacity at constant volume C_v , are used to drive a heat engine. Their initial temperatures are T_1 and T_2 . The maximum amount of work which can be obtained from the system is

[Question ID = 7345]

1. $C_v \{ 2 (T_1 T_2)^{1/2} - (T_1 + T_2) \}$

[Option ID = 29374]

2. $C_v (T_1 + T_2)$

[Option ID = 29375]

3. $C_v (T_1 T_2)^{1/2}$

[Option ID = 29376]

4. 0

[Option ID = 29377]

Correct Answer :-

- $C_v \{ 2 (T_1 T_2)^{1/2} - (T_1 + T_2) \}$

[Option ID = 29374]

38) For a system of bosons, we can write the Bose-Einstein distribution function as $f(E_i) = \frac{1}{\exp(\alpha + \beta E_i) - 1}$, Where, $\beta = 1/k_B T$ and $\alpha = \mu/k_B T$ (k_B = Boltzmann constant). If μ represents the chemical potential, then which one of the following is true?

[Question ID = 7346]

1. $\mu \leq 0$

[Option ID = 29378]

2. $\mu \geq 0$

[Option ID = 29379]

3. $\mu \leq 1$

[Option ID = 29380]

4. $\mu \geq 1$

[Option ID = 29381]

Correct Answer :-

- $\mu \geq 0$

[Option ID = 29379]

39) An ideal capacitor C is charged to a voltage V_0 and connected at $t = 0$ across an ideal inductor L (The circuit now consists of a capacitor and an inductor only). If the resonant frequency

$$\omega_0 = \frac{1}{\sqrt{LC}}$$

, the voltage across the capacitor at time $t > 0$ is given by

[Question ID = 7347]

1. V_0 [Option ID = 29382]

2. $V_0 \cos(\omega_0 t)$ [Option ID = 29383]

3. $V_0 \sin(\omega_0 t)$ [Option ID = 29384]

4. $V_0 e^{-\omega_0 t} \cos(\omega_0 t)$

[Option ID = 29385]

Correct Answer :-

- $V_0 \cos(\omega_0 t)$ [Option ID = 29383]

40) Magnetic moment of proton (μ_p) in terms of nuclear magneton (μ_N) is

[Question ID = 7348]

1. $\mu_p = 1.9\mu_N$

[Option ID = 29386]

2. $\mu_p = 2.7\mu_N$

[Option ID = 29387]

3. $\mu_p = 3.8\mu_N$

[Option ID = 29388]

4. $\mu_p = 5.4\mu_N$

[Option ID = 29389]

Correct Answer :-

- $\mu_p = 2.7\mu_N$

[Option ID = 29387]

41) Find the eigenvalues of $A+4I$ where I is identity matrix and $A = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$

[Question ID = 7349]

1. 1,3

[Option ID = 29390]

2. 5,7

[Option ID = 29391]

3. 4,4

[Option ID = 29392]

4. None of these

[Option ID = 29393]

Correct Answer :-

- 5,7

[Option ID = 29391]

42) The limit $\lim_{n \rightarrow \infty} \left(\frac{1}{n}\right)^{\frac{1}{n}}$ is

[Question ID = 7350]

1. $\frac{1}{e}$

[Option ID = 29394]

2. 1

[Option ID = 29395]

3. 0

[Option ID = 29396]

4. e

[Option ID = 29397]

Correct Answer :-

- 1

[Option ID = 29395]

43) $\left(\frac{1+i}{\sqrt{2}}\right)^{49}$

is equal to

[Question ID = 7351]

1. $\left(\frac{1+i}{\sqrt{2}}\right)$

[Option ID = 29398]

2. $\left(\frac{2+98i}{\sqrt{2^{49}}}\right)$

[Option ID = 29399]

3. $\left(\frac{1-i}{\sqrt{2}}\right)$

[Option ID = 29400]

4. $\left(\frac{2-98i}{\sqrt{2^{49}}}\right)$

[Option ID = 29401]

Correct Answer :-

- $\left(\frac{1+i}{\sqrt{2}}\right)$

[Option ID = 29398]

44) $\sin\left(\frac{\pi}{4} + i\right)$ is equal to

[Question ID = 7352]

1. $\frac{\sqrt{2}}{4} \left(e - \frac{1}{e}\right) + \frac{\sqrt{2}}{4} \left(e - \frac{1}{e}\right) i$

[Option ID = 29402]

2. $\frac{\sqrt{2}}{4} \left(e + \frac{1}{e}\right) + \frac{\sqrt{2}}{4} \left(e - \frac{1}{e}\right) i$

[Option ID = 29403]

3. $\frac{\sqrt{2}}{4} \left(e + \frac{1}{e} \right) + \frac{\sqrt{2}}{4} \left(e + \frac{1}{e} \right) i$

[Option ID = 29404]

4. $\frac{\sqrt{2}}{4} \left(e - \frac{1}{e} \right) + \frac{\sqrt{2}}{4} \left(e + \frac{1}{e} \right) i$

[Option ID = 29405]

Correct Answer :-

• $\frac{\sqrt{2}}{4} \left(e + \frac{1}{e} \right) + \frac{\sqrt{2}}{4} \left(e - \frac{1}{e} \right) i$

[Option ID = 29403]

45) Two students are working on a math problem. The first student has probability $\frac{1}{2}$ of solving it and the second student has probability $\frac{3}{4}$ of solving it. What is the probability that at least one of them solves the problem?

[Question ID = 7353]

1. $\frac{3}{8}$

[Option ID = 29406]

2. $\frac{5}{8}$

[Option ID = 29407]

3. $\frac{7}{8}$

[Option ID = 29408]

4. $\frac{9}{8}$

[Option ID = 29409]

Correct Answer :-

• $\frac{7}{8}$

[Option ID = 29408]

46) Expansion of the function $f(z) = \frac{1}{z^2-3z+2}$ in the region defined by $|z| > 2$ is

[Question ID = 7354]

1. $z^{-2} + 3z^{-4} + 7z^{-6} + \dots$

[Option ID = 29410]

2. $z^{-2} + 3z^{-3} + 7z^{-4} + \dots$

[Option ID = 29411]

3. $z^{-1} + 3z^{-2} + 7z^{-3} + \dots$

[Option ID = 29412]

4. $z^{-3} + 3z^{-4} + 7z^{-5} + \dots$

[Option ID = 29413]

Correct Answer :-

• $z^{-2} + 3z^{-3} + 7z^{-4} + \dots$

[Option ID = 29411]

47) The Fourier transformation of the function $f(x) = 1$ for $|x| < a$

= 0 for $|x| > a$

is

[Question ID = 7355]

1. $\sqrt{\frac{2}{\pi}} \frac{\sin sa}{s}$

[Option ID = 29414]

2. $\sqrt{\frac{2 \cos sa}{\pi s}}$

[Option ID = 29415]

3. $\sqrt{\frac{\pi \sin sa}{2 s}}$

[Option ID = 29416]

4. $\sqrt{\frac{\pi \cos sa}{2 s}}$

[Option ID = 29417]

Correct Answer :-

• $\sqrt{\frac{2 \sin sa}{\pi s}}$

[Option ID = 29414]

48) The Laplace transformation of the function $f(t) = 2^t$ is

[Question ID = 7356]

1. $\frac{\ln 2}{s - \ln 2}$

[Option ID = 29418]

2. $\frac{1}{s - \ln 2}$

[Option ID = 29419]

3. $\frac{1}{s \ln 2 - 1}$

[Option ID = 29420]

4. $\frac{\ln 2}{s \ln 2 - 1}$

[Option ID = 29421]

Correct Answer :-

• $\frac{1}{s - \ln 2}$

[Option ID = 29419]

49) Consider a collection of non-interacting particles, each of mass m in a volume where the gravitational force is a -ve (z-direction). Consider the system is in thermal equilibrium at a temperature T . Find the partition function

[Question ID = 7357]

1. $Q_N = \left[\frac{(kT)^3}{2\pi m g \hbar^2} \right]^{N/2}$

[Option ID = 29422]

2. $Q_N = \left[\frac{2\pi m g \hbar^2}{(kT)^3} \right]^{N/2}$

[Option ID = 29423]

3. $Q_N = \left[\frac{(kT)^3}{2\pi m g \hbar^2} \right]^N$

[Option ID = 29424]

4. $Q_N = \left[\frac{2\pi m g \hbar^2}{(kT)^3} \right]^N$

[Option ID = 29425]

Correct Answer :-

$$\bullet Q_N = \left[\frac{(kT)^3}{2\pi m g h^2} \right]^{N/2}$$

[Option ID = 29422]

50) The quantum distribution function for any gas atom which follows MB, BE and FD statistics is given as a generalized single form

$$f_i = g_i / [\exp(\epsilon_i - \mu) / (kT + J)]$$

If the distribution function follows the MB statistics in a classical limit then what will be the condition of the following. Symbols have their usual meanings

[Question ID = 7358]

1. $\frac{f_i}{g_i} \ll 1 ; J = 1$

[Option ID = 29426]

2. $\frac{f_i}{g_i} \ll 1 ; J = 0$

[Option ID = 29427]

3. $\frac{g_i}{f_i} \gg 1 ; J = 1$

[Option ID = 29428]

4. $\frac{g_i}{f_i} \gg 1 ; J = -1$

[Option ID = 29429]

Correct Answer :-

$$\bullet \frac{f_i}{g_i} \ll 1 ; J = 0$$

[Option ID = 29427]