




Marking scheme – 2017 (Compartment)



CHEMISTRY (043)/ CLASS XII

Set 56/1/1




Q.No	Value Points	Marks
1	hcp	1
2	$AlCl_3 / Al^{3+}$	1
3	Orbital splitting energies are not sufficiently large for forcing pairing	1
4	2,3-dinitro phenol	1
5	Having no α - hydrogen	1
6	Vapour pressure of the solvent decreases in the presence of non – volatile solute (glucose) hence boiling point increases	2
7	(i) First order (ii) $s^{-1} / \text{time}^{-1}$	1 1
8	Hypophosphorous acid is a good reducing agent as it contains two P-H bonds. There is no P-H bond in orthophosphoric acid, so it is not a reducing agent Example : It reduces $AgNO_3$ to metallic silver/ chemical equation	1 1
	OR	
8	a) 4 b) Due to lower bond dissociation enthalpy of BiH_3 as compared to SbH_3	1 1
9	i. Due to resonance the two O-O bond lengths are identical. ii. Due to strong bond formed by it with other elements.	1 1
10.	i) (b) is chiral ii) (a) will undergo S_N2 reaction faster	1 1
11	In bcc, $z=2$; $d = (zxM) / a^3 \times N_A$ (i) No. of atoms = $\frac{w}{M} \times N_A$ $2.5 \times 10^{24} = \frac{500g}{M} \times N_A$ $M = [500 \times N_A] / 2.5 \times 10^{24}$ (ii) Putting values of M in equation (i) $d = 2 \times 500 g \times N_A / [2.5 \times 10^{24} \text{ atoms} \times (400 \times 10^{-10} \text{ cm})^3 \times N_A]$ $d = 6.25 \text{ g/ cm}^3$ (or any other correct method)	$\frac{1}{2}$ 1 $\frac{1}{2}$ 1
12	$p_{\text{total}} = p_1^{\circ} + (p_2^{\circ} - p_1^{\circ}) x_2$ $600 = 450 + (700 - 450) x_2$ $x_2 = 0.6$ $x_2 = 1 - 0.6 = 0.4$	1 1 $\frac{1}{2}$ $\frac{1}{2}$
13	$P_A = 2P_o - P_t$ $= (2 \times 0.4) - 0.7 = 0.1$ $k = \frac{2.303}{t} \log P_o/P_A$ $k = \frac{2.303}{100} \log 0.4/0.1$ $k = \frac{2.303}{100} \times 0.6021$ $= 1.39 \times 10^{-2} s^{-1}$	$\frac{1}{2}$ $\frac{1}{2}$ 1 1
14	i) The process of removing an adsorbed substance from a surface on which it is	1



	<p>adsorbed.</p> <p>ii) The formation of micelles takes place only above a particular concentration called CMC.</p> <p>iii) The catalytic reaction that depends upon the pore structure of the catalyst and size of the reactant and product molecules.</p>	<p>1</p> <p>1</p>
15	<p>a) The metal is converted into its volatile compound and collected elsewhere. It is then decomposed to get the pure metal.</p> <p>b) i) Ni ii) Ti/Zr</p> <p>c) It is used to separate two sulphide ores by preventing one to form froth.</p>	<p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>1</p>
16	<p>a) $H_2O < H_2S < H_2Se < H_2Te$, because of decrease in bond dissociation enthalpy.</p> <p>b) </p>	<p>1,1</p> <p>1</p>
OR		
	<p>a) i) Due to higher oxidation state of P in PCl_5</p> <p>ii) Liberation of hydrogen prevents the formation of $FeCl_3$</p> <p>b) </p>	<p>1</p> <p>1</p> <p>1</p>
17	<p>Hybridisation : sp^3d^2</p> <p>Magnetic character : Paramagnetic</p> <p>Spin nature: High spin</p>	<p>1</p> <p>1</p> <p>1</p>
18.	<p>a) A: $CH_3-CH=CH_2$ B: $CH_3-CH_2-CH_2Br$ C: $CH_3-CH_2-CH_2I$ D: $CH_3-CH_2-CH_2MgI$</p> <p></p> <p>b)</p>	<p>$\frac{1}{2} \times 4$</p> <p>1</p>
19.	<p>a) $CH_3-O-CH_3 + HI \longrightarrow CH_3-OH + CH_3-I$</p> <p>b) .</p>	<p>1</p>

	<p>Protonation of alkene to form carbocation by electrophilic attack of H_3O^+.</p> $\text{H}_2\text{O} + \text{H}^+ \rightarrow \text{H}_3\text{O}^+$ $\text{>C=C<} + \text{H}-\overset{\text{H}}{\underset{\cdot\cdot}{\text{O}}^+}-\text{H} \rightleftharpoons \text{-}\overset{\text{H}}{\text{C}}-\overset{+}{\text{C}}\text{-} + \text{H}_2\ddot{\text{O}}$ <p>Nucleophilic attack of water on carbocation.</p> $\text{-}\overset{\text{H}}{\text{C}}-\overset{+}{\text{C}}\text{-} + \text{H}_2\ddot{\text{O}} \rightleftharpoons \text{-}\overset{\text{H}}{\text{C}}-\overset{\text{H}}{\text{C}}-\overset{+}{\text{O}}-\text{H}$ <p>Deprotonation to form an alcohol.</p> $\text{-}\overset{\text{H}}{\text{C}}-\overset{\text{H}}{\text{C}}-\overset{+}{\text{O}}-\text{H} + \text{H}_2\ddot{\text{O}} \rightarrow \text{-}\overset{\text{H}}{\text{C}}-\overset{\cdot\cdot}{\text{O}}\text{H}- + \text{H}_3\ddot{\text{O}}^+$	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p>
20.	<p>i) A: $\text{CH}_3\text{-CH}_2\text{CN}$; B: $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{NH}_2$; C: $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-NH-COCH}_3$</p> <p>ii) A: $\text{Ar}-\overset{+}{\text{N}}_2\text{BF}_4^-$; B:  ; C: </p>	<p>$\frac{1}{2} \times 3$</p> <p>$\frac{1}{2} \times 3$</p>
21	<p>a) Glycosidic linkage</p> <p>b) Source : Meat, Fish, egg, curd (any one) ; Pernicious anaemia</p> <p>c) DNA is double strand while RNA is single strand molecule (or any other correct difference)</p>	<p>1</p> <p>$\frac{1}{2}, \frac{1}{2}$</p> <p>1</p>
22	<p>i) Treatment of hyperacidity Class : Antacids</p> <p>ii) Relieve pain and produce sleep Class: Narcotic analgesics</p> <p>iii) Relieve pain and reduce fever Class: Non- Narcotic analgesics / Analgesics</p>	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
23	<p>a) <i>Poly β-hydroxybutyrate – co-β-hydroxy valerate / (PHBV)</i></p> <p>Monomers : $\text{CH}_3-\overset{\text{OH}}{\text{CH}}-\text{CH}_2-\text{COOH}$, $\text{CH}_3-\text{CH}_2-\overset{\text{OH}}{\text{CH}}-\text{CH}_2-\text{COOH}$</p> <p>Repeating unit :</p> $\left(\text{O}-\underset{\text{CH}_3}{\text{CH}}-\text{CH}_2-\overset{\text{O}}{\underset{\text{O}}{\text{C}}}-\text{O}-\underset{\text{CH}_2\text{CH}_3}{\text{CH}}-\text{CH}_2-\overset{\text{O}}{\underset{\text{O}}{\text{C}}} \right)_n$ <p>b) PHBV is used in speciality packaging, orthopaedic devices and in controlled release of drugs.(any two)</p> <p>c) Concern for environment , caring (or any other)</p>	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}, \frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}, \frac{1}{2}$</p> <p>$\frac{1}{2}, \frac{1}{2}$</p>
24	<p>a) E^0 value of silver is lower than that of gold, hence silver displaces gold which gets deposited on the silver object. E^0 value of copper is lower than that of silver, hence silver cannot displace copper from its solution.</p> <p>b) i) Electrons flow from Zn to Ag plate. ii) Zn as anode and Ag acts as cathode iii) Cell will stop functioning iv) Concentration of Zn^{2+} ions will increase and that of Ag^+ ions will decrease. v) No change</p>	<p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}, \frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
	OR	



24	<p>a) When concentration approaches zero, the molar conductivity is known as limiting molar conductivity</p> <p>The change in Δm with dilution is due to the increase in the degree of dissociation and consequently the number of ions in the total volume of the solution that contains 1 mol of electrolyte, hence Δm increases steeply.</p> <p>b) $E_{\text{cell}} = E^{\circ}_{\text{cell}} - \frac{0.059}{n} \log \frac{[Mg^{2+}]}{[Cu^{2+}]}$</p> $= 2.71 \text{ V} - \frac{0.059}{2} \log \frac{0.1}{0.001}$ $= 2.71 \text{ V} - \frac{0.059}{2} \log 10^2$ $= 2.651 \text{ V}$	1 1 1 1 1
25	<p>a) A: Na_2CrO_4 ; B: $\text{Na}_2\text{Cr}_2\text{O}_7$; C : $\text{K}_2\text{Cr}_2\text{O}_7$</p> $4 \text{ FeCr}_2\text{O}_4 + 8 \text{ Na}_2\text{CO}_3 + 7 \text{ O}_2 \rightarrow 8 \text{ Na}_2\text{CrO}_4 + 2 \text{ Fe}_2\text{O}_3 + 8 \text{ CO}_2$ $2\text{Na}_2\text{CrO}_4 + 2 \text{ H}^+ \rightarrow \text{Na}_2\text{Cr}_2\text{O}_7 + 2 \text{ Na}^+ + \text{H}_2\text{O}$ $\text{Na}_2\text{Cr}_2\text{O}_7 + 2 \text{ KCl} \rightarrow \text{K}_2\text{Cr}_2\text{O}_7 + 2 \text{ NaCl}$	$\frac{1}{2}$, $\frac{1}{2}$, 1 1 1 1
OR		
25	<p>a) i) Copper; Due to high $\Delta_a H^\ominus$ and low $\Delta_{\text{hyd}} H^\ominus$</p> <p>ii) Cerium ; Due to stable $4f^0$ configuration / Tb ; Due to stable $4f^7$ configuration</p> <p>b) i) Due to ability of oxygen to form multiple bonds to metal</p> <p>ii) HCl is oxidized to chlorine</p> <p>iii) Due to strong interatomic metallic bonding.</p>	$\frac{1}{2}$, $\frac{1}{2}$ $\frac{1}{2}$, $\frac{1}{2}$ 1 1 1
26	<p>a) i).</p>  <p>ii).</p>  <p>b) i) Because it is a deactivating group / Due to electron withdrawing carboxylic group resulting in decreased electron density at o- and p- position.</p> <p>ii) Due to extensive association of carboxylic acid molecules through intermolecular hydrogen bonding.</p> <p>iii) Due to steric and +I effect of two methyl groups in propanone</p>	1 1 1 1 1
OR		
26	<p>a) i).</p>  <p>ii) CH_3COOH</p> <p>iii) $\text{CH}_3\text{-CH(Br)-COOH}$</p> <p>b) i) Add ammoniacal solution of silver nitrate / Tollen's reagent to both the compounds, propanal will give silver mirror while propanone does not.</p> <p>ii) Add NaHCO_3 solution to both the compounds, benzoic acid will give effervescence and liberate CO_2 while benzaldehyde will not. (Or any other suitable test)</p>	1 1 1 1 1

1	Dr. (Mrs.) Sangeeta Bhatia		6	Sh. Rakesh Dhawan	
2	Dr. K.N. Uppadhya		7	Dr. (Mrs.) Sunita Ramrakhiani	
3	Prof. R.D. Shukla		8	Mrs. Preeti Kiran	
4	Sh. S.K. Munjal		9	Dr. Azhar Aslam Khan	
5	Sh. D.A. Mishra		10	Ms. Garima Bhutani	



collegedunia.com
India's largest Student Review Platform

**These answers are meant to be used by evaluators*



collegedunia.com
India's largest Student Review Platform