POST GRADUATE COMMON ENTRANCE TEST - 2011

COURSE		SUBJECT
MBA (In	frastructure Management	
MAXIMUM MARKS		MAXIMUM TIME FOR ANSWERING
		120 Minutes
ET NO.	QUESTION I	BOOKLET DETAILS
	VERSION CODE	SERIAL NUMBER
	A ₁	00000105
	MBA (In	ME / M. Tech / M. Arch / MBA (Infrastructure Management courses offered by VTU / UVCE / UBDT) TOTAL DURATION 150 Minutes ET NO. QUESTION E VERSION CODE

DOs

- 1. Check whether the PGCET No. has been entered and shaded in the respective circles on the OMR answer sheet.
- 2. This question booklet is issued to you by the invigilator after the 2nd Bell, i.e. after 10:25 am.
- 3. The serial number of this guestion booklet should be entered on the OMR answer sheet.
- The version code of this question booklet should be entered on the OMR answer sheet and the respective circles should also be shaded completely.
- Compulsorily sign at the bottom portion of the OMR answer sheet in the space provided.

DON'Ts

- 1. The timing and marks printed on the OMR answer sheet should not be damaged / mutilated / spoiled.
- 2. The 3rd Bell rings at 10:30 am, till then;
 - Do not remove the seals of this question booklet.
 - Do not look inside this question booklet.
 - Do not start marking on the OMR answer sheet.

IMPORTANT INSTRUCTIONS TO CANDIDATES

- This question booklet contains 75 (items) questions and each question will have one statement and four answers. (Four different options / responses.)
- After the 3rd bell is rung at 10:30 am, remove the seals of this question booklet and check that this booklet does not
 have any unprinted or torn or missing pages or items etc., if so, get it replaced by a complete test booklet. Read
 each item and start marking on the OMR answer sheet.
- During the subsequent 120 minutes :
 - · Read each question (item) carefully.
 - Choose one correct answer from out of the four available responses (options / choices) given under each
 question / item. In case you feel that there is more than one correct response, mark the response which you
 consider the best. In any case, choose only one response for each question / item.
 - Completely darken / shade the relevant circle with a blue or black ink ballpoint pen against the question number on the OMR answer sheet.
- Please note that even a minute unintended ink dot on the OMR answer sheet will also be recognized and recorded by the scanner. Therefore, avoid multiple markings of any kind on the OMR answer sheet.
- Use the space provided at the bottom on each page of the question booklet for Rough Work. Do not use the OMR answer sheet for the same.
- 6. After the last bell is rung at 12:30 pm, stop marking on the OMR answer sheet and affix your left hand thumb impression on the OMR answer sheet as per the instructions.
- 7. Hand over the OMR answer sheet to the room invigilator as it is.
- 8. After separating the top sheet (KEA copy), the invigilator will return the bottom sheet replica (candidate's copy) to you to carry home for self evaluation.
- 9. Preserve the replica of the OMR answer sheet for a minimum period of ONE year.
- 10. Only Non-programmable calculators are allowed.

Marks Distribution

PART II: 50 Questions carry one mark each (1 to 50)
PART II: 25 Questions carry two marks each (51 to 75)

SEAL



PART - I

Each question carries one mark.

 $50 \times 1 = 50$

- 1. Poise is the unit of
 - (A) velocity gradient

(B) dynamic viscositý

(C) kinematic viscosity

- (D) mass density.
- 2. Venturimeter is used to measure
 - (A) discharge

(B) average velocity

(C) velocity at a point

- (D) pressure at a point.
- If fluid particles move in straight lines and all the lines are parallel to the surface, the flow is called
 - (A) uniform

(B) compressible

(C) laminar

- (D) steady.
- 4. A Newtonian fluid is defined as the fluid which
 - (A) is incompressible and non-viscous
 - (B) obeys Newton's law of viscosity
 - (C) is highly viscous
 - (D) is compressible and non-viscous.
- 5. Bernoulli's equation is derived by making assumption that
 - (A) the flow is uniform and incompressible
 - (B) the flow is non-viscous, uniform and steady
 - (C) the flow is steady, non-viscous, incompressible and irrotational
 - (D) the flow is non-uniform, steady and compressible.

- 6. Reynolds number is defined as the
 - (A) ratio of inertial force to gravity force
 - (B) ratio of viscous force to gravity force
 - (C) ratio of viscous force to elastic force
 - (D) ratio of inertial force to viscous force.
- 7. Which one of the following is not true?
 - (A) Internal energy is a thermodynamic property of a system
 - (B) Internal energy is a state function
 - (C) Internal energy is independent of temperature and pressure of the system
 - (D) Internal energy of a system does not change in a cyclic process.
- 8. The average molecular weight of air is
 - (A) 0.0288 kg/mol

(B) 0.0320 kg/mol

(C) 0.0160 kg/mol

- (D) 0.0148 kg/mol.
- 9. The heat supplied to a system under constant pressure is equal to
 - (A) the work done by the system
- (B) the change in internal energy
- (C) the change in enthalpy
- (D) the change in kinetic energy.
- 10. With increase in pressure, the heat of vaporization of liquids
 - (A) remains unchanged

(B) increases

(C) decreases

O.T.9

- (D) may increase or decrease.
- 11. There exists a unique solution for a set of material balance equations if
 - (A) the number of degrees of freedom is positive
 - (B) the number of degrees of freedom is negative
 - (C) the number of degrees of freedom is non-zero
 - (D) the number of degrees of freedom is zero.

A	1	5		406
12.	The approximate density of water	at room tem	perature is at zont hand to a	19. The un
	(A) 1 kg/m ³	(B)	10 kg/m ³	
	(C) 1000 kg/m ³	(D)	0.01 kg/m ³ .	W (O)
13.	supplied for			
	(A) Isobaric process	(B)	Isothermal process	
	(C) Adiabatic process	(D)	Isochoric process.	
14.	Fugacity coefficient of a substance	is the ratio		21. Therms
	(A) pressure	(B)	activity	
	(C) mole fraction	(D)	activity coefficient.	
15.	An isolated system can exchange .	ottage a lo	r et eram molecular weights	
10.				
	(A) matter	(B)	energy	
	(C) both matter and energy	(D)	neither matter nor energy.	an (O)
16.	Change of heat content when one pressure is called		mpound is burnt in oxygen	
	(A) heat of reaction	(B)	heat of formation	
	(C) heat of combustion	(D)	calorific value.	
17.	PV^{γ} = constant is valid for	proces	ettes of polyesterinearion r.s	24 The ldr
	(A) Isothermal	(B)	Adiabatic	
	(C) Isobaric	(D)	Isotropic.	m (A)
18.	Which of the following is an extens	ive property	of a system ?	mi (H)
	(A) Heat capacity	(B)	Pressure and moderning	
	(C) Concentration	(D)	Volume.	



4.				
19.	The	unit of heat flux is		The approximate density of water a
	(A)	J/s m ²	(B)	W/m ²
	(C)	W/mm ²	(D)	both (A) and (B).
20.	The	thermal conductivity of polyethylene	is	that of aluminium.
	(A)	equal to	(B)	more than
	(C)	less than	(D)	data insufficient.
21.	The	ermal resistance is given by		
	(A)	K/BA	(B)	B/KA
	(C)	KA/B	(D)	B/K.
22.	Nur	nber of gram molecular weights of a	solute	dissolved in 1 kg of solvent is referre
	to a	S		
, -	(A)	molality	(B)	molarity
	(C)	normality	(D)	formality.
23.	Fra	ctional distillation is an example of m	ass tra	ansfer in phase.
	(A)	liquid-liquid	(B)	gas-gas
	(C)	liquid-gas	(D)	solid-liquid.
24.	The	kinetics of polyesterification reaction	n indica	ates that the functional group reactivity
	is	attedation (8		ed colours service immeration. Vi
	(A)	independent on the size of the mone	omer m	nolecule
	(B)	independent concentration of mono	mer mo	olecule and all and the storage of
	(C)	dependent on the size of monomer i	molecu	le witangan Jashi (A)
	(D)	independent on temperature.		



A 1	A 1			7		406
200	25	The	rate of polymerization is proport	ional to		, Q
		(A)	initiator concentration and first	power of	monomer concentration	
		(B)	square root of initiator concent	ration and	first power of monomer concentration	n
13.		(C)	second power of monomer cond	centration	Pittas conduct produced by using ex-	
		(D)	first power of initiator concentr	ation.	(A). Calmarension moldring Victoria	
	26.	In c	ase of second order reaction the	rate of th	e reaction depends on the	
		(A)	concentration of one of the read	ctants		
		(B)	concentration of a reactant rais	sed to the	second power	
		(C)	concentration of a reactant rais	sed to the	first power	
ed		(D)	concentration of reactant raise	d to the n	th power.	
	27.	Unit	for first order reaction rate is			
		(A)	sec - 1	(B)	°C/sec	
		(C)	g/mol	(D)	mole/sec.	
	28.	The	rate of reaction	with	increase in concentration of reac	ting
		spec	cies.	100		
		(A)	increases	(B)	decreases	
		(C)	is unaltered	(D)	increases followed by decreases.	
	29.	The	rate of chemical reaction is dir	rectly prop	portional to the product of the effect	tive
		cond	centration of each participating n	nolecule. T	This theory is known as	
		(A)	Collision theory	(B)	Transition state theory	
		(C)	Law of mass action	(D)	Adsorption theory.	

[P.T.O.

406			8	A
30.	Wat	ter tanks can be manufactured by		The rate of polymerication is propo
	(A)	of monomer concentration MTS	(B)	Compression molding
	(C)	nd first power of monomer MIRRs	(D)	Injection molding.
31.	Filn	ns can be produced by using		
	(A)	Compression molding	(B)	Thermofoaming
	(C)	the reaction depends on the MIS	(D)	Blown film extrusion.
32.	Whi	ich process is referred as primary	process '	. (v) remuestration of one of the r.
	(A)	Calendering 194000 binone so	(B)	Extrusion
	(C)	Injection molding	(D)	Coating.
33.	Clar	mping pressure in injection molding	g is	(D) - concerns allon of respons trus
	(A)	higher than injection pressure		
	(B)	equal to injection pressure	H)	
	(C)	lower than injection pressure	(I)	
getti	(D)	does not have any significance.		
34.	Twi	n screw extruders are extensively	used	Maria de la compansión
	(A)	for mixing and compounding	(B) '	to produce thermofoamed products
	(C)	to get anisotropic behaviour	(D)	to produce pipes.
35.	Hyd	rogen bonding exists in and and		
	(A)	PVC Transition state theory OV9	(B)	(A) Collision (heavy 99
	(C)	Adsorption theory. 34QH	(D)	Nylon.

FP.T.O.



A	ı		9 .	406
36.	Tefl	lon is a trade name for		
	(A)	PVDF	(B)	PCTFE
	(C)	PTT	(D)	PTFE.
37.	Wh	ich of the following is not a Vinyl pol	ymer?	
	(A)	PVC	(B)	PMMA
	(C)	PVA	(D)	PET.
38.	Crit	ical micelle concentration is related i	to	tie servicing returnations (8)
	(A)	plasma polymerization	(B)	gas phase polymerization
	(C)	solution polymerization	(D)	emulsion polymerization.
39.	Poly	mers generally cannot be purified by	y	
	(A)	Fractional precipitation	(B)	Partial dissolution
	(C)	Crystallization and distillation	(D)	Solvent extraction.
40.	Cro	sslinked PS can be obtained when s	tyrene	is polymerized along with small amount
	(A)	Divinyl benzene	(B)	Acrylonitrile
	(C)	Hydroquinone	(D)	Butadiene.
41.	Geo	metric isomerism was obtained due	to the	pressure of
	(A)	carbon-carbon double bond	(B)	carbon-carbon single bond
	(C)	asymmetric carbon atoms	(D)	symmetric carbon atoms.
42.	Aru	ibber that has good biocompatibility	is	
	(A)	NR .	(B)	PDMS
	(C)	NBR	(D)	PVC.

406			10	0		
43.	Mor	nomer to prepare nylon 6 is				
	(A)	Vinyl acetate		(B)	Caprolactum	
	(C)	Lactonide		(D)	Oxirane.	
44.	In s	suspension polymerization initiator	sho	ould be	orV o female unaversity but he state	
	(A)	media soluble		(B)	monomer insoluble	
	(C)	monomer soluble		(D)	monomer and media soluble.	
45.	Zeig	gler-Natta Catalyst is used to produ	ice			
	(A)	Stereoregular polymers		(B)	Branched polymers	
	(C)	Copolymers		(D)	Amorphous polymers.	
46.	Gel	point in condensation polymerizati	ion	is use	d to express	
	(A)	end of reaction		(B)	control of reaction rate	
	(C)	start of crosslinking	(cl)	(D)	start of degradation.	
47.	Atmospheric oxygen in polymerization reaction acts as a / an					
	(A)	Inhibitor		(B)	Initiator	
	(C)	Co-catalyst		(D)	Oxidising agent.	
48.	Chain carrier in case of anionic polymerization is					
	(A)	Carbanion		(B)	Carbonium ion	
	(C)	Carboxylate ion		(D)	Carbonyl.	
49.	Exp	andable polystyrene beads are pro	duc	ced by		
	(A)	bulk polymerization		(B)	interfacial polymerization	
	(C)	plasma polymerization		(D)	suspension polymerization.	
50.	Whi	ch of the following polymers is trai	nspa	arent '	? agent weld borg and had recon	
	(A)	Iso PP	100	(B)	HDPE	
	(C)	PMMA		(D)	ABS.	



PART - II

Each question carries two marks.

 $25 \times 2 = 50$

- 51. The specific weight of pure water is
 - (A) 1.24 kg/m³

(B) 1000 kg/m^3

(C) 9810 kg/cm³

- (D) $13.6 \times 10^{3} \text{ kg/m}^{3}$.
- 52. What is the mass in grams of 0.4 mole oxygen molecules?
 - (A) 12·8

(B) 12·0

(C) 16·0

- (D) 12·2.
- 53. A heat engine operates between a heat source at 700 K and a heat sink at 300 K.
 What is the maximum efficiency of the engine?
 - (A) 60%

(B) 52%

(C) 68%

- (D) 57%.
- 54. What is the change in entropy when 1 k mol of an ideal gas at 335 K and 10 bar is expanded irreversibly to 300 K and 1 bar? ($C_p = 29.3 \text{ kJ/mol.K}$)
 - (A) 15.0 kJ/k.mol.K

(B) 15.91 kJ/k.mol.K

(C) 16.9 kJ/k.mol.K

- (D) 16.2 kJ/k.mol.K.
- 55. Molecular weight of Na 2 CO 3 is
 - (A) 116

(B) 100

(C) 106

- (D) 110.
- 56. If a bucket holds 2 kg of NH 3, it contains
 - (A) 1 k mol of NH₃

(B) 0.117 k mol of NH 3

(C) 0.99 k mol of NH₃

- (D) 34 k mol of NH 3.
- 57. If PE has a molecular weight (M_n) of 2,80,000, what is its degree of polymerization?
 - (A) 100

(B) 1000

(C) 10000

(D) 2800.

- 58. Functionalities of butadiene and ethylene glycol are
 - (A) 2 and 2

(B) 4 and 2


(C) 4 and 3

- (D) 2 and 4.
- 59. The epoxy/glass fibre composite contains 40% of glass fibre. The densities of epox and glass fibre are 1.2 and 2.2 respectively. The density of the epoxy composite in the units of g/c.c. is
 - (A) 1.7


(B) 1·75


(C) 1.6

- (D) 1.55.
- 60. The structure of caprolactum is









- 61. Crystallinity of the three different types of PE follows the order
 - (A) HDPE > LLDPE > LDPE
- (B) LDPE > LLDPE > HDPE
- (C) HDPE > LDPE > LLDPE
- (D) LLDPE > LDPE > HDPE.
- 62. The temperature at which thermoforming is best carried out is
 - (A) softening temperature
 - (B) melting temperature
 - (C) glass transition temperature
 - (D) 20°C 30°C above melting temperature.

(B)

(D)

PTFE & PVDF

HDPE & PTFE.

PP & PS

(C) HDPE & PAN

- 68. Examples for free radical initiators are
 - (A) Benzoyl peroxide, BF 3, hydroquinone, AlCl 3
 - (B) Benzoyl peroxide, AIBN, potassium persulphate, H 2O 2
 - (C) AIBN, nitrobenzene, hydroquinone, DCP
 - (D) Benzoyl peroxide, AIBN, dinitrobenzene, lauroyl peroxide.
- 69. The rate constant for half change for a first order reaction is

(A)
$$k = \frac{1}{t_{1/2}}$$

(B)
$$\ln k = \frac{0.693}{t_{1/2}}$$

(C)
$$k = \frac{0.693}{t_{1/2}}$$

(D)
$$k = (t_{1/2})^2$$
.

- 70. The group of polymers consisting of PS, PMMA, PVA, HDPE and PAN is best categorised as
 - (A) Engineering polymers
 - (B) Polymers of addition polymerization
 - (C) Polymers of condensation polymerization
 - (D) Polymers of ring opening polymerization.
- 71. Which polymerization methods adopted to prepare nylon 66 and nylon 6?
 - (A) Condensation and ring opening
 - (B) Condensation and polyaddition
 - (C) Condensation and condensation
 - (D) Condensation and insertion.

- 72. How much chloride (chlorine) can be obtained from 58.5 kg of NaCl?
 - (A) 23

(B) 35·5

(C) 35·0

- (D) 37·0.
- 73. The starting material to prepare polyurethane is
 - (A) Diol + di-isocyanate

(B) Diol + diacid

(C) Diacid + diamine

- (D) Diol + diol.
- 74. The melting points of nylon 66 and nylon 6 are
 - (A) 260°C & 220°C

(B) 220°C & 200°C

(C) 260°C & 200°C

- (D) 220°C & 260°C.
- 75. Addition polymerization is also known as
 - (A) chain and vinyl polymerization
 - (B) chain and insertion polymerization
 - (C) chain and polyaddition polymerization
 - (D) chain and melt polycondensation.