CY: CHEMISTRY

Duration: Three Hours

Maximum Marks: 100

Read the following instructions carefully.

- 1. This question paper contains 24 pages including blank pages for rough work. Please check all pages and report discrepancy, if any.
- Write your registration number, your name and name of the examination centre at the specified locations on the right half of the Optical Response Sheet (ORS).
- Using HB pencil, darken the appropriate bubble under each digit of your registration number and the letters corresponding to your paper code.
- 4. All questions in this paper are of objective type.
- 5. Questions must be answered on the ORS by darkening the appropriate bubble (marked A, B, C. D) using HB pencil against the question number on the left hand side of the ORS. For each question darken the bubble of the correct answer. In case you wish to change an answer, crase the old answer completely. More than one answer bubbled against a question will be treated as an incorrect response.
- 6. There are a total of 65 questions carrying 100 marks.
- 7. Questions Q.1 Q.25 will carry 1-mark each, and questions Q.26 Q.55 will carry 2-marks each.
- 8. Questions Q.48 Q.51 (2 pairs) are common data questions and question pairs (Q.52, Q.53) and (Q.54, Q.55) are linked answer questions. The answer to the second question of the linked answer questions depends on the answer to the first question of the pair. If the first question in the linked pair is wrongly answered or is un-attempted, then the answer to the second question in the pair will not be evaluated.
- Questions Q.56 Q.65 belong to General Aptitude (GA). Questions Q.56 Q.60 will carry 1-mark each, and questions Q.61 - Q.65 will carry 2-marks each. The GA questions will begin on a fresh page starting from page 15.
- Un-attempted questions will carry zero marks.
- 11. Wrong answers will carry NEGATIVE marks. For Q.1 Q.25 and Q.56 Q.60, % mark will be deducted for each wrong answer. For Q.26 Q.51 and Q.61 Q.65, % mark will be deducted for each wrong answer. The question pairs (Q.52, Q.53), and (Q.54, Q.55) are questions with linked answers. There will be negative marks only for wrong answer to the first question of the linked answer question pair i.e. for Q.52 and Q.54, % mark will be deducted for each wrong answer. There is no negative marking for Q.53 and Q.55.
- Calculator (without data connectivity) is allowed in the examination hall.
- 13. Charts, graph sheets or tables are NOT allowed in the examination hall.
- 14. Rough work can be done on the question paper itself. Additionally, blank pages are provided at the end of the question paper for rough work.







Some Useful Data

1. Physical Constants


- (a) Planck Constant, $h = 6.626 \times 10^{-14} \text{ J s}$
- (b) Avogadro number, $N = 6.023 \times 10^{-23}$
- (c) Speed of light, $c = 3 \times 10^{8}$ m s⁻¹
- (d) Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mole}^{-1} = 2 \text{ cal K}^{-1} \text{ mole}^{-1}$

2. Atomic Numbers

В	5	Mn	25
Mg	12	Fc	26
P	15	Co	27
As	33	Cu	29
Sb	51	Ζπ	30
Bi	83	Мо	42
Ca	20		

Q.1 - Q.25 carry one mark each.

- Q.IThe Lewis acidity of BF3 is less than BCl3 even though fluorine is more electronegative than chlorine. It is due to
 - (A) stronger 2p(B)-2p(F) σ -bonding
- (B) stronger $2p(B)-2p(F)\pi$ -bonding
- (C) stronger 2p(B)-3p(Cl) \u03c3-bonding
- (D) stronger $2p(B)-3p(CI)\pi$ -bonding
- Q.2 Pyroxenes are a class of silicate minerals, which exhibit a polymeric chain structure, as shown below.



Its simplest repeat unit is

- (A) [SiO₄] 4
- (B) $[SiO_3]^{2-}$
- (C) [Si₂O₇]⁶⁻
- (D) $[Si_4O_{11}]^6$
- Q.3Among the following pentachlorides the one which does not exist due to the 'inert-pair effect' is
 - (A) PCl₃
- (B) BiCls
- (C) SbCl₅
- (D) AsCl₅
- Q.4 Band theory predicts that magnesium is an insulator. However, in practice it acts as a conductor due 10
 - (A) presence of filled 3s orbital
- (B) overlap of filled 2p and filled 3s orbital
- (C) overlap of filled 3s and empty 3p orbital (D) presence of unfilled 3p orbital
- The number of 'framework electron pairs' present in the borane cluster $\{B_{12}H_{12}\}^{2-}$ is Q.5
 - (A) 10
- (B) 11
- (C) 12
- (D) 13
- The reaction between [PdCl₄]²⁻ and C₂H₄ produces a new compound. Compared to free C₂H₄, the Q.6 C'-C' bond order of the product is
 - (A) between 1 and 2

(B) less than I

(C) unaltered

- (D) greater than 2
- Q.7 Among the following pair of metal ions present in Nature, the first one functions as an electrontransfer agent and the second one catalyzes the hydrolysis reactions. The correct pair is
 - (A) Fe and Zn

(B) Mg and Fe

(C) Co and Mo

- (D) Ca and Cu
- Q.8Structurally nickellocene is similar to ferrocene. Nickellocene attains stability due to the formation of
 - (A) a monocation

(B) a dication

(C) a monoanion


(D) a dianion


Q.9 The absolute configurations for compounds X and Y, respectively, are


- (A) R. S
- (B) S, R
- (C) R, R
- (D) S, S

Q.10 In the reaction

the major product [X] is









Q.11 Among the following, a pair of resolvable configurational enantiomers is given by


- (A) cis-1.2-dimethylcyclohexane
- (B) cis-1.3-dimethylcyclohexane
- (C) cis-1,4-dimethylcyclohexane
- (D) trans-1,3-dimethylcyclohexane


Q.12 In the reaction

the major product [X] is









Q.13 The decreasing order of isoelectric point for the following \alpha-amino acids is

(I)

Lysine

Alanine (II) Glutamic acid

(111)


(A) I > II > JJI

(B) II > I > III

(C) III > I > II

 $(D) \ 1 > III > II$

Q.14 The decreasing order of the reactivity of the following compounds towards electrophiles is





Π



Ш

I


111 < 1 < 11 (A)

(B) Ii > III > I

(C) iff > I > II


 $(D) 1 > \Pi > \Pi$


Q.15 In the reaction





the major product [X] is

- (A)
- (B)
- (C)
- (D)












Q.16 The decreasing order of acidity of the marked H of the following molecules is





[]



I

III

 $(A) \mid I > II > III$

(B) III > I > II

(C) III > II > 1

III < I < II (II)

CY

The decreasing order of nucleophilicity for the following anions is Q.17

- (A) $CH_3CO_2^- > CH_3O^- > C_6H_5O^- > NO_3^-$
- (B) $CH_1O' > NO_2' > C_2H_2O' > CH_2CO_2'$
- $(C) CH_1O^- > C_2H_2O^- > CH_1CO_- > NO_1^-$
- (D) $C_0H_3O^- > CH_3O^- > NO_7^- > CH_3CO_2^-$
- The molar entropy of crystalline CO at absolute zero is Q.18
 - (A) Zero
- (B) -Rln 2
- (C) Rin 2
- (D) 2RIn 2

- Q.19 For an ideal gas
 - (A) $(\partial P/\partial T)_V (\partial T/\partial V)_P (\partial V/\partial P)_T = 0$
 - $(B) (\partial P/\partial T)_{V} (\partial T/\partial V)_{P} (\partial V/\partial P)_{T} = -1$
 - (C) $(\partial P/\partial T)_V (\partial T/\partial V)_P (\partial V/\partial P)_T = +1$
 - (D) $(\partial P/\partial T)_V (\partial T/\partial V)_P (\partial V/\partial P)_T = +2$
- Among W (work), Q (heat), U (internal energy) and S (entropy) Q.20
 - (A) W and U are path functions but Q and S are state functions
 - (B) W and S are path functions but Q and U are state functions
 - (C) S and U are path functions but Q and W are state functions
 - (D) W and Q are path functions but U and S are state functions
- For eigen functions $\psi_1 = \sqrt{\frac{1}{b}} \sin\left(\frac{\pi x}{b}\right)$ and $\psi_2 = \sqrt{\frac{2}{b}} \sin\left(\frac{2\pi x}{b}\right)$ of particle in a 1-D box of length

 $b(0 \le x \le b)$

- (A) ψ₁ is normalized and orthogonal to ψ₂
- (B) ψ₁ is normalized but not orthogonal to ψ₂
- (C) ψ₂ is normalized and orthogonal to ψ₁
- (D) ψ₂ is neither normalized nor orthogonal to ψ₁
- Q.22 The bond order of C_2 molecule is
 - (A) 0
- (B) 1
- (C) 2
- (D) 3
- Sulfur can exist in four phases. The possible number of triple points is Q.23
 - (A) 1
- (B) 2
- (C) 3
- (D) 4
- The standard reduction potentials at 298 K for single electrodes are given below: Q.24

Electrode	Electrode Potential (volt) -2.34		
Mg ²⁺ /Mg			
Zn*2 / Zn	-0.76		
Fe ⁺² / Fe	-0.44		

From this we can infer that

- (A) Zn can reduce both Mg²⁺ and Fe²⁺
 (B) Fe can reduce both Mg²⁺ and Zn²⁺
 (C) Mg can reduce both Zn²⁺ and Fe²⁺

- (D) Mg can reduce Zn2+ but not Fe2+

Q.25 For the pair of reactions given below

i)
$$N_2(g) + 3H_2(g) = 2NH_1(g)$$

ii)
$$\frac{1}{2}N_2(g) + \frac{3}{2}H_2(g) \longrightarrow NH_3(g)$$

if at a particular temperature, K_{Pi} and K_{Pi} are the equilibrium constants for reactions i) and ii) respectively then,

(A)
$$K_{P_1} = 2K_{P_2}$$

(B)
$$K_{p_1} = K_{p_2}^2$$

(C)
$$2K_{p_1} = K_{p_2}$$

(D)
$$K_{P1}^1 = K_{P2}$$

Q.26 - Q.55 carry two marks each.

Q.26 According to VSEPR model, the shape of [XeOF₅] is

(A) octahedral

(B) trigonal bipyramidal

(C) square pyramidal

(D) pentagonal monopyramidal

Q.27 The number of unpaired electron(s) present in the species [Fe(H₂O)₃(NO)]²⁺ which is formed during 'brown ring test' is

- (A) 2
- (B) 3
- (C) 4
- (D) 5

Q.28 Fe₃O₄ and Co₃O₄ are metal oxides having spinel structure. Considering their CFSEs, the correct statement regarding their structure is

- (A) both have normal spinel structure
- (B) both have inverse spinel structure
- (C) Fe₃O₄ has normal and Co₃O₄ has inverse spinel structure
- (D) Fe₃O₄ has inverse and Co₃O₄ has normal spinel structure

Q.29 The mechanism of the reaction between $[Fe(CN)_b]^4$ and $[Fe(bpy)_3]^{3+}$ (bpy = 2,2'-bipyridine) is

- (A) outer-sphere electron-transfer
- (B) inner-sphere electron-transfer
- (C) self-exchange reaction
- (D) ligand-exchange followed by electron-transfer

Q.30 The d-d absorption band of [Fe(H2O)6]2+ is split due to

- (A) presence of octahedral geometry
- (B) static Jahn-Teller distortion
- (C) dynamic Jahn-Teller distortion
- (D) presence of trigonal bipyramidal geometry

Q.31 The crystal-field symbol for the ground-state of [Mn(CN)₆]⁴ is

- $(A)^{2}T_{2p}$
- (B) A12
- (C) 5E,
- (D) Aig



7/24

In the following reactions: Q.32

the reagent/conditions X and Y are

- (A) $X = BF_3$; Y = heating at 1250°C
- (B) X = NaF; Y = heating at 250°C

(C) $X = NH_4F$; Y = HCI

- (D) $X = CF_3SO_3H$; $Y = H_2SO_4$
- Q.33 [CoCl₄]²⁻ is a blue coloured complex. Controlled-treatment of this complex with water generates two isomeric light pink coloured complexes of composition $[Co(H_2O)_4Cl_2]$. Identify the correct point groups for $[CoCl_4]^{2-}$ and two isomeric complexes $[Co(H_2O)_4Cl_2]$.
 - (A) D_{4h} and $(C_{2v}$ and $C_{2h})$

(B) T_d and (C_{2v} and D_{4h})

(C) $D_{4\rm h}$ and (C $_{2\nu}$ and $D_{4\rm h})$

(D) T_d and $(C_{2v}$ and $C_{4v})$

In the reaction Q.34

EtO₂C
$$CO_2H$$
 $i. B_2H_6$ [X]


the major product [X] is


(A)




(C)

(D)








In the reaction Q.35


the major product [X] is

(A)

(B)

(C)





Q.36 In the following reaction sequence

the major product [X] is

Q.37 In the reactions

the major products, [X] and (Y), respectively, are

(D) and
$$\downarrow$$
 coordinate \downarrow and \downarrow coordinate \downarrow

Q.38 In the reaction


the major product (X) is

(A)



(B)


(C)




Q.39 In the reaction sequence

the major products. [X] and [Y], respectively, are


(A)




(B)



(C)



(D)



Q.40 The change in entropy when two moles of Argon gas are heated at constant volume from 300 K to 500 K is

(A) -12.74 J K⁻¹ mole⁻¹ (C) 6.37 J K⁻¹ mole⁻¹

(B) -6.37 J K⁻¹ mole⁻¹ (D) 12.74 J K⁻¹ mole⁻¹

CY

10/24

At any temperature T, the fugacity coefficient (γ) is given by Q.41

$$\ln \gamma = \int_{0}^{P} \frac{Z - 1}{P'} dP'$$

where Z is the compressibility factor. The fugacity coefficient of a real gas governed by equation of state P(V-b) = RT with b a constant is given by

(A)
$$\frac{RT}{bP}$$

(B)
$$e^{\frac{RT}{bP}}$$

(A)
$$\frac{RT}{bP}$$
 (B) $e^{\frac{RT}{bP}}$ (C) $\frac{bP}{RT}$

(D)
$$e^{\frac{hP}{RT}}$$

The specific rate constant of decomposition of a compound is represented by

$$\ln k = 5.0 - \frac{12000}{T}$$

The activation energy of decomposition for this compound at 300 K is

- (A) 24 kcal/mole
- (B) 12 kcal/mole
- (C) 24 cal/mole
- (D) 12 cal/mole

The commutator (x3, p1) is equal to

(A)
$$-\frac{3hx^2}{2\pi i}$$
 (B) $\frac{hx}{2\pi i}$ (C) $\frac{hx^2}{2\pi i}$ (D) $\frac{3hx^2}{2\pi i}$

(B)
$$\frac{hx}{2\pi i}$$

(C)
$$\frac{hx^2}{2\pi i}$$

(D)
$$\frac{3hx^2}{2\pi i}$$

An electron of mass 'm' is confined to a one dimensional box of length 'b'. If it makes a radiative Q.44 transition from second excited state to the ground state, the frequency of the photon emitted is

(A)
$$\frac{9h}{8mb^2}$$

(B)
$$\frac{3h}{8mb^2}$$
 (C) $\frac{h}{mb^2}$ (D) $\frac{2h}{mb^2}$

(C)
$$\frac{h}{mb^2}$$

(D)
$$\frac{2h}{mb^2}$$

The point group of CIF3 molecule and its corresponding number of irreducible representations are respectively

- (A) Cac and 4
- (B) C_{2v} and 4 (C) C_{3v} and 3
- (D) C₂, and 3

Q.46 The most populated rotational state for HCl (B = 8.5 cm⁻¹) at 300 K is

- (A) 2
- (B) 3
- (C) 5
- (D) 7

The ratio of life times of two states that give rise to line widths of 1.0 cm⁻¹ and 0.2 cm⁻¹ respectively Q.47

- (A) 1:2
- (B) 1:5
- (C) 2:1
- (D) 5:1

Common Data Questions

Common Data for Questions 48 and 49:

A six-coordinate transition-metal complex is ESR and Mösshauer active. The effective magnetic moment of this complex is ~5.9 B.M.

The metal-ion along with its oxidation state and the number of unpaired electron present are

(A) Fe(II) and 4

(B) Mn(II) and 5

(C) Fe(III) and 1

(D) Fc(III) and 5

The complex is Q.49

- (A) $[Mn(H_2O)_6]^{2+}$
- (B) $[Fe(CN)_6]^3$
- (C) $[Fe(H_2O)_6]^{24}$
- (D) $[Fe(H_2O)_6]^{3}$

Common Data for Questions 50 and 51:

An organic compound [X] ($C_{12}H_{16}O_3$) exhibits the following spectral data IR: ~1720 cm⁻¹

^tH NMR: 2.35 (s. 6H), 3.30 (s, 3H), 3.83 (t, 2H), 4.42 (t, 2H), 7.07 (s, 1H), 7.58 (s, 2H)

The compound [X] with an excess of MeMgBr gives a 1:1 mixture of compounds [Y] and [Z]. The compound [Z] exhibits the following ¹H NMR data: 2.0 (bs. 1H), 3.30 (s, 3H), 3.56 (t, 2H), 3.70 (t, 2H)

Q.50 The compound [X] is

Q.51 The compound [Y] is

Linked Answer Questions

Statement for Linked Questions 52 and 53:

In the reaction sequence


Q.52 The compound [X] is

Q.53 The compound [Z] is

2010 C'Y

Statement for Linked Questions 54 and 55:

In the μ vs. T diagram for different phases of the same substance at one atmospheric pressure, the lines A. B and C compound to



Q.54 Based on the above diagram:

- (A) A represents the change in chemical potential as a function of temperature for the solid phase.

 B for the liquid and C for the gas
- (B) A represents the change in chemical potential as a function of temperature for the liquid phase, B for the gas and C for the solid
- (C) A represents the change in chemical potential as a function of temperature for the gas phase. B for the liquid and C for the solid
- (D) A represents the change in chemical potential as a function of temperature for the gas phase.

 B for solid and C for the liquid

Q.55 From the same diagram

- (A) D represents boiling point. E sublimation point and F melting point
- (B) E represents boiling point, D sublimation point and F melting point
- (C) E represents melting point, F sublimation point and D boiling point
- (D) D represents melting point, F boiling point and E sublimation point

collegedunia

2010					
Gener	ral Aptitude (C	GA) Questions			
Q.56 -	- Q.60 сагту о	ne mark each.			
Q.56	25 persons are in a room. 15 of them play hockey. 17 of them play football and 10 of them play both hockey and football. Then the number of persons playing neither hockey nor football is:				
	(A) 2	(B) 17	(C) 13	(D) 3	
Q.57	sentence:	Cor a	from the options give		2
	(A) uphold (B) restrain (C) cherish (D) conserve				
Q.58		epresses the relation in	of related words follow the original pair.	ed by four pairs of word	ls. Select the
	(A) fallow: land (B) unaware: si (C) wit: jester (D) renovated:	eeper			
Q.59	Which of the fol	lowing options is the c	closest in meaning to the	word below:	
	(A) cyclic (B) indirect (C) confusing (D) crooked				
Q.60	sentence:		from the options give	The state of the s	1000 (0.100000-1000 D- 40.00
	(A) masked (B) belied (C) betrayed (D) suppressed				
Q.61 -	- Q.65 carry to	wo marks each.			
Q.61	1" January, The is less than 3 ye	age difference between ars. Given the following Hari's age + Gita's	age > Irfan's age + Saira between Gita and Saira i	blings (that is born one a a's age.	fter another)

iii. There are no twins.

(B) SGHI

In what order were they born (oldest first)?

(D) IHSG

(C) IGSH

(A) HSIG

		200				
Q.62	5 skilled workers can build a wall in 20 days; 8 semi-skilled workers can build a wall in 25 days; 10 unskilled workers can build a wall in 30 days. If a team has 2 skilled, 6 semi-skilled and 5 unskilled workers, how long will it take to build the wall?					
	(A) 20 days	(B) 18 days	(C) 16 days	(D) 15 days		
Q.63	Modern warfare has changed from large scale clashes of armies to suppression of civilian populations. Chemical agents that do their work silently appear to be suited to such warfare; and regretfully, there exist people in military establishments who think that chemical agents are useful tools for their cause.					
	Which of the following statements best sums up the meaning of the above passage:					
	 (A) Modern warfare has resulted in civil strife. (B) Chemical agents are useful in modern warfare. (C) Use of chemical agents in warfare would be undesirable. (D) People in military establishments like to use chemical agents in war. 					
Q.64	Given digits 2, 2, 3, 3, 3, 4, 4, 4, 4 how many distinct 4 digit numbers greater than 3000 can be formed?					
	(A) 50	(B) 51	(C) 52	(D) 54		
Q.65	If 137 + 276 = 435 how much is 731 + 672?					
	(A) 534	(B) 1403	(C) 1623	(D) 1513		

END OF THE QUESTION PAPER









2610 ______CY





collegedunia India's largest Student Review Platform

CY

Space for Rough Work



21/24









.

