

Question No.	Questions
1.	<p>Which one of the following high spin complexes has the largest CSFE Crystal field stabilization energy</p> <p>(1) $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$ (2) $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$ (3) $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ (4) $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$</p>
2.	<p>The number of 3c, 2e BHB and B-B bonds present in B_4H_{10} respectively are</p> <p>(1) 2, 4 (2) 3, 2 (3) 4, 1 (4) 4, 0</p>
3.	<p>The most unstable species among the following is</p> <p>(1) $\text{Ti}(\text{C}_2\text{H}_5)_4$ (2) $\text{Ti}(\text{CH}_2\text{Ph})_4$ (3) $\text{Pb}(\text{CH}_3)_4$ (4) $\text{Pb}(\text{C}_2\text{H}_5)_4$</p>
4.	<p>The acid catalyzed hydrolysis of $\text{trans-}[\text{Co}(\text{en})_2\text{AX}]^{n+}$ can give <i>cis</i>-product also due to the formation of</p> <p>(1) Square pyramidal intermediate (2) Trigonal bipyramidal intermediate (3) Pentagonal bipyramidal intermediate (4) Face capped octahedral intermediate</p>
5.	<p>Total number of lines expected in ^{31}P NMR spectrum of HPF_2 is ($I = 1/2$ for both ^{19}F and ^{31}P)</p> <p>(1) Six (2) Four (3) Five (4) Three</p>

Question No.	Questions
20.	In the iodometric titration of sodium thiosulfate ($\text{Na}_2\text{S}_2\text{O}_3$) with acidic dichromate solution, 25 mL of 0.1 M dichromate requires 50 mL of 'x' M thiosulfate. The value of 'x' is (1) 0.6 (2) 0.3 (3) 0.1 (4) 0.4
21.	The room temperature magnetic moment (μ_{eff} in BM) for a monomeric Cu(II) complex is greater than 1.73. This may be explained using the expression (1) $\mu_{\text{eff}} = \mu_s (1 - \alpha\lambda/\Delta)$ (2) $\mu_{\text{eff}} = [n(n+2)]^{1/2}$ (3) $\mu_{\text{eff}} = [4s(s+1) + L(L+1)]^{1/2}$ (4) $\mu_{\text{eff}} = g[J(J+1)]^{1/2}$
22.	The numbers of P-S and P-P bonds in the compound P_4S_3 are, respectively, (1) 3 and 6 (2) 4 and 3 (3) 6 and 3 (4) 6 and 2
23.	In the absence of bound globin chain, heme group on exposure to O_2 gives the iron-oxygen species (1) Fe(III)-O-Fe(III) (2) Fe(III)-O-O^- (3) $\text{Fe(III)-O-O-Fe(III)}$ (4) Fe(IV)-O-
24.	The complex $[\text{Cr}(\text{bipyridyl})_3]^{2+}$, shows a red phosphorescence due to transition (1) ${}^4\text{T}_{1g} \leftarrow {}^4\text{A}_{2g}$ (2) ${}^2\text{E}_g \leftarrow {}^4\text{A}_{2g}$ (3) ${}^4\text{T}_{2g} \leftarrow {}^4\text{A}_{2g}$ (4) ${}^4\text{A}_{2g} \leftarrow {}^2\text{E}_g$

Question No.	Questions
38.	Due to Frenkel defect, the density of the ionic solids (1) increases (2) decreases (3) does not change (4) none of the above
39.	What is the simplest formula of a solid whose cubic unit cell has the atom A at each corner, the atom B at each face centre and a C atom at the body centre (1) AB_2C (2) A_2BC (3) AB_3C (4) ABC_3
40.	Which of the following thermodynamic function is called as the arrow of "time" (1) Enthalpy (2) Gibbs free energy (3) Entropy (4) Helmholtz free energy
41.	For a potentiometric titration in the curve of emf (E) v/s volume (V) of the titrant added, the equivalence point is indicated by (1) $ dE/dV = 0, d^2E/dV^2 = 0$ (2) $ dE/dV = 0, d^2E/dV^2 > 0$ (3) $ dE/dV > 0, d^2E/dV^2 = 0$ (4) $ dE/dV > 0, d^2E/dV^2 > 0$
42.	If the concentration (c) is increased to 4 times its original value (c), the change in molar conductivity for strong electrolytes is (where b is Kohlrausch's constant) - (1) 0 (2) $b\sqrt{c}$ (3) $2b\sqrt{c}$ (4) $4b\sqrt{c}$



Question No.	Questions																																			
43.	<p>The energy levels of the harmonic oscillator (neglecting zero point energy) are $\epsilon_n = nh\nu$ for $n = 0, 1, 2, \dots$. Assuming $h\nu = k_B T/3$; the partition function is</p> <p>(1) e (2) $e^{1/3} (e^{1/3} - 1)$ (3) $1/3e$ (4) $3e/(3e^3 - 1)$</p>																																			
44.	<p>The ground state of hydrogen atom is -13.598 eV. The expectation values of kinetic energy $\langle T \rangle$ and potential energy, $\langle V \rangle$, in units of eV, are</p> <p>(1) $\langle T \rangle = 13.598, \langle V \rangle = -27.196$ (2) $\langle T \rangle = -27.196, \langle V \rangle = 13.598$ (3) $\langle T \rangle = -6.799, \langle V \rangle = -6.799$ (4) $\langle T \rangle = 6.799, \langle V \rangle = -20.397$</p>																																			
45.	<p>The correct expression for the product $(\overline{M}_n)(\overline{M}_w)$ [where \overline{M}_n and \overline{M}_w are the number average and weight average molar masses, respectively, of a polymer] is</p> <p>(1) $N^{-1} \sum_i N_i M_i$ (2) $N^{-1} \sum_i N_i M_i^2$ (3) $N / \sum_i N_i M_i$ (4) $N / \sum_i N_i M_i^2$</p>																																			
46.	<p>Match the following columns :</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%; text-align: center;">Column-1</th> <th style="width: 50%; text-align: center;">Column-2</th> </tr> </thead> <tbody> <tr> <td>A. Energy of the ground state of He+</td> <td>1. -6.04 eV</td> </tr> <tr> <td>B. Potential energy of 1st orbit of H-atom</td> <td>2. -27.2 eV</td> </tr> <tr> <td>C. Kinetic energy of II excited state of He+</td> <td>3. 8.68×10^{-18} J</td> </tr> <tr> <td>D. Ionisation potential of He+</td> <td>4. -54.4 eV</td> </tr> </tbody> </table> <p>Codes.</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th>A</th> <th>B</th> <th>C</th> <th>D</th> </tr> </thead> <tbody> <tr> <td>(1)</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> </tr> <tr> <td>(2)</td> <td>4</td> <td>3</td> <td>2</td> <td>1</td> </tr> <tr> <td>(3)</td> <td>4</td> <td>2</td> <td>1</td> <td>3</td> </tr> <tr> <td>(4)</td> <td>2</td> <td>3</td> <td>1</td> <td>4</td> </tr> </tbody> </table>	Column-1	Column-2	A. Energy of the ground state of He+	1. -6.04 eV	B. Potential energy of 1st orbit of H-atom	2. -27.2 eV	C. Kinetic energy of II excited state of He+	3. 8.68×10^{-18} J	D. Ionisation potential of He+	4. -54.4 eV		A	B	C	D	(1)	1	2	3	4	(2)	4	3	2	1	(3)	4	2	1	3	(4)	2	3	1	4
Column-1	Column-2																																			
A. Energy of the ground state of He+	1. -6.04 eV																																			
B. Potential energy of 1st orbit of H-atom	2. -27.2 eV																																			
C. Kinetic energy of II excited state of He+	3. 8.68×10^{-18} J																																			
D. Ionisation potential of He+	4. -54.4 eV																																			
	A	B	C	D																																
(1)	1	2	3	4																																
(2)	4	3	2	1																																
(3)	4	2	1	3																																
(4)	2	3	1	4																																

Code-A

Question No.	Questions
47.	The protecting power of lyophilic colloidal sol is expressed in terms of (1) Critical miscelle concentration (2) Oxidation number (3) Coagulation value (4) Gold number
48.	Which one of the following is an example for homogenous catalysis ? (1) Hydrogenation of oil (2) Manufacture of ammonia by Haber's process (3) Manufacture of sulphuric acid by Contact process (4) Hydrolysis of sucrose in presence of dilute hydrochloric acid
49.	The energy of a hydrogen atom in a state is $(-hcR_H/25)$, where R_H = Rydberg Constant). The degeneracy of the state will be - (1) 25^1 (2) 25^2 (3) 25^3 (4) 25^4
50.	The value of the commutator $[x, p_x^2]$ is (1) $2i$ (2) $2i\hbar p_x$ (3) $2ixp_x$ (4) $\hbar ip_x/\pi$
51.	The number of the lines in the ESR spectrum of CD_3 is (the spin of D is 1). (1) 1 (2) 3 (3) 4 (4) 7
52.	Colligative properties are used for the determination of (1) molar mass (2) equivalent weight (3) arrangement of molecules (4) melting and boiling point



Question No.	Questions
53.	Which of the following does not contain a C_3 axis ? (1) $POCl_3$ (2) NH_4^+ (3) H_3O^+ (4) ClF_3
54.	Franck Condon principle is related to (1) time required for electronic transition to occur (2) absorption of light (3) time of electronic transition and change in internuclear distance (4) symmetry of molecules
55.	Which pairing of molecule and point group is correct ? (1) BCl_3, C_{3v} (2) $SiCl_4, D_{4h}$ (3) H_2S, C_{2v} (4) SF_4, C_{4v}
56.	The symmetric stretching mode of the SiF_4 molecule : (1) IR active (2) IR inactive (3) generates a change in molecular dipole moment (4) gives rise to a strong absorption in IR spectrum
57.	Match the following columns : LIST-1 1. Sol 2. Gel 3. Emulsion 4. Foam Codes (1) 1-A 2-B 3-C 4-D (2) 1-B 2-C 3-D 4-A (3) 1-C 2-A 3-D 4-B (4) 1-B 2-D 3-A 4-C LIST-2 A. Liquid dispersed in solid B. gas dispersed in liquid C. Solid dispersed in liquid D. liquid dispersed in liquid


Question No.	Questions
72.	Keto-enol tautomerism is observed in : (1) $\text{C}_6\text{H}_5-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3$ (2) $\text{C}_6\text{H}_5-\overset{\text{O}}{\parallel}{\text{C}}-\text{C}_6\text{H}_5$ (3) $\text{CH}_3\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ (4) $\text{C}_6\text{H}_5-\overset{\text{O}}{\parallel}{\text{C}}-\text{H}$
73.	Which of the following gases is mainly responsible for acid rain ? (1) NO_2 and CO_2 (2) CO_2 and SO_2 (3) SO_2 and NO_2 (4) None of these
74.	Which of the following compound displays two singlets at $\delta_{2,3}$ and 7.1 ppm. (1) 1, 2-dimethylbenzene (2) 1, 3-dimethyl benzene (3) 1, 4-dimethyl benzene (4) methyl benzene
75.	A single strong and sharp absorption near 1650 cm^{-1} in IR spectra indicates the presence of (1) Acid chlorides (2) Amides (3) Anhydrides (4) Aldehydes
76.	The proteins in which prosthetic group is carbohydrate are known as (1) Lipo-protein (2) Mucoprotein (3) Chromoprotein (4) Nucleoprotein

Code-A

Question No.	Questions		
77.	Match the List I and List II and select the correct answer using codes given below : <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top;"> List I 1 Nerol 2 Citral 3 Pinol 4 Lupeol </td> <td style="width: 50%; vertical-align: top;"> List II A Lemon grass oil B Geraniol C Amyrin D α-pinene </td> </tr> </table> Correct answer is : (1) 1-C, 2-B, 3-A, 4-D (2) 1-B, 2-A, 3-D, 4-C (3) 1-D, 2-C, 3-A, 4-D (4) 1-A, 2-D, 3-B, 4-D	List I 1 Nerol 2 Citral 3 Pinol 4 Lupeol	List II A Lemon grass oil B Geraniol C Amyrin D α -pinene
List I 1 Nerol 2 Citral 3 Pinol 4 Lupeol	List II A Lemon grass oil B Geraniol C Amyrin D α -pinene		
78.	Hydrolysis product of sucrose is : (1) Fructose (2) Glucose + Galactose (3) Glucose (4) Glucose + Fructose		
79.	The mass spectrum of primary amides shows a moderate molecular ion and an Intense peak at $m/z = 44$ due to : (1) Loss of an alkyl radical (2) Loss of HCN (3) Loss of CO (4) Loss of methyl radical		
80.	Which one of the following is bacteriostatic drug ? (1) Chloramphenicol (2) Penicillin (3) Streptomycin (4) Phenacetin		
81.	Heating 1, 4-dicarbonyl compounds in the presence of phosphorus pentoxide (P_2O_5) gives : (1) Pyrrole (2) Furan (3) Thiophene (4) Quinoline		
82.	The Acetylation of thiophene occurs at : (1) C_3 -position (2) C_4 -position (3) C_2 -position (4) both at C_2 and C_4 -positions		

Question No.	Questions
83.	Pyridine is basic in nature having (1) $pK_a = 5.21$ (2) $pK_a = -0.27$ (3) $pK_a = 5.81$ (4) $pK_a = -0.35$
84.	Least stable carbocation among the following is (1) $(CH_3)_3C^+$ (2) $(CH_3)_2CH^+$ (3) $CH_3CH_2^+$ (4) CH_3^+
85.	Due to the presence of an unpaired electron, free radicals are (1) Anions (2) Cations (3) Chemically reactive (4) Chemically inactive
86.	Benzoyl peroxide undergoes homolytic cleavage to produce (1) Phenyl radical (2) Methyl radical (3) Phenyl chloride (4) Methyl chloride
87.	S_N1 mechanism for the hydrolysis of an alkyl halide involves the formation of intermediate (1) Free radical (2) Carbanion (3) Carbocation (4) None of these
88.	Which of the following is <u>NOT</u> polar protic solvent ? (1) H_2O (2) C_2H_5OH (3) Fumaric acid (4) Acetone
89.	A new carbon-carbon bond formation is possible in (1) Clemmensen reduction (2) Wurtz reduction (3) Friedel-Craft alkylation (4) Oppenauer oxidation

Question No.	Questions
90.	Give the name of reaction given below :  (1) Perkin reaction (2) Pechmann condensation (3) Benzoin condensation (4) Claisen-Schmidt reaction
91.	What is meant by a reaction going in 94% enantiomeric excess ? (1) The product contains 94% of one enantiomer and 6% of other enantiomer (2) The product contains an enantiomer which is 94% pure (3) The product contains 94% of one enantiomer and 6% of the products (4) The product contains 97% of one enantiomer and 3% of other enantiomer .
92.	Which of the following functional group is <u>not</u> reduced by sodium borohydride (NaBH ₄) (1) >C=O (2) $\begin{array}{c} \text{-C-Cl} \\ \\ \text{O} \end{array}$ (3) $\begin{array}{c} \text{-C-H} \\ \\ \text{O} \end{array}$ (4) $\begin{array}{c} \text{-C-OH} \\ \\ \text{O} \end{array}$
93.	The given reaction is the example of :  (1) 2 + 4 cycloaddition (2) 2 + 2 cycloaddition (3) 2 + 2 + 2 cycloaddition (4) 2S + 2S cycloaddition
94.	A photo chemical reaction is : (1) catalysed by light (2) Initiated by light (3) accompanied with the emission of light (4) used to convert heat energy into light

Question No.	Questions
95.	Which of the following solvents is unacceptable on large scale ? (1) Dimethoxy ethane (2) Xylene (3) Diethyl ether (4) Heptane
96.	For the reaction given below, which reaction condition are not suitable ? <div style="text-align: center;">  </div> (1) $\text{LiAlH}_4 / \text{Et}_2\text{O}$ (2) $\text{H}_2\text{N NH}_2 / \text{NaOH}$ (3) Zn (Hg) / HCl (4) $\text{HSCH}_2\text{CH}_2\text{CH}_2\text{SH} / \text{H}^+, \text{H}_2 / \text{Ni}$
97.	Which of the following statements is <u>not</u> correct ? (1) The molecule to be synthesised is a target molecule. (2) Synthetic equivalent is a real chemical compound resulting from disconnection (3) Regioselective reaction does not produce one of several possible structural isomers (4) Synthons are idealised fragments (usually cation or anion) resulting from a disconnection.
98.	How many oxygen atoms lined up in a row would fit in a one nanometre space ? (1) Seventy (2) One (3) Seven (4) None
99.	The role of catalyst in chemical reaction is (1) Lowers the activation energy (2) Alters the amount of products (3) Increases ΔH of Forward reaction (4) Decreases ΔH of Forward reaction
100.	Secondary pollutant is (1) SO_2 (2) CO (3) PAN (4) Aerosol

