


PHYSICS

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer:

1. Bob *P* is released from the position of rest at the moment shown. If it collides elastically with an identical bob *Q* hanging freely then velocity of *Q*, just after collision is ($g = 10 \text{ m/s}^2$)



- (1) 1 m/s (2) 4 m/s
(3) 2 m/s (4) 8 m/s

Answer (3)

Sol. Velocity of *P* just before collision is $= \sqrt{2gl}$
 $= 2 \text{ m/sec}$

As collision is elastic and the mass of *P* and *Q* are equal therefore just after collision velocity of *P* is 0 and that of *Q* is 2 m/sec.

2. Choose the option showing the correct relation between Poisson's ratio (σ), Bulk modulus (*B*) and modulus of rigidity (*G*).

- (1) $\sigma = \frac{3B - 2G}{2G + 6B}$ (2) $\sigma = \frac{6B + 2G}{3B - 2G}$
(3) $\sigma = \frac{9BG}{3B + G}$ (4) $B = \frac{3\sigma - 3G}{6\sigma + 2G}$

Answer (1)

Sol. $E = 2G(1 + \sigma)$ (1)

$E = 3B(1 - 2\sigma)$ (2)


$$1 = \frac{2G}{3B} \left(\frac{1 + \sigma}{1 - 2\sigma} \right)$$

$$\Rightarrow 3B - 6B\sigma = 2G + 2G\sigma$$

$$\Rightarrow 3B - 2G = \sigma (2G + 6B)$$

$$\sigma = \left(\frac{3B - 2G}{2G + 6B} \right)$$

3. Two conducting solid spheres (*A* & *B*) are placed at a very large distance with charge densities and radii as shown:



When the key *K* is closed, find the ratio of final charge densities.

- (1) 4 : 1 (2) 1 : 2
(3) 2 : 1 (4) 1 : 4

Answer (3)


Sol. Final potential is same

$$\Rightarrow \frac{1}{4\pi\epsilon_0} \frac{Q_1}{R} = \frac{1}{4\pi\epsilon_0} \frac{Q_2}{2R} \quad \dots(1)$$

$$\text{Also, } Q_1 + Q_2 = \sigma \cdot 4\pi R^2 + \sigma \cdot 4\pi(2R)^2 \quad \dots(2)$$

$$\Rightarrow \frac{\sigma_1}{\sigma_2} = 2.$$

4. Position-time graph for a particle is parabolic and is as shown:



Choose the corresponding *v* - *t* graph

- (1) (2)
(3) (4)

Answer (2)

Sol. Since $x \propto t^2$

$$\Rightarrow v = \frac{dx}{dt} \propto t'$$

\Rightarrow Option 2 is correct

5. For a system undergoing isothermal process, heat energy is supplied to the system. Choose the option showing correct statements
- (a) Internal energy will increase
 - (b) Internal energy will decrease
 - (c) Work done by system is positive
 - (d) Work done by system is negative
 - (e) Internal energy remains constant
- (1) (a), (c), (e) (2) (b), (d)
 (3) (c), (e) (4) (a), (d), (e)

Answer (3)

Sol. For isothermal process,

$$dT = 0$$

so, $dU = 0 \Rightarrow$ Internal energy remains same

$$dQ = dW$$

as dQ is positive,

so dW is positive

6. The heat passing through the cross-section of a conductor, varies with time ' t ' as $Q(t) = \alpha t - \beta t^2 + \gamma t^3$. (α , β and γ are positive constants.) The minimum heat current through the conductor is

- (1) $\alpha - \frac{\beta^2}{2\gamma}$ (2) $\alpha - \frac{\beta^2}{3\gamma}$
 (3) $\alpha - \frac{\beta^2}{\gamma}$ (4) $\alpha - \frac{3\beta^2}{\gamma}$

Answer (2)

Sol. Heat through cross section of rod

$$Q = \alpha t - \beta t^2 + \gamma t^3$$

$$\text{so heat current} = \frac{dQ}{dt}$$

$$\text{heat current} = \frac{dQ}{dt} = \alpha - 2\beta t + 3\gamma t^2$$

for heat current to be minimum

$$\frac{d^2Q}{dt^2} = -2\beta + 6\gamma t = 0$$

$$t = \frac{2\beta}{6\gamma} = \left(\frac{\beta}{3\gamma} \right)$$


so minimum heat current

$$\left. \frac{dQ}{dt} \right|_{\text{minimum}} = \alpha - 2\beta \times \frac{\beta}{3\gamma} + 3\gamma \times \frac{\beta^2}{9\gamma^2}$$

$$= \alpha - \frac{2\beta^2}{3\gamma} + \frac{\beta^2}{3\gamma}$$

$$= \left(\alpha - \frac{\beta^2}{3\gamma} \right)$$


7. Momentum-time graph of an object moving along a straight line is as shown in figure. If $(P_2 - P_1) < P_1$ and $(t_2 - t_1) = t_1 < (t_3 - t_2)$ then at which points among A, B and C the magnitude of force experienced by the object is maximum and minimum respectively.



- (1) A, B (2) A, C
 (3) B, C (4) B, A

Answer (2)

Sol.



$$F_A = \frac{P_1}{t_1}$$

$$F_B = \frac{P_2 - P_1}{t_2 - t_1}$$

$$F_C = \frac{P_2 - P_1}{t_3 - t_2}$$


Therefore the maximum force is at A and minimum force is at C.

8. A particle moving in unidirectional motion travels half of the total distance with a constant speed of 15 m/s. Now first half of the journey time it travels at 10 m/s and second half of the remaining journey time it travels at 5 m/s. Average speed of the particle is

- (1) 12 m/s (2) 10 m/s
 (3) 7 m/s (4) 9 m/s

Answer (2)

Sol.




$$v_{av} = \frac{2x}{\frac{x}{15} + 2t}$$

$$= \frac{2x}{\frac{x}{15} + \frac{2x}{10+5}}$$

$$= 10 \text{ m/sec}$$

9. A bullet strikes a stationary ball kept at a height as shown. After collision, range of bullet is 120 m and that of ball is 30 m. Find initial speed of bullet. Collision is along horizontal direction.

Take $g = 10 \text{ m/s}^2$



- (1) 150 m/s (2) 90 m/s
(3) 240 m/s (4) 360 m/s

Answer (4)

Sol. $m_1v + m_2(0) = m_1v_1' + m_2v_2'$... (1)

$$\Delta t = \sqrt{\frac{2h}{g}} = 2s \quad \dots(2)$$

$$\Rightarrow v_1' = \frac{120 \text{ m}}{2s} = 60 \text{ m/s}$$


$$\& v_2' = \frac{30 \text{ m}}{2s} = 15 \text{ m/s}$$

$$\Rightarrow v = 360 \text{ m/s}$$

10. If an inductor with inductive reactance, $X_L = R$ is connected in series with resistor R across an A.C voltage, power factor comes out to be P_1 . Now, if a capacitor with capacitive reactance, $X_C = R$ is also connected in series with inductor and resistor in the same circuit, power factor becomes P_2 . Find $\frac{P_1}{P_2}$

- (1) $\sqrt{2} : 1$ (2) $1 : \sqrt{2}$
(3) $1 : 1$ (4) $1 : 2$

Answer (2)




$$Z = \sqrt{R^2 + R^2}$$

$$= \sqrt{2}R$$

$$P_1 = \cos\phi = \text{power factor} = \frac{R}{Z} = \left(\frac{1}{\sqrt{2}}\right)$$

When capacitor is also connected in series



The LCR circuit is in resonance stage

$$\text{So, } Z = \sqrt{R^2 + (X_L - X_C)^2}$$

$$Z = R$$

$$P_2 = \cos\phi = \text{power factor} = \frac{R}{Z} = \frac{R}{R} = 1$$

$$\text{So, } \frac{P_1}{P_2} = \frac{\left(\frac{1}{\sqrt{2}}\right)}{1} = \frac{1}{\sqrt{2}}$$

11. Electromagnetic wave beam of power 20 mW is incident on a perfectly absorbing body for 300 ns. The total momentum transferred by the beam to the body is equal to

- (1) $2 \times 10^{-17} \text{ Ns}$ (2) $1 \times 10^{-17} \text{ Ns}$
(3) $3 \times 10^{-17} \text{ Ns}$ (4) $5 \times 10^{-17} \text{ Ns}$

Answer (1)

Sol. Total energy incident = Pt

$$\text{So total initial momentum} = \frac{Pt}{c}$$

$$\text{Total final momentum} = 0$$

$$\text{Total momentum transferred} = \frac{Pt}{c}$$

$$= \frac{20 \times 10^{-3} \times 300 \times 10^{-9}}{3 \times 10^8}$$

$$= 2 \times 10^{-17} \text{ Ns}$$

12. The velocity of an electron in the seventh orbit of hydrogen-like atom is $3.6 \times 10^6 \text{ m/s}$. Find the velocity of the electron in the 3rd orbit.

- (1) $4.2 \times 10^6 \text{ m/s}$ (2) $8.4 \times 10^6 \text{ m/s}$
(3) $2.1 \times 10^6 \text{ m/s}$ (4) $3.6 \times 10^6 \text{ m/s}$

Answer (2)

Sol. For hydrogen like atom,

$$v \propto \frac{1}{n}$$

$$\left(\frac{v_1}{v_2}\right) = \left(\frac{n_2}{n_1}\right)$$

$$\Rightarrow \frac{3.6 \times 10^6}{v_2} = \frac{3}{7}$$

$$\Rightarrow v_2 = \frac{7}{3} \times 3.6 \times 10^6$$

$$= 8.4 \times 10^6 \text{ m/s}$$

13. Electric field in a region is given by $\vec{E} = \frac{a}{x^2} \hat{i} + \frac{b}{y^3} \hat{j}$,

where x & y are co-ordinates. Find SI units of a & b .

- (1) $a - \text{Nm}^2\text{C}^{-1}$ (2) $a - \text{Nm}^3\text{C}^{-1}$
 $b - \text{Nm}^3\text{C}^{-1}$ $b - \text{Nm}^2\text{C}^{-1}$
 (3) $a - \text{NmC}^{-1}$ (4) $a - \text{Nm}^2\text{C}^{-1}$
 $b - \text{Nm}^2\text{C}^{-1}$ $b - \text{Nm}^2\text{C}^{-1}$

Answer (1)

Sol. $E - \text{NC}^{-1}$

$$x^2 - \text{m}^2$$

$$y^3 - \text{m}^3$$

$$\Rightarrow a - \text{Nm}^2\text{C}^{-1}$$

$$\& \quad b - \text{Nm}^3\text{C}^{-1}$$

14. Coil A of radius 10 cm has N_A number of turns and I_A current is flowing through it. Coil B of radius 20 cm has N_B number of turns and I_B current is flowing through it. If magnetic dipole moment of both the coils is same then

- (1) $I_A N_A = 4 I_B N_B$ (2) $I_A N_A = \frac{1}{4} I_B N_B$
 (3) $I_A N_A = 2 I_B N_B$ (4) $I_A N_A = \frac{1}{2} I_B N_B$

Answer (1)

Sol. Magnetic dipole moment $\mu = NIA = NI\pi R^2$

$$\text{So } \frac{\mu_A}{\mu_B} = \frac{N_A I_A R_A^2}{N_B I_B R_B^2} = 1$$

$$\frac{N_A I_A (10^2)}{N_B I_B (20^2)} = 1$$

$$N_A I_A = 4 N_B I_B$$

15. An ideal gas undergoes a thermodynamic process following the relation $PT^2 = \text{constant}$. Assuming symbols have their usual meaning then volume expansion coefficient of the gas is equal to

- (1) $\frac{2}{T}$ (2) $\frac{3}{T}$
 (3) $\frac{1}{2T}$ (4) $\frac{1}{T}$

Answer (2)

Sol. Volume expansion coefficient $= \frac{dV}{VdT}$

For $PT^2 = \text{constant}$




$$\text{Or } \frac{T^3}{V} = \text{constant}$$

$$\text{Or } \frac{dV}{dT} = (C) 3T^2$$

$$\text{Or } \frac{dV}{VdT} = \frac{3T^2}{T^3}$$

$$\frac{dV}{VdT} = \frac{3}{T}$$

16. Consider a combination of gates as shown :



Answer (1)

Sol. $y = (A'B') = A + B$

\Rightarrow OR gate


\Rightarrow Option 1

17. For the given YDSE setup. Find the number of fringes by which the central maxima gets shifted from point O.

(Given $d = 1 \text{ mm}$)

$$D = 1 \text{ m}$$


$$\lambda = 5000 \text{ \AA}$$



- (1) 10 (2) 15
 (3) 8 (4) 12

Answer (1)

Sol.



at central position, path difference, is,

$$(\mu - 1)t_1 - (\mu - 1)t_2$$

$$\Delta x = (\mu - 1)(t_1 - t_2)$$

$$\Delta x = \left(\frac{3}{2} - 1\right)(5.11 - 5.10) \text{ mm}$$

$$= \frac{1}{2} \times (0.01) \text{ mm}$$

$$= 0.005 \text{ mm}$$

$$= 5 \times 10^{-6} \text{ m}$$

$$\text{No. of fringes shifted} = \frac{\Delta x}{\lambda} = \frac{5 \times 10^{-6} \text{ m}}{5 \times 10^{-7} \text{ m}}$$

$$= 10$$

18.


19.

20.

SECTION - B

Numerical Value Type Questions: This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g. 06.25, 07.00, -00.33, -00.30, 30.27, -27.30) using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.

21. In a part of a circuit shown:



Find $V_A - V_B$ in volts. It is given that current is decreasing at a rate of 1 ampere/s.

Answer (18)

Sol. $V_A - iR - L \frac{di}{dt} - 12 = V_B$

$$\Rightarrow V_A - V_B = +18 \text{ volts}$$

22. A particle undergoing SHM follows the position-time equation given as $x = A \sin\left(\omega t + \frac{\pi}{3}\right)$. If the SHM motion has a time period of T , then velocity will be maximum at time $t = \frac{T}{\beta}$ for first time after $t = 0$. Value of β is equal to

Answer (03.00)

Sol. $x = A \sin\left(\omega t + \frac{\pi}{3}\right)$

$$\Rightarrow v = A\omega \cos\left(\omega t + \frac{\pi}{3}\right)$$

For maximum value of v

$$\cos\left(\omega t + \frac{\pi}{3}\right) = \pm 1$$


$$\Rightarrow \omega t + \frac{\pi}{3} = \pi \text{ (for nearest value of } t)$$

$$\omega t = \frac{2\pi}{3}$$


$$t = \frac{T}{3}$$

So $\beta = 3$

23. A block of mass 1 g is equilibrium with the help of a current carrying square loop which is partially lying in constant magnetic field (B) as shown. Resistance of the loop is 10Ω . Find the voltage (V) (in volts) of the battery in the loop.



Answer (10.00)



$$i l B = mg$$

$$i = \left(\frac{mg}{lB} \right) = \frac{(1 \times 10^{-3} \text{ kg}) \times (10 \text{ m/s}^2)}{(0.1 \text{ m}) \times (0.1 \text{ T})}$$

$$= 1 \times 10^{-3} \times 10^3$$

$$i = 1 \text{ A}$$

As resistance of loop = 10Ω

$$i = \frac{V}{R} = 1 \text{ A}$$

$$V = (1 \times 10) \text{ V} = 10 \text{ V}$$

24. Initial volume of 1 mole of a monoatomic gas is 2 litres. It is expanded isothermally to a volume of 6 litres. Change in internal energy is xR . Find x .

Answer (00)

Sol. $\Delta U = nC_V \Delta T$

$$= nC_V(0) \quad (\because \text{isothermal})$$

$$\Rightarrow \Delta U = 0$$

25. An object is placed at a distance of 40 cm from the pole of a converging mirror. The image is formed at a distance of 120 cm from the mirror on the same side. If the focal length is measured with a scale where each 1 cm has 20 equal divisions. If the fractional error in the measurement of focal length

is $\frac{1}{10k}$ Find k .

Answer (60.00)

Sol. $u = -40 \text{ cm}$

$$v = -120 \text{ cm}$$

$$\frac{1}{v} + \frac{1}{u} = \frac{1}{f}$$

$$\Rightarrow -\frac{1}{120} - \frac{1}{40} = \frac{1}{f}$$

$$\frac{1}{f} = \left(\frac{-1-3}{120} \right) = -\frac{4}{120}$$


$$f = -30 \text{ cm}$$

$$\text{Least count of scale} = \left(\frac{1}{20} \right) \text{ cm}$$

$$\text{Fractional error} = \left(\frac{1}{\frac{20}{30}} \right) = \left(\frac{1}{600} \right)$$


$$\text{as } \frac{1}{10k} = \frac{1}{600}$$

$$k = 60$$



In two circuit shown above the value of current I_1 (in amperes) is equal to $-\frac{y}{5}$ A. Value of y is equal to

Answer (11.00)



Using Kirchoff's law.

$$I_1 + I_3 - I_2 = -2 \quad \dots(i)$$

$$I_3 + 2I_2 = 5 \quad \dots(ii)$$

$$2I_2 - (I_3 - I_2) - (I_1 + I_3 - I_2) = 5 \quad \dots(iii)$$

$$\Rightarrow I_1 = -\frac{11}{5} \text{ A}$$

$$\Rightarrow y = 11$$

27. ??

28. ??

29. ??

30. ??