

- Draw the graph of  $f(x) = x^3 - 3x$ .
  - From the graph in (a), or otherwise, find the set of real numbers  $q$  such that the equation  $x^3 - 3x + q = 0$  has three distinct real roots.
- Let  $\mathbf{A}$  and  $\mathbf{B}$  be  $n \times n$  real matrices. Let  $\mathbf{I}_n$  denote the identity matrix of order  $n$ . Show that the matrix  $\begin{bmatrix} \mathbf{A} & \mathbf{I}_n \\ \mathbf{I}_n & \mathbf{B} \end{bmatrix}$  has rank  $n$  if and only if  $\mathbf{A}$  is nonsingular and  $\mathbf{B} = \mathbf{A}^{-1}$ .
- Let  $x_1, x_2, \dots, x_{10}$  be non-negative real numbers such that  $x_1^2 + x_2^2 + \dots + x_{10}^2 = 10$ . Find the maximum value of the product  $x_1^3 x_2^3 \dots x_{10}^3$ .

- The joint probability density function of the bivariate random vector  $(X, Y)$  is

$$f(x, y) = \frac{1}{2\pi} e^{-\sqrt{x^2 + y^2}}, \quad x, y \in \mathbb{R}.$$

Calculate  $P(X < 2Y)$ .

- Consider a random variable  $X$  having probability density function

$$f(x) = \begin{cases} \frac{k(p)}{x^p} & \text{for } x \geq 1, \\ 0 & \text{otherwise,} \end{cases}$$

where  $p > 0$  and  $k(p)$  is a suitable positive constant. Find the set of possible values of  $p$  for which  $\text{Var}(X)$  exists but the fourth moment of  $X$  does not exist.

- A box contains  $N$  balls numbered  $1, \dots, N$ , where  $N$  is unknown. From this box,  $n$  balls are drawn using simple random sampling with replacement and their numbers recorded. Let  $X_i$  denote the number recorded at the  $i^{\text{th}}$  draw,  $i = 1, \dots, n$ .
  - Find a statistic which is complete and sufficient for  $N$ .
  - Find the maximum likelihood estimator of  $N$ .
- Let  $X_1, X_2, \dots, X_n$  ( $n > 2$ ) be independent and identically distributed random variables having the uniform distribution over  $(\theta_1 - \theta_2, \theta_1 + \theta_2)$ , where  $\theta_1 \in \mathbb{R}$  and  $\theta_2 > 0$ . Find the uniformly minimum variance unbiased estimator of  $\frac{\theta_1}{\theta_2}$ .

8. Let  $X_1, X_2, \dots, X_n$  be independent and identically distributed random variables with probability density function given by

$$f(x|\theta) = \begin{cases} 3\theta^{-3}x^2 \exp(-(x/\theta)^3) & \text{if } x > 0, \\ 0 & \text{otherwise,} \end{cases}$$

where  $\theta > 0$  is unknown. We wish to test the hypothesis  $H_0 : \theta \leq 1$  against  $H_1 : \theta > 1$ . Let  $0 < \alpha < 1$ . Find a uniformly most powerful test of size  $\alpha$  for testing  $H_0$  against  $H_1$ .

9. Consider the linear model

$$\begin{aligned} E(Y_1) &= \theta_1 + \theta_2 + \theta_3, \\ E(Y_2) &= \theta_1 + \theta_2 - \theta_3, \\ E(Y_3) &= \theta_1 + \theta_2 - 2\theta_3, \end{aligned}$$

where  $Y_1, Y_2, Y_3$  are uncorrelated, each with unknown variance  $\sigma^2$  and  $\theta_1, \theta_2, \theta_3$  are three unknown parameters.

- (a) Is  $\theta_1$  estimable? Give reasons.  
(b) Is  $\theta_1 + \theta_2 - 4\theta_3$  estimable? Give reasons.
10. Let  $X_i, i \geq 1$ , be independent and identically distributed random variables having the uniform distribution over  $(0, 1)$ . Let  $X$  be defined as  $X = \sum_{i=1}^N X_i$ , where  $N$  is an unknown integer.
- (a) Find an unbiased estimator  $T(X)$  of  $N$ .  
(b) Decide, with adequate reasons, if  $\frac{T(X)}{N}$  converges to 1 almost surely, as  $N$  goes to infinity.