

QUESTION PAPER CODE 65/2/RU

EXPECTED ANSWERS/VALUE POINTS

SECTION - A

Marks

1. $|2\hat{a} + \hat{b} + \hat{c}|^2 = (2\hat{a})^2 + (\hat{b})^2 + (\hat{c})^2 + 2(2\hat{a} \cdot \hat{b} + \hat{b} \cdot \hat{c} + \hat{c} \cdot 2\hat{a})$ $\frac{1}{2}$ m

$\therefore |2\hat{a} + \hat{b} + \hat{c}| = \sqrt{6}$ $\frac{1}{2}$ m

2. $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{vmatrix} = -\hat{i} + \hat{j}$ $\frac{1}{2}$ m

unit vector is $-\frac{\hat{i}}{\sqrt{2}} + \frac{\hat{j}}{\sqrt{2}}$ $\frac{1}{2}$ m

3. $\frac{x - 3/5}{1/5} = \frac{y + 7/15}{1/15} = \frac{z - 3/10}{-1/10}$ $\frac{1}{2}$ m

Direction cosines are $\frac{6}{7}, \frac{2}{7}, \frac{-3}{7}$ or $\frac{-6}{7}, \frac{-2}{7}, \frac{3}{7}$ $\frac{1}{2}$ m

4. $\Delta = \begin{vmatrix} x+y+z & x+y+z & x+y+z \\ z & x & y \\ -3 & -3 & -3 \end{vmatrix}$ $\frac{1}{2}$ m

$$= 0$$
 $\frac{1}{2}$ m

5. order 2, degree 1 $\frac{1}{2}$ m
sum = 3 $\frac{1}{2}$ m

6. $\frac{dx}{dy} + \frac{2y}{1+y^2} \cdot x = \cot y$ $\frac{1}{2}$ m

Integrating factor = $e^{\log(1+y^2)}$ or $(1+y^2)$ $\frac{1}{2}$ m



SECTION - B

7. $y = e^{m \sin^{-1} x}$, differentiate w.r.t. "x", we get $\frac{dy}{dx} = \frac{m e^{m \sin^{-1} x}}{\sqrt{1-x^2}}$ $1\frac{1}{2}$ m

$$\Rightarrow \sqrt{1-x^2} \frac{dy}{dx} = my, \text{ Differentiate again w.r.t. "x"} \quad \text{[Ans]}$$

$$\Rightarrow \sqrt{1-x^2} \frac{d^2y}{dx^2} - \frac{x}{\sqrt{1-x^2}} \frac{dy}{dx} = m \frac{dy}{dx} \quad \text{[Ans]}$$

$$\Rightarrow (1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = m \left(\sqrt{1-x^2} \frac{dy}{dx} \right) = m(my) \quad \text{[Ans]}$$

$$\Rightarrow (1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - m^2y = 0 \quad \text{[Ans]}$$

8. $f(x) = \sqrt{x^2 + 1}$, $g(x) = \frac{x+1}{x^2 + 1}$, $h(x) = 2x - 3$

Differentiating w.r.t. "x", we get

$$f'(x) = \frac{x}{\sqrt{x^2 + 1}}, \quad g'(x) = \frac{1-2x-x^2}{(x^2+1)^2}, \quad h'(x) = 2 \quad \text{[Ans]}$$

$$\therefore f'(h'(g'(x))) = \frac{2}{\sqrt{5}} \quad \text{[Ans]}$$

9. $\int (3-2x) \sqrt{2+x-x^2} dx = 2 \int \sqrt{\left(\frac{3}{2}\right)^2 - \left(x - \frac{1}{2}\right)^2} dx + \int (1-2x) \sqrt{2+x-x^2} dx \quad \text{[Ans]}$

$$= 2 \cdot \left\{ \frac{x-\frac{1}{2}}{2} \sqrt{2+x-x^2} + \frac{9}{8} \sin^{-1} \left(\frac{x-\frac{1}{2}}{\frac{3}{2}} \right) \right\} + \frac{2}{3} (2+x-x^2)^{\frac{3}{2}} + c \quad \text{[Ans]}$$

$$\text{or } \left(\frac{2x-1}{2} \sqrt{2+x-x^2} + \frac{9}{4} \sin^{-1} \left(\frac{2x-1}{3} \right) + \frac{2}{3} (2+x-x^2)^{\frac{3}{2}} + c \right)$$



OR

$$\int \frac{x^2 + x + 1}{(x^2 + 1)(x + 2)} dx = \frac{1}{5} \int \frac{2x + 1}{x^2 + 1} dx + \frac{3}{5} \int \frac{1}{x + 2} dx \quad 2 \text{ m}$$

$$= \frac{1}{5} \int \frac{2x}{x^2 + 1} dx + \frac{1}{5} \int \frac{1}{x^2 + 1} dx + \frac{3}{5} \int \frac{1}{x + 2} dx \quad \frac{1}{2} \text{ m}$$

$$= \frac{1}{5} \log |x^2 + 1| + \frac{1}{5} \tan^{-1} x + \frac{3}{5} \log |x + 2| + c \quad 1\frac{1}{2} \text{ m}$$

10.
$$\begin{pmatrix} 400 & 300 & 100 \\ 300 & 250 & 75 \\ 500 & 400 & 150 \end{pmatrix} \begin{pmatrix} 50 \\ 20 \\ 40 \end{pmatrix} = \begin{pmatrix} 30000 \\ 23000 \\ 39000 \end{pmatrix} \quad 2 \text{ m}$$

cost incurred respectively for three villages is Rs. 30,000, Rs. 23,000, Rs. 39,000 1 m

One value : Women welfare or Any other relevant value 1 m

11.
$$\tan^{-1} \left(\frac{x+1+x-1}{1-(x+1)(x-1)} \right) = \tan^{-1} \left(\frac{8}{31} \right) \quad 2 \text{ m}$$

$$\Rightarrow \frac{2x}{2-x^2} = \frac{8}{31} \quad \therefore 4x^2 + 31x - 8 = 0 \quad 1 \text{ m}$$

$$\therefore x = \frac{1}{4}, -8 \text{ (Rejected)} \quad 1 \text{ m}$$

OR

$$\text{L.H.S.} = \tan^{-1} \left(\frac{x-y}{1+xy} \right) + \tan^{-1} \left(\frac{y-z}{1+yz} \right) + \tan^{-1} \left(\frac{z-x}{1+zx} \right) \quad 2 \text{ m}$$

$$= \tan^{-1} x - \tan^{-1} y + \tan^{-1} y - \tan^{-1} z + \tan^{-1} z - \tan^{-1} x \quad \left. \right\} \quad 2 \text{ m}$$

$$= 0 = \text{RHS}$$



$$12. \begin{vmatrix} a^2 & bc & ac + c^2 \\ a^2 + ab & b^2 & ac \\ ab & b^2 + bc & c^2 \end{vmatrix} = abc \begin{vmatrix} a & c & a+c \\ a+b & b & a \\ b & b+c & c \end{vmatrix}$$

Taking a, b & c common from C_1 , C_2 and C_3 1 m

$$= 2abc \begin{vmatrix} a+c & c & a+c \\ a+b & b & a \\ b+c & b+c & c \end{vmatrix}$$

$C_1 \rightarrow C_1 + C_2 + C_3$ and taking 2 common from C_1 1 m

$$= 2abc \begin{vmatrix} a+c & c & 0 \\ a+b & b & -b \\ b+c & b+c & -b \end{vmatrix} C_3 \rightarrow C_3 - C_1 1 m$$

$$= 2abc \begin{vmatrix} a+c & c & 0 \\ a-c & -c & 0 \\ b+c & b+c & -b \end{vmatrix} R_2 \rightarrow R_2 - R_3 \frac{1}{2} m$$

Expand by C_3 , $= 2abc (-b)(-ac - c^2 - ac + c^2) = 4a^2 b^2 c^2 \frac{1}{2} m$

$$13. \text{ Adj } A = \begin{pmatrix} -3 & 6 & 6 \\ -6 & 3 & -6 \\ -6 & -6 & 3 \end{pmatrix}; |A| = 27 \quad 2+1 m$$

$$\text{A. Adj } A = \begin{pmatrix} -1 & -2 & -2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{pmatrix} \begin{pmatrix} -3 & 6 & 6 \\ -6 & 3 & -6 \\ -6 & -6 & 3 \end{pmatrix} = 27 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = |A| I_3 \quad 1 m$$

$$14. f(x) = |x-1| + |x+1|$$

$$L f'(-1) = \lim_{x \rightarrow (-1)^-} \frac{\{-(x-1)-(x+1)\}-2}{x-(-1)} = \lim_{x \rightarrow (-1)^-} \frac{-2(x+1)}{x+1} = -2 \quad 1 m$$



$$R f'(-1) = \lim_{x \rightarrow (-1)^+} \frac{-(x-1)+(x+1)-2}{x-(-1)} = \lim_{x \rightarrow (-1)^+} \frac{0}{x+1} = 0 \quad 1 \text{ m}$$

$-2 \neq 0 \therefore f(x)$ is not differentiable at $x = -1$

$$L f'(1) = \lim_{x \rightarrow 1^-} \frac{-(x-1)+(x+1)-2}{x-1} = \lim_{x \rightarrow 1^-} \frac{0}{x-1} = 0 \quad 1 \text{ m}$$

$$R f'(1) = \lim_{x \rightarrow 1^+} \frac{x-1+x+1-2}{x-1} = \lim_{x \rightarrow 1^+} \frac{2(x-1)}{x-1} = 2 \quad 1 \text{ m}$$

$0 \neq 2 \therefore f(x)$ is not differentiable at $x = 1$

15. let the equation of line passing through $(1, 2, -4)$ be

$$\vec{r} = \hat{i} + 2\hat{j} - 4\hat{k} + \lambda(a\hat{i} + b\hat{j} + c\hat{k}) \quad 1 \text{ m}$$

Since the line is perpendicular to the two given lines \therefore

$$\therefore 3a - 16b + 7c = 0$$

$$3a + 8b - 5c = 0$$

$$\text{Solving we get, } \frac{a}{24} = \frac{b}{36} = \frac{c}{72} \text{ or } \frac{a}{2} = \frac{b}{3} = \frac{c}{6} \quad 1 \text{ m}$$

$$\therefore \text{Equation of line is : } \vec{r} = \hat{i} + 2\hat{j} - 4\hat{k} + \lambda(2\hat{i} + 3\hat{j} + 6\hat{k}) \quad \frac{1}{2} \text{ m}$$

OR

$$\text{Equation of plane is : } \begin{vmatrix} x+1 & y-2 & z \\ 2+1 & 2-2 & -1 \\ 1 & 1 & -1 \end{vmatrix} = 0 \quad 3 \text{ m}$$

$$\text{Solving we get, } x + 2y + 3z - 3 = 0 \quad 1 \text{ m}$$

16. Let $x = \text{No. of spades in three cards drawn}$

$$\begin{array}{ccccccc} x & : & 0 & 1 & 2 & 3 & 1 \text{ m} \\ P(x) & : & \frac{3}{4^3} & \frac{3}{4} \left(\frac{1}{4}\right) \left(\frac{3}{4}\right)^2 & \frac{3}{4} \left(\frac{1}{4}\right)^2 \frac{3}{4} & \frac{3}{4} \left(\frac{1}{4}\right)^3 \left(\frac{3}{4}\right)^0 & 2 \text{ m} \\ & & = \frac{27}{64} & = \frac{27}{64} & = \frac{9}{64} & = \frac{1}{64} & \\ x \cdot P(x) & : & 0 & \frac{27}{64} & \frac{18}{64} & \frac{3}{64} & \frac{1}{2} \text{ m} \end{array}$$



$$\text{Mean} = \sum x \cdot P(x) = \frac{48}{64} = \frac{3}{4}$$
½ m

OR

let p = probability of success ; q = Probability of failure

$$\text{then, } 9 P(x=4) = P(x=2)$$

$$\Rightarrow 9 \cdot {}^6C_4 p^4 \cdot q^2 = {}^6C_2 \cdot p^2 \cdot q^4$$
2 m

$$\Rightarrow 9p^2 = q^2 \therefore q = 3p$$
1 m

$$\text{Also, } p + q = 1 \Rightarrow p + 3p = 1 \therefore p = \frac{1}{4}$$
1 m

17. 

$$\int_0^{\pi/4} \frac{1}{\cos^3 x \sqrt{2 \sin 2x}} dx = \int_0^{\pi/4} \frac{1}{\cos^4 x 2 \sqrt{\tan x}} dx$$
1 m

$$= \int_0^{\pi/4} \frac{(1 + \tan^2 x)}{2 \sqrt{\tan x}} \sec^2 x dx$$
1 m

$$= \frac{1}{2} \int_0^1 \frac{1+t^2}{\sqrt{t}} dt \quad \text{Taking, } \tan x = t;$$
1 m

$$= \frac{1}{2} \left[2\sqrt{t} + \frac{2}{5} t^{5/2} \right]_0^1$$
½ m

$$= \frac{1}{2} \left[2 + \frac{2}{5} \right] = \frac{6}{5}$$
½ m

18. $\int \log x \cdot \frac{1}{(x+1)^2} dx = \log x \cdot \frac{-1}{x+1} + \int \frac{1}{x} \cdot \frac{1}{x+1} dx$

2 m

$$= \frac{-\log x}{x+1} + \int \frac{1}{x} dx - \int \frac{1}{x+1} dx$$
1 m



$$= \frac{-\log x}{x+1} + \log x - \log(x+1) + c \quad 1 \text{ m}$$


$$\text{or } \frac{-\log x}{x+1} + \log \left(\frac{x}{x+1} \right) + c$$

19. $\vec{a} - \vec{b} = -\hat{i} + \hat{j} + \hat{k}; \vec{c} - \vec{b} = \hat{i} - 5\hat{j} - 5\hat{k} \quad 1\frac{1}{2} \text{ m}$

$$(\vec{a} - \vec{b}) \times (\vec{c} - \vec{b}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 1 & 1 \\ 1 & -5 & -5 \end{vmatrix} = -4\hat{j} + 4\hat{k} \quad 1\frac{1}{2} \text{ m}$$

\therefore Unit vector perpendicular to both of the vectors $= -\frac{\hat{j}}{\sqrt{2}} + \frac{\hat{k}}{\sqrt{2}}$ 1 m

SECTION - C

20.  $\frac{dy}{dx} = \frac{y^2}{xy - x^2} \Rightarrow \frac{dy}{dx} = \frac{\left(\frac{y}{x}\right)^2}{\frac{y}{x} - 1}$, Hence the differential equation is homogeneous 1 m

Put $y = vx$ and $\frac{dy}{dx} = v + x \frac{dv}{dx}$, we get $v + x \frac{dv}{dx} = \frac{v^2}{v-1}$ $1+1 \text{ m}$

$$\therefore x \frac{dv}{dx} = \frac{v^2}{v-1} - v = \frac{v}{v-1} \quad 1 \text{ m}$$

$$\int \frac{v-1}{v} dv = \int \frac{1}{x} dx \Rightarrow v - \log v = \log x + c \quad 1 \text{ m}$$

$$\therefore \frac{y}{x} - \log \frac{y}{x} = \log x + c \quad \left(\text{or, } \frac{y}{x} = \log y + c \right) \quad 1 \text{ m}$$

OR



Given differential equation can be written as $\frac{dx}{dy} + \frac{1}{1+y^2} x = \frac{\tan^{-1}y}{1+y^2}$ 1 m

Integrating factor = $e^{\tan^{-1}y}$ and solution is : $x e^{\tan^{-1}y} = \int \frac{\tan^{-1}y \cdot e^{\tan^{-1}y}}{1+y^2} dy$ 1+1½ m

$x e^{\tan^{-1}y} = \int te^t dt = te^t - e^t + c = e^{\tan^{-1}y} (\tan^{-1}y - 1) + c$ (where $\tan^{-1}y = t$) 1½ m

$x = 1, y = 0 \Rightarrow c = 2 \therefore x \cdot e^{\tan^{-1}y} = e^{\tan^{-1}y} (\tan^{-1}y - 1) + 2$ 1 m

$$\text{or } x = \tan^{-1}y - 1 + 2 e^{-\tan^{-1}y}$$

21. Equation of line through A and B is $\frac{x-3}{-1} = \frac{y+4}{1} = \frac{z+5}{6} = \lambda$ (say) 2 m

General point on the line is $(-\lambda + 3, \lambda - 4, 6\lambda - 5)$ 1 m

If this is the point of intersection with plane $2x + y + z = 7$

then, $2(-\lambda + 3) + \lambda - 4 + 6\lambda - 5 = 7 \Rightarrow \lambda = 2$ 1 m

\therefore Point of intersection is $(1, -2, 7)$ 1 m

Required distance = $\sqrt{(3-1)^2 + (4+2)^2 + (4-7)^2} = 7$ 1 m

22. $f: R_+ \rightarrow [-9, \infty); f(x) = 5x^2 + 6x - 9; f^{-1}(y) = \frac{\sqrt{54+5y}-3}{5}$

$$f \circ f^{-1}(y) = 5 \left\{ \frac{\sqrt{54+5y}-3}{5} \right\}^2 + 6 \left\{ \frac{\sqrt{54+5y}-3}{5} \right\} - 9 = y \quad 3 \text{ m}$$

$$f^{-1} \circ f(x) = \frac{\sqrt{54+5(5x^2+6x-9)}-3}{5} = x \quad 2\frac{1}{2} \text{ m}$$

Hence 'f' is invertible with $f^{-1}(y) = \frac{\sqrt{54+5y}-3}{5}$ ½ m

OR

20



(i) commutative : let $x, y \in R - \{-1\}$ then

$$x * y = x + y + xy = y + x + yx = y * x \therefore * \text{ is commutative}$$

1½ m

(ii) Associative : let $x, y, z \in R - \{-1\}$ then

$$x * (y * z) = x * (y + z + yz) = x + (y + z + yz) + x(y + z + yz)$$

$$= x + y + z + xy + yz + zx + xyz$$

1½ m

$$(x * y) * z = (x + y + xy) * z = (x + y + xy) + z + (x + y + xy) \cdot z$$

$$= x + y + z + xy + yz + zx + xyz$$

1 m

$$x * (y * z) = (x * y) * z \therefore * \text{ is Associative}$$

(iii) Identity Element : let $e \in R - \{-1\}$ such that $a * e = e * a = a \forall a \in R - \{-1\}$

$$\therefore a + e + ae = a \Rightarrow e = 0$$

½ m

(iv) Inverse : let $a * b = b * a = e = 0 ; a, b \in R - \{-1\}$

½ m

$$\Rightarrow a + b + ab = 0 \therefore b = \frac{-a}{1+a} \text{ or } a^{-1} = \frac{-a}{1+a}$$

½ m

23. Solving the two curves to get the points of intersection $(\pm 3\sqrt{p}, 8)$

1½ m

$$m_1 = \text{slope of tangent to first curve} = \frac{-2x}{9p}$$

1½ m

$$m_2 = \text{slope of tangent to second curve} = \frac{2x}{p}$$

1½ m

$$\text{curves cut at right angle iff } \frac{-2x}{9p} \times \frac{2x}{p} = -1$$

½ m

$$\Leftrightarrow 9p^2 = 4x^2 \text{ (Put } x = \pm 3\sqrt{p})$$

$$\Leftrightarrow 9p^2 = 4(9p)$$

$$\therefore p = 0 ; p = 4$$

1 m


24.

correct figure

1½ m

$$\text{ar (ABDOA)} = \frac{1}{4} \int_0^4 y^2 dy = \left[\frac{y^3}{12} \right]_0^4 = \frac{16}{3} \dots\dots(i) \quad 1\frac{1}{2} \text{ m}$$





$$\text{ar(OEBDO)} = \int_0^4 2\sqrt{x} \, dx - \int_0^4 \frac{x^2}{4} \, dx = \left[\frac{4}{3}x^{3/2} - \frac{x^3}{12} \right]_0^4$$

$$= \frac{32}{3} - \frac{16}{3} = \frac{16}{3} \quad \dots\dots\dots \text{(ii)} \quad 1\frac{1}{2} \text{ m}$$

$$\text{ar (OEBCO)} = \frac{1}{4} \int_0^4 x^2 dx = \left[\frac{x^3}{12} \right]_0^4 = \frac{16}{3} \dots \dots \dots \text{(iii)} \quad 1\frac{1}{2} \text{ m}$$

From (i), (ii) and (iii) we get $\text{ar}(\text{ABDOA}) = \text{ar}(\text{OEBDO}) = \text{ar}(\text{OEBCO})$

25. E_1 : Bolt is manufactured by machine A
 E_2 : Bolt is manufactured by machine B
 E_3 : Bolt is manufactured by machine C
 A : Bolt is defective

$$P(E_1) = \frac{30}{100}; P(E_2) = \frac{50}{100}; P(E_3) = \frac{20}{100};$$

$$P(A/E_1) = \frac{3}{100}; P(A/E_2) = \frac{4}{100}; P(A/E_3) = \frac{1}{100}$$

$$P(E_2/A) = \frac{\frac{30}{100} \times \frac{3}{100} + \frac{50}{100} \times \frac{4}{100} + \frac{20}{100} \times \frac{1}{100}}{\frac{50}{100} \times \frac{4}{100}} = \frac{200}{90 + 200 + 20} = \frac{20}{31}$$

$$P(\bar{E}_2/A) = 1 - P(E_2/A) = \frac{11}{31}$$


26. Let the two factories I and II be in operation for x and y

days respectively to produce the order with the minimum cost

then, the LPP is :

Minimise cost : $z = 12000x + 15000y$

Subject to :



$$50x + 40y \geq 6400 \quad \text{or} \quad 5x + 4y \geq 640$$

$$50x + 20y \geq 4000 \quad \text{or} \quad 5x + 2y \geq 400 \quad 2 \text{ m}$$

$$30x + 40y \geq 4800 \quad \text{or} \quad 3x + 4y \geq 480$$

$$x, y \geq 0$$

correct graph 2 m

Vertices are A (0, 200) ; B (32, 120)

C (80, 60) ; D (160, 0) $\frac{1}{2} \text{ m}$

$z(A) = \text{Rs. } 30,00,000; z(B) = \text{Rs. } 21,84,000;$

$z(C) = \text{Rs. } 18,60,000 \text{ (Min.)}; z(D) = \text{Rs. } 19,20,000;$

On plotting $z < 1860000$

or $12x + 15y < 1860$, we get no

point common to the feasible region

\therefore Factory I operates for 80 days $\frac{1}{2} \text{ m}$

Factory II operates for 60 days

