


Marking scheme – 2020




CHEMISTRY (043) / CLASS XII





56/C/1






Q. No	Expected Answer / Value Points	Marks
SECTION A		
1	Starch/ cellulose/ proteins / nucleic acids / natural rubber (or any other suitable example)	1
2	$\left[\text{CH}_2 - \overset{\text{Cl}}{\underset{ }{\text{C}}} = \text{CH} - \text{CH}_2 \right]_n$ <p style="text-align: center;">Neoprene</p>	1
3	Homopolymer	1
4	Teflon / PTFE	1
5	PHBV / Nylon-2-nylon-6 / any natural polymer (or any other suitable example)	1
6	Nucleoside	1
7	Smoke/dust (or any other suitable example)	1
8	Alitame	1
9	Molar conductivity increases.	1
10	Kraft Temperature	1
11	(D)	1
12	(C) or (D)	1
13	(A)	1
14	(D)	1
15	(D)	1
16	(i)	1
17	(i)	1



18	(iii)	1
19	(iii)	1
20	(iv)	1
SECTION B		
21	<p>a) At Anode: $2\text{H}_2\text{O}_{(l)} \longrightarrow 4\text{H}^+_{(aq)} + 4\text{e}^- + \text{O}_2$ At Cathode: $\text{Cu}^{2+}_{(aq)} + 2\text{e}^- \longrightarrow \text{Cu}_{(s)}$. / Copper is deposited at cathode and Oxygen gas is liberated at anode.</p> <p>b) At Anode: $2\text{H}_2\text{O}_{(l)} \longrightarrow 4\text{H}^+_{(aq)} + 4\text{e}^- + \text{O}_2$ At Cathode: $\text{Ag}^+_{(aq)} + \text{e}^- \longrightarrow \text{Ag}_{(s)}$. / Silver is deposited at cathode and oxygen gas is liberated at anode.</p> <p style="text-align: center;">OR</p> $\text{Fe}^{3+} + \text{e}^- \longrightarrow \text{Fe}^{2+}$, so 1 mol of Fe^{3+} requires 1 F 3 moles of Fe^{3+} require 3 F $Q = I \times t$ $t = 3 \times 96500 / 2$ $t = 144750 \text{ sec}$	<p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
22	<p>Vitamins are certain organic compounds, required in small amounts in our diet but their deficiency causes specific diseases / organic compounds required in the diet in small amounts to perform specific biological functions for normal maintenance of optimum growth and health of the organism. Vitamins are classified into two groups depending upon their solubility in water or fat. (i) Fat soluble vitamins (ii) Water soluble vitamins.</p> <p style="text-align: center;">OR</p> <p>Proteins are polymers of α-amino acids. (or any other correct answer) They are classified as Fibrous and Globular proteins on the basis of their shape.</p>	<p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>
23	<p>Antiseptics the chemicals which either kill or prevent the growth of microorganisms but are applied to the living tissues such as wounds, cuts, ulcers and diseased skin surfaces. Examples are furacine, soframincine, etc. Disinfectants are also the chemicals which either kill or prevent the growth of microorganisms but applied to inanimate objects such as floors, drainage system, instruments, etc. e.g. Concentrated acids, Phenol (above 1% conc.)</p> <p style="text-align: center;">(OR any other suitable point of difference and example)</p>	<p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>
24.	 <p style="text-align: center;">Cis Trans</p>	1+1
25.	<p>It is the rate of reaction when concentration of each reactant is taken as unity. / It is the proportionality constant in the rate law expression or in differential rate equation or in the rate of reaction. $K = 0.693 / t_{1/2}$</p>	<p>1</p> <p>1</p>
26.	<p>a) Electrolytic refining -The more basic metal remains in the solution and the less basic ones go to the anode mud. / Anode is impure metal and pure metal strip is cathode while aqueous solution of the metal salt acts as the electrolyte. b) Zone refining - Impurities are more soluble in the melt than in the solid state of the metal.</p>	<p>1</p> <p>1</p>
27.	<p>$\text{Mn}^{3+} = 4$ unpaired electrons $\text{Cr}^{3+} = 3$ unpaired electrons Cr^{3+} is more stable due to half filled t_{2g}^3 configuration</p>	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>
SECTION C		
28	$\Lambda_m = \frac{k}{c} \times 1000$ $= \frac{8 \times 10^{-5}}{0.002} \times 1000$ $= 40 \text{ Scm}^2\text{mol}^{-1}$	<p>$\frac{1}{2}$</p> <p>1</p>

	$\alpha = \frac{\Delta_m^c}{\Delta_m^o}$ $= 40/390.5 = 0.102$	$\frac{1}{2}$ 1
29	$K = \frac{2.303}{t} \log \frac{[R_0]}{[R]}$ $4.9 \times 10^{-3} = \frac{2.303}{t} \log \frac{4}{3}$ $t = \frac{2.303}{4.9 \times 10^{-3}} \log (0.6020 - 0.4771)$ $= 58.7 \text{ sec}$ <p style="text-align: center;">OR</p> a) Molecularity = 2 or bimolecular. b) Order = 1 or pseudo first order. c) Rate = $\frac{-\Delta[C_{12}H_{22}O_{11}]}{\Delta t} = + \frac{\Delta[C_6H_{12}O_6]}{\Delta t} = + \frac{\Delta[C_6H_{12}O_6]}{\Delta t}$	$\frac{1}{2}$ 1 1 $\frac{1}{2}$ 1 1 1
30	a) Electrophoresis / coagulation / neutralisation of dispersed phase particles / movement of colloidal particles towards oppositely charged electrode / precipitation. b) Demulsification / coagulation / separation of components of emulsion / breaking down of emulsion. c) Coagulation / mutual coagulation / precipitation.	1 1 1
31	$4 \text{ FeCr}_2\text{O}_4 + 8 \text{ Na}_2\text{CO}_3 + 7 \text{ O}_2 \rightarrow 8 \text{ Na}_2\text{CrO}_4 + 2 \text{ Fe}_2\text{O}_3 + 8 \text{ CO}_2$ $2\text{Na}_2\text{CrO}_4 + 2 \text{ H}^+ \rightarrow \text{Na}_2\text{Cr}_2\text{O}_7 + 2 \text{ Na}^+ + \text{H}_2\text{O}$ $\text{Na}_2\text{Cr}_2\text{O}_7 + 2 \text{ KCl} \rightarrow \text{K}_2\text{Cr}_2\text{O}_7 + 2 \text{ NaCl}$ <p style="text-align: right;">(Ignore balancing)</p>	1 1 1
32	a) It reacts with moisture readily / It is highly reactive with any source of proton / it forms hydrocarbons or alkanes / Grignard's reagent gets hydrolysed easily / $\text{RMgX} + \text{H}_2\text{O} \rightarrow \text{RH} + \text{Mg(OH)X}$. b) They can't form hydrogen bonds with water / water can't compensate for the intermolecular forces of alkyl halides / interactions between water molecules and alkyl halides molecules are not strong / less energy is released when alkyl halide and water are mixed. c) Chloroform gets oxidised in air and sunlight / poisonous gas or phosgene is formed / $\text{CHCl}_3 + \frac{1}{2} \text{ O}_2 \rightarrow \text{COCl}_2 + \text{HCl}$.	1 1 1
	(or any other correct reason)	
33	a)  b) $\text{CH}_3\text{CH}_2\text{OH}$ c)  <p style="text-align: center;">OR</p> a) i) $\text{H}_3\text{C}-\text{CH}=\text{CH}_2 \xrightarrow{\text{H}_2\text{O}, \text{H}^+} \text{H}_3\text{C}-\underset{\text{OH}}{\text{CH}}-\text{CH}_3$ ii)  b) 4-Methylphenol < phenol < 4-Nitrophenol / 4-Methylphenol, phenol, 4-Nitrophenol	1 x 3 1 1 1
34	Hinsberg Test: Methyl amine or 1° amine reacts with Hinsberg's reagent or Benzene Sulphonyl Chloride to give a product which is soluble in alkali, Dimethyl amine or 2° amine reacts with Hinsberg's reagent or Benzene Sulphonyl Chloride to give	$\frac{1}{2}$ $\frac{1}{2}$

	<p>a product which is insoluble in alkali while Trimethylamine or 3° amine doesn't react with Hinsberg's reagent or Benzene Sulphonyl Chloride.</p> <p> $\text{H}_3\text{C}-\text{NH}_2 + \text{C}_6\text{H}_5\text{SO}_2\text{Cl} \rightarrow \text{C}_6\text{H}_5\text{SO}_2\text{NHCH}_3$ This compound is soluble in alkali </p> <p> $(\text{CH}_3)_2\text{NH} + \text{C}_6\text{H}_5\text{SO}_2\text{Cl} \rightarrow \text{C}_6\text{H}_5\text{SO}_2\text{N}(\text{CH}_3)_2$ This compound is insoluble in alkali </p> <p> $(\text{CH}_3)_3\text{N} + \text{C}_6\text{H}_5\text{SO}_2\text{Cl} \rightarrow \text{No reaction}$ </p> <p>(Or any other suitable chemical test to identify different degree methyl amines)</p>	<p>½</p> <p>½</p> <p>½</p>
SECTION D		
35	<p>a) i)  ii) </p> <p>b) i)  ii) </p> <p>c) Carboxylate ion is more stabilised than phenoxide ion / conjugate base of carboxylic acid is more stable than that of phenol / carboxylate ion has two equivalent resonating structures while the structures are non equivalent in phenoxide ion/ negative charge in carboxylate ion is delocalised over more electronegative two Oxygen atoms while in phenoxide ion negative charge is delocalised over one Oxygen atom and less electronegative Phenyl ring (or C-atoms). /Carboxylic acid reacts with NaHCO_3 to give brisk effervescence of CO_2 while phenol doesn't or reaction given by the student.</p>	<p>1+1</p> <p>1+1</p> <p>1</p>
35	<p>a) $\text{CH}_3\text{COOCH}_2\text{CH}_3 \xrightarrow{\text{H}_2\text{SO}_4 \text{ dil}} \text{CH}_3\text{COOH} + \text{CH}_3\text{CH}_2\text{OH}$</p> <p style="text-align: center;">OR</p> <p>$\text{CH}_3\text{CH}_2\text{OH} \xrightarrow{\text{oxidation}} \text{CH}_3\text{COOH}$</p> <p>$\text{CH}_3\text{CH}_2\text{OH} \xrightarrow{\text{dehydration}} \text{CH}_2=\text{CH}_2$</p> <p>A = Ethyl acetate ($\text{CH}_3\text{COOCH}_2\text{CH}_3$), B = Acetic or Ethanoic acid (CH_3COOH), C = Ethanol ($\text{CH}_3\text{CH}_2\text{OH}$)</p>	<p>½ x 3</p> <p>½ x 3</p>

	<p>b) i) $\text{CH}_3\text{CHO} \xrightarrow{\text{LiAlH}_4, \text{dry ether}} \text{CH}_3\text{CH}_2\text{OH}$</p> <p>ii) $\text{CH}_3\text{CHO} \xrightarrow{\text{Zn/Hg in HCl}_{\text{conc.}}} \text{CH}_3\text{CH}_3$ Or NH_2NH_2 with</p> <p>(or any other suitable reagent for the above reactions or any other correct method for conversion)</p>	1
	<p>ii) $\text{CH}_3\text{CHO} \xrightarrow{\text{Zn/Hg in HCl}_{\text{conc.}}} \text{CH}_3\text{CH}_3$ Or NH_2NH_2 with</p> <p>(or any other suitable reagent for the above reactions or any other correct method for conversion)</p>	1
36	<p>a) i) There are two -O-H bonds or groups / ii) Due to lower bond dissociation enthalpy of Te-H than H-O/ due to large size of Te / longer bond length of Te-H than H-O. iii) Due to highest electronegativity / due to absence of vacant d-orbitals / It can show only one oxidation state / it can't show higher positive oxidation state.</p> <p>b) i) $2\text{XeF}_6 + 2\text{H}_2\text{O} \longrightarrow \text{XeO}_2\text{F}_2 + 4\text{HF}$ (ignore balancing) ii) $\text{I}_2 + 3\text{Cl}_2 \longrightarrow 2\text{ICl}_3$ (ignore balancing)</p> <p>OR</p> <p>a) i) $\text{I}_2 < \text{F}_2 < \text{Br}_2 < \text{Cl}_2$ ii) $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$ iii) $\text{H}_2\text{O} > \text{H}_2\text{S} > \text{H}_2\text{Se} > \text{H}_2\text{Te}$</p>  <p>Sulphurous acid (H_2SO_3)</p>  	1 1 1 1 1
36	<p>a) i) $\text{I}_2 < \text{F}_2 < \text{Br}_2 < \text{Cl}_2$ ii) $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$ iii) $\text{H}_2\text{O} > \text{H}_2\text{S} > \text{H}_2\text{Se} > \text{H}_2\text{Te}$</p>  	1 1 1
37	<p>a) Henry's law : The law states that at a constant temperature, the solubility (mole fraction) of a gas in a liquid is directly proportional to the partial pressure of the gas present above the surface of liquid or solution. Applications: To avoid bends, in the condition of anoxia and to fill CO_2 in cold drink bottles (any of the two applications)</p> <p>b) $\frac{P_o - P}{P_o} = X_2$</p> $\frac{P_o - P}{P_o} = \frac{n_2}{n_1 + n_2} \approx \frac{n_2}{n_1}$ $\frac{760 - 745}{760} = \frac{w_2}{M_2} \times \frac{M_1}{w_1} = \frac{15}{760} = \frac{5}{M_2} \times \frac{18}{95}$ $M_2 = \frac{760 \times 5 \times 18}{95 \times 15} = 48 \text{ gmol}^{-1} \text{ or u}$	1 $\frac{1}{2} + \frac{1}{2}$ 1 1 1

37	OR		
	a)		
	Ideal Solution	Non-Ideal solution	½ x 4
	It obeys Raoult's law over the entire range of concentration.	It doesn't obey Raoult's law over the entire range of concentration.	
	$\Delta V_{\text{mixing}} = 0$ and $\Delta H_{\text{mixing}} = 0$	$\Delta V_{\text{mixing}} \neq 0$ and $\Delta H_{\text{mixing}} \neq 0$.	
	(Any other two points of difference between the two).		
	b) $\Delta T_f = iK_f m$, for NaCl, $i=2$, $2 = 2 \times K_f \frac{Wb \times 1000}{Mb \times Wa}$ $1 = 1.86 \times \frac{w \times 1000}{58.5 \times 100}$ $w = 3.147 \text{ g}$ (1/2 mark to be deducted for incorrect or no units)		½ ½ ½ 1
	(or by any other correct method)		



collegedunia
India's largest Student Review Platform