

**NARAYANA GRABS  
THE LION'S SHARE IN JEE-ADV.2022**

**5 RANKS in OPEN CATEGORY  
ONLY FROM NARAYANA  
IN TOP 10 AIR**



JEE MAIN (APRIL) 2023 (10-04-2023-AN)  
*Memory Based Question Paper*  
**MATHEMATICS**



## MATHEMATICS


### SECTION - A

**Multiple Choice Questions:** This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

**Choose the correct answer:**

1. Let a circle  $x^2 + y^2 = 16$  and line passing through  $(1, 2)$  cuts the curve at  $A$  and  $B$  then the locus of the mid-point of  $AB$  is
  - (1)  $x^2 + y^2 + x + y = 0$
  - (2)  $x^2 + y^2 - x + 2y = 0$
  - (3)  $x^2 + y^2 - x - 2y = 0$
  - (4)  $x^2 + y^2 + x + 2y = 0$

**Answer (3)**



Let  $P(x_1, y_1)$  be the mid-point of  $AB$

Then  $T = S_1$

$$x_1^2 + y_1^2 - 16 = xx_1 + yy_1 - 16$$

$$\Rightarrow xx_1 + yy_1 = x_1^2 + y_1^2 \quad \dots(i)$$

$\therefore$  (i) passes through  $(1, 2)$

$$\therefore x_1 + 2y_1 = x_1^2 + y_1^2$$

$\therefore$  required locus

$$x^2 + y^2 - x - 2y = 0$$

2. Consider  $f(x) = \sec^{-1}\left(\frac{2x}{5x+3}\right)$ , domain of  $f(x)$  is  $[\alpha, \beta] \cup [\gamma, \delta]$ , then the value of  $|3\alpha + 10\beta + 5\gamma + 21\delta|$  is
  - (1) 22
  - (2) 23
  - (3) 21
  - (4) 19

**Answer (3)**

$$\text{Sol. } \frac{2x}{5x+3} \geq 1 \text{ OR } \frac{2x}{5x+3} \leq -1$$

$$\Rightarrow \frac{2x}{5x+3} - 1 \geq 0$$

$$\Rightarrow \frac{-3x-3}{5x+3} \geq 0$$

$$\Rightarrow \frac{x+1}{5x+3} \leq 0 \Rightarrow x \in \left[-1, \frac{-3}{5}\right)$$

$$\frac{2x}{5x+3} + 1 \leq 0$$

$$\Rightarrow \frac{7x+3}{5x+3} \leq 0 \Rightarrow x \in \left(-\frac{3}{5}, \frac{-3}{7}\right]$$

$$\therefore x \in \left[-1, \frac{-3}{5}\right] \cup \left(\frac{-3}{5}, \frac{-3}{7}\right]$$

$$\alpha = -1, \beta = \frac{-3}{5}, \gamma = \frac{-3}{5}, \delta = \frac{-3}{7}$$

$$|3\alpha + 10\beta + 5\gamma + 21\delta|$$

$$= |-3 - 6 - 3 - 9| = 21$$

3. 8 persons has to travel from  $A$  to  $B$  in 3 allotted cars. If a car can carry maximum 3 persons. Then find the number of ways they can travel.
  - (1) 1880
  - (2) 1800
  - (3) 1680
  - (4) 1600

**Answer (3)**

$$\text{Sol. } \begin{matrix} C_1 & C_2 & C_3 \\ 3 & 3 & 2 \end{matrix} \rightarrow$$

$$\text{Total } \frac{8!}{3! 3! 2! 2!} \text{ groups}$$

So, they can travel in

$$\frac{8!}{3! 3! 2! 2!} \times 3! \text{ ways}$$

$$= 1680$$

4. If  $\frac{z+i}{4z+zi}$  is purely real  $\Delta z = x + iy$  ( $x, y \in \mathbb{R}$ ) then one of the possibility is

- (1)  $x \neq 0, y \neq -1$
- (2)  $x \neq 0, y = -1$
- (3)  $x = -1, y = 1$
- (4)  $x = 1, y \neq -1$

**Answer (2)**

$$\text{Sol. } \operatorname{Im} \left( \frac{x+i(y+1)}{2x+i(y+1)} \cdot \frac{2x-i(y+1)}{2x-i(y+1)} \right)$$

$$= \frac{2x(y+1) - x(y+1)}{4x^2 + (y+1)^2}$$

$$= \frac{x(y+1)}{4x^2 + (y+1)^2} = 0$$

$$\Rightarrow x = 0 \text{ or } y = -1$$

5. If  $\int \left( \left( \frac{x}{e} \right)^{2x} + \left( \frac{e}{x} \right)^{2x} \right) \ln x \, dx = \alpha \left( \frac{x}{e} \right)^{2x} + \beta \left( \frac{e}{x} \right)^{2x} + c$

where  $c$  is constant of integration, then

(1)  $\alpha + \beta = 0$       (2)  $\alpha + \beta = 1$

(3)  $\alpha\beta = \frac{1}{2}$       (4)  $\alpha\beta = \frac{1}{4}$

**Answer (1)**

**Sol.** Let  $\left( \frac{x}{e} \right)^{2x} = t$

$$2x(\ln x - 1) = \ln t$$

$$\left[ 2(\ln x - 1) + 2x \left( \frac{1}{x} \right) \right] dx = \frac{1}{t} dt$$

$$\ln x = \frac{1}{2t} dt$$

$$I = \int \left( t + \frac{1}{t} \right) \times \frac{1}{2t} dt = \frac{1}{2} \int \left( 1 + \frac{1}{t^2} \right) dt$$

$$= \frac{1}{2} \left( t - \frac{1}{t} \right) + c$$

$$= \frac{1}{2} \left( \left( \frac{x}{e} \right)^{2x} - \left( \frac{e}{x} \right)^{2x} \right) + c$$

$$\Rightarrow \alpha = \frac{1}{2}, \beta = -\frac{1}{2}$$

6. If a dice is thrown  $n$ -times and probability of getting 7 times odd is equal to 9 times even. Then

$$P(2 \text{ even}) \text{ is } \frac{K}{2^{15}} \text{ then } K \text{ is}$$

(1) 58      (2) 60  
 (3) 48      (4) 65

**Answer (2)**

**Sol.**  $P(\text{getting odd 7 times}) = P(\text{getting even 9 times})$

$${}^n C_7 \left( \frac{1}{2} \right)^7 \left( \frac{1}{2} \right)^{n-7} = {}^n C_9 \left( \frac{1}{2} \right)^9 \left( \frac{1}{2} \right)^{n-9}$$

$${}^n C_7 = {}^n C_9$$

$$n = 9 + 7 = 16$$

$$P(2 \text{ times even}) = {}^{16} C_2 \left( \frac{1}{2} \right)^{14} \left( \frac{1}{2} \right)^2$$

$$= \frac{{}^{16} C_2}{2^{16}} = \frac{16!}{2! \times 14!} \times \frac{1}{2^{16}}$$

$$= \frac{15 \times 16}{2 \times 2^{16}}$$

$$= \frac{15 \times 4}{2^{15}} = \frac{K}{2^{15}}$$

$$\Rightarrow K = 60$$

7. If  $\int_0^{t^2} (f(x) + x^2) dx = \frac{4}{3} t^3$ , then  $f(x)$  is

(1)  $x^2 - 2\sqrt{x}$       (2)  $x^2 + 2\sqrt{x}$   
 (3)  $x^2$       (4)  $-x^2 + 2\sqrt{x}$

**Answer (4)**

**Sol.**  $(f(t^2) + t^4) 2t = 4t^2$

$$f(t^2) + t^4 = 2t$$

$$f(x^2) = -x^4 + 2x$$

$$\text{Let } x^2 = u$$

$$f(u) = -u^2 + 2\sqrt{u}$$


$$f(x) = -x^2 + 2\sqrt{x}$$

8. The equation of conic is  $19x^2 + 15y^2 = 285$ . A concentric circle with radius 4 units is given then angle of common tangent made by minor axis of ellipse is

(1)  $\frac{\pi}{3}$       (2)  $\frac{\pi}{2}$   
 (3)  $\frac{\pi}{6}$       (4)  $\frac{\pi}{4}$

**Answer (1)**

**Sol.**





$$T_n = 4 + \frac{(n-1)}{2} [14 + (n-2)3]$$

$$= 4 + \frac{(n-1)}{2} [8 + 3n]$$

$$T_n = 4 + \frac{1}{2} (3n^2 + 5n - 8)$$

$$\sum T_n = S_n = \frac{3}{2} \sum n^2 + \frac{5}{2} \sum n$$

$$= \frac{3}{2} \frac{n(n+1)(2n+1)}{6} + \frac{5}{2} \frac{n(n+1)}{2}$$


$$\frac{S_{29} - S_9}{60} = 223$$

23. In  $\triangle ABC$ ,  $P$  is circumcentre,  $Q$  is orthocentre, then  $\overrightarrow{PA} + \overrightarrow{PB} + \overrightarrow{PC}$  is

- (1)  $2\overrightarrow{PQ}$       (2)  $\overrightarrow{PQ}$   
 (3)  $3\overrightarrow{PQ}$       (4)  $\frac{1}{2}\overrightarrow{PQ}$

**Answer (2)**

**Sol.**



Let  $P$  be origin then

$$\overrightarrow{PA} + \overrightarrow{PB} + \overrightarrow{PC} = \vec{a} + \vec{b} + \vec{c}$$

$$\overrightarrow{PD} = \frac{\vec{b} + \vec{c}}{2}$$

$$\overrightarrow{PQ} = 2\overrightarrow{PD}$$

$$= \vec{b} + \vec{c}$$

$$\overrightarrow{PQ} = \vec{a} + (\vec{b} + \vec{c})$$

$$= \vec{a} + \vec{b} + \vec{c}$$

24. Let  $S = \left\{ x \in \left[ -\frac{\pi}{2}, \frac{\pi}{2} \right] : 9^{1-\tan^2 x} + 9^{\tan^2 x} = 10 \right\}$  and

$$\beta = \sum_{x \in S} \left( \frac{x}{3} \right). \text{ Then } \frac{1}{7}(\beta - 14)^2 \text{ is}$$

**Answer (28)**

**Sol.** Let  $9^{\tan^2 x} = t$

$$\frac{9}{t} + t = 10$$

$$t^2 - 10t + 9 = 0$$

$$\Rightarrow t = 9 \text{ or } 1$$

$$9^{\tan^2 x} = 9$$

$$\Rightarrow \tan^2 x = 1$$

$$= \tan x = \pm 1$$

$$x = \pm \frac{\pi}{4}$$

or

$$\tan x = 0$$

$$\Rightarrow x = 0$$

$$\beta = \frac{0}{3} + \frac{\pi}{12} - \frac{\pi}{12} = 0$$

$$\therefore \frac{1}{7}(\beta - 14)^2 = \frac{14^2}{7} = 28$$

25. The coefficient of  $x$  and  $x^2$  in  $(1+x)^p (1-x)^q$  are 4 and -5, then  $2p+3q$  is

**Answer (63)**

$$\text{Sol. } (1+x)^p (1-x)^q = (1+px + \frac{p(p-1)}{2} x^2 + \dots)$$

$$(1-qx + q \frac{(q-1)}{2} x^2 + \dots)$$

$$\therefore \text{Coefficient of } x = p - q = 4 \quad \dots(i)$$

$$\text{Coefficient of } x^2 = \frac{q(q-1)}{2} - pq + \frac{p(p-1)}{2} = -5$$

$$\Rightarrow (p-q)^2 - (p+q) = -10$$

$$\therefore p+q = 26 \quad \dots(ii)$$

$$\text{By (i) and (ii) } p = 15, q = 11$$

$$\text{So, } 2p+3q = 30+33 = 63$$

26. Let  $\alpha$  be the remainder  $(22)^{2022} + (2022)^{22}$  is divided by 3 and  $\beta$  be the remainder when the same is divided by 7 then  $\alpha^2 + \beta^2$  is

**Answer (05)**

**Sol.**  $(22)^{2022} + (2022)^{22}$

For  $\alpha$

$$(21+1)^{2022} + \underbrace{(2022)^{22}}_{\text{divisible by 3}}$$

$$= (3k_1 + 1)$$

For  $\beta$

$$(21+1)^{2022} + (2023-1)^{22}$$

$$= (7\lambda + 1) + (7\mu + 1)$$

$$= 7k_2 + 2$$


$$\text{So, } \alpha = 1, \beta = 2$$

$$\alpha^2 + \beta^2 = 5$$

27. If area bounded by region  $\{(x, y) | |x^2 - 2| \leq y \leq x\}$  is A, then  $6A + 16\sqrt{2}$  is

**Answer (27)**

**Sol.**



$\therefore$  Required area

$$= \int_1^{\sqrt{2}} x - \{- (x^2 - 2)\} dx + \int_{\sqrt{2}}^2 \{x - (x^2 - 2)\} dx$$

$$= \int_1^{\sqrt{2}} (x^2 + x - 2) dx + \int_{\sqrt{2}}^2 (-x^2 + x + 2) dx$$

$$= \left( \frac{x^3}{3} + \frac{x^2}{2} - 2x \right)_1^{\sqrt{2}} + \left( \frac{-x^3}{3} + \frac{x^2}{2} + 2x \right)_{\sqrt{2}}^2$$

$$= \left( \frac{2\sqrt{2}}{3} + 1 - 2\sqrt{2} \right) - \left( \frac{1}{3} + \frac{1}{2} - 2 \right)$$

$$+ \left( \frac{-8}{3} + 2 + 4 \right) - \left( \frac{-2\sqrt{2}}{3} + 1 + 2\sqrt{2} \right)$$

$$\therefore 6A + 16\sqrt{2} = 27 - 16\sqrt{2} + 16\sqrt{2}$$

$$= 27$$

28.

29.

30.

□ □ □