



**NARAYANA GRABS
THE LION'S SHARE IN JEE-ADV.2022**

**5 RANKS in OPEN CATEGORY
ONLY FROM NARAYANA
IN TOP 10 AIR**




JEE MAIN (APRIL) 2023 (13-04-2023-FN)

Memory Based Question Paper
PHYSICS



PHYSICS

- 1.** Train A takes 35 sec less than train B. Find length of tunnel.




Ans. 1800

$$\text{Sol. } \frac{60\ell + 4\ell}{20} - \frac{61\ell}{30} = 35$$

$$\Rightarrow \ell = \frac{1050}{35}$$

$$\Rightarrow L = 60\ell = \frac{1050}{35} \times 60 = 1800 \text{ m}$$

- 2.** Find $P_1 - P_2$.



Ans. 175

$$\text{Sol. } A_1 V_1 = A_2 V_2$$

$$1.5 \times V_1 = 25 \times 10^{-2} \times 60$$

$$V_1 = \frac{25 \times 60 \times 10^{-2} \times 10}{1.5}$$

$$V_1 = 10 \text{ cm/s}$$

By Bernoulli's

$$P_1 + \frac{1}{2} \times 1000 \times (0.1)^2 = P_2 + \frac{1}{2} \times 1000 \times (0.6)^2$$

$$P_1 + 5 = P_2 + \frac{1}{2} \times 1000 \times 36 \times 10^{-2}$$

$$P_1 + 5 = P_2 + 180$$

$$P_1 - P_2 = 175 \text{ Pas.}$$

3. For a polytropic process $P = av^{-3}$. Find Bulk Modulus :

Ans. (3)

$$\text{Sol. } B = -\frac{dP}{dv/v}$$

$$Pv^3 = a$$

difference wrt to pressure





$$v^3 + P_3 v^2 \frac{dv}{dP} = 0$$

$$v = -3 \frac{Pdv}{dP} = 0$$

$$v = -3 \frac{P dv}{dP}$$

$$\frac{dP.v}{dv} = -3P$$


4. In an SHM draw the graph of T.E. – P.E. Vs x.







Ans. (3)

Sol. T.E. – P.E. = K.E.

$$\text{K.E.} = \frac{1}{2} m\omega^2 (A^2 - x^2)$$



5. Increasing order of power dissipation?



- (1) iv > i > ii > iii (2) iv > i > iii > ii (3) i > ii > iii > iv (4) iv > iii > ii > i

Ans. (1)

Sol. $P = i^2 R$

$$R_1 = \frac{3R}{2}, R_2 = \frac{2R}{3}, R_3 = \frac{R}{3}, R_4 = 3R : R_4 > R_1 > R_2 > R_3$$

6. Which of the following Maxwell equation is time dependant?

- (1) $\oint \vec{E} \cdot d\vec{s} = \frac{q_{in}}{\epsilon_0}$ (2) $\oint \vec{B} \cdot d\vec{s} = 0$ (3) $\oint \vec{E} \cdot d\vec{l} = -\frac{dQ_B}{dt}$ (4) $\oint \vec{B} \cdot d\vec{l} = \mu_0 I_{en}$

Ans. (4)

Sol. n^n : \rightarrow Ampere's Circuital law charges in time varying condition. Ans. (4)

7. If ratio of kinetic energy of two particle is $\frac{16}{9}$. If linear momentum of two particle are same then

ratio of mass $\frac{m_1}{m_2}$ is:

- (1) $\frac{9}{16}$ (2) $\frac{16}{9}$ (3) $\frac{4}{3}$ (4) $\frac{3}{4}$

Ans. (1)

Sol. $\frac{K_1}{K_2} = \frac{p_1^2}{2m_1} \times \frac{2m_2}{p_2^2} = \frac{m_2}{m_1} = \frac{16}{9}$

$$\frac{m_1}{m_2} = \frac{9}{16}$$

8. Mass of a planet is equal to 9 times of mass of earth and radius is 4 times the radius of earth. Find escape velocity (in km/sec.) of the planet. [Given; escape velocity of earth $V_E = 11.2$ km/sec]

Ans. 16.8

Sol. $V_P = \sqrt{\frac{2GM_P}{R_P}}$ $V_E = \sqrt{\frac{2GM_E}{R_E}}$

$$\frac{V_P}{V_E} = \frac{\sqrt{\frac{2GM_P}{R_P}}}{\sqrt{\frac{2GM_E}{R_E}}} = \sqrt{\frac{R_E}{R_P} \times \frac{M_P}{M_E}}$$

$$V_P = \sqrt{\frac{1}{4} \times 9} \times V_E = \frac{3}{2} V_E$$

$$V_P = \frac{3}{2} \times 11.2 \text{ km/sec.}$$

$$= 16.8 \text{ km/sec}$$

9. Find the value of x , if elastic potential energy per unit volume is $x \times 10^9$ J stored in the wire of length $L = 50$ mm. Young's modulus $Y = 2 \times 10^{11}$ N/m² and change in length ΔL is the wire is 10 mm.

Ans. 4×10^9 J/m³

Sol. $\frac{\text{Energy}}{\text{Volume}} = \frac{1}{2} \times \text{stress} \times \text{strain}$

$$= \frac{1}{2} \times Y \times (\text{strain})^2$$

$$= \frac{1}{2} \times 2 \times 10^{11} \times \left[\frac{10 \times 10^{-3}}{50 \times 10^{-3}} \right]^2$$

$$= 10^{11} \times \left[\frac{1}{5} \right]^2 = 4 \times 10^9 \text{ J/m}^3$$

10. Find the ratio of radius of 2nd orbit of He⁺ and 4th orbit of Be³⁺

(1) $\frac{1}{2}$

(2) $\frac{2}{1}$

(3) $\frac{4}{1}$

(4) $\frac{1}{4}$

Ans. (1)

Sol. $r \propto \frac{n^2}{z}$

$$\frac{r_{He^+}}{r_{Be^{3+}}} = \frac{2^2 \times 4}{2 \times 4 \times 4} = \frac{1}{2}$$

11. If the height of the tower used for L.D.S. is increased by 21% then percentage change in range is :

(1) 10%

(2) 21%

(3) 19%


(4) 42%

Ans. (1)

Sol. $R_l = \sqrt{2H Re}$... (1)

$$R_2 = \sqrt{2\left(\frac{Hl21}{100}\right)Re} \quad \dots (2)$$

$$\% \text{ change} = \frac{(R_2 - R_1)}{R_1} \times 100 = \frac{\sqrt{\frac{121}{100}} - 1}{1} \times 100 = \frac{\frac{11}{10} - 1}{1} \times 100 = 10\%$$



Given $R_B = 10 \text{ k}\Omega$, $R_C = 1 \text{ k}\Omega$, power gain is 10^x . Find x

(1) 1

(2) 2

(3) 0

(4) 3

Ans. (1)

Sol. Power gain = $A_v \cdot A_I = B \frac{R_C}{R_B} \cdot B = B^2 \frac{R_C}{R_B} = \left(\frac{(20-10) \times 10^3}{(200-100) \times 10^{-6}} \right)^2 \times \frac{1 \times 10^3}{10 \times 10^3} = 10$

Hence x = 1

13. Mass (m) = (5 ± 0.5) kg, speed (v) = (20 ± 0.4) m/s. Find the kinetic energy.

- (1) $(1000 + 70)$ J (2) (1000 ± 140) J
 (3) (500 ± 140) J (4) (500 ± 70) J

Ans. (2)


Sol. $k = mv^2$

$$k = \frac{1}{2} \times 5 \times 400 = 5 \times 200 = 1000 \text{ J}$$

$$\frac{\Delta k}{2k} = \frac{\Delta m}{m} + \frac{2\Delta v}{v} = \frac{0.5}{5} + \frac{2 \times 0.4}{20}$$

$$\Delta k = 1000 \left(\frac{1}{10} + \frac{4}{100} \right) = 1000 \left(\frac{10+4}{100} \right) = 140 \text{ J}$$


14. Radius of the cylinder is R find displacement of point B in half rotation. [Cylinder performs pure rolling]



- (1) $2R$ (2) $R\sqrt{4+\pi^2}$ (3) R (4) $R\sqrt{1+\pi^2}$


Ans. (2)

Sol.



$$\text{Displacement} = \sqrt{(2R)^2 + (\pi R)^2} = R\sqrt{4+\pi^2}$$


15. Find the apparent depth of the bottom surface of the tank, when seen from above (in air)?



- (1) $\frac{d}{2} \left[\frac{n_1 + n_2}{n_1 n_2} \right]$ (2) $\frac{d}{2} \left[\frac{n_2 - n_1}{n_2 n_1} \right]$ (3) $d \left[\frac{n_1 + n_2}{n_1 n_2} \right]$ (4) $d \left[\frac{n_2 - n_1}{n_2 n_1} \right]$

Ans. (1)


Sol.



Formula used : $d_{app} = \frac{d_1}{n_1} + \frac{d_2}{n_2}$

$$d_{app} = \frac{d}{2} \left[\frac{n_1 + n_2}{n_1 n_2} \right]$$

- 16.** Find out apparent speed of bird as seen by fish :



Ans. (2)

$$\text{Sol. } \mathbf{v} = \mathbf{v}_{\text{fish}} + \frac{\mathbf{v}_{\text{Bird}} \times 4 / 3}{1}$$

$$= 9 + 12 \times \frac{4}{3}$$

$$= 9 + 16 = 25 \text{ m/s}$$

17. If a wire of resistance R is connected across V_0 , then power is P . The wire is cut into two equal parts 2^{nd} connected with V_0 individually then sum of power P_2 . Find out $\frac{P}{P_2}$ is $\frac{1}{x}$ find out x ?

Ans. 4

$$\text{Sol. } P = \frac{V_0^2}{R}$$

$$P_2 = \frac{V_0^2}{R/2} + \frac{V_0^2}{R/2} = \frac{4V_0^2}{R} = 4P$$

$$\frac{P}{P_2} = \frac{1}{4}$$

- 18.** A particle is performing SHM having position $x = A \cos 30^\circ$, and $A = 40$ cm. If its kinetic energy at this position is 200 J, the value of force constant (in kilo-N/m) is

Ans. 10

Sol. $\frac{1}{2}k(A^2 - x^2) = 200$ $[x = \frac{\sqrt{3}A}{2}]$

$$\frac{1}{2}k\left(A^2 - \frac{3A^2}{4}\right) = 200 \quad [\omega = \sqrt{\frac{k}{m}}]$$

$$\frac{1}{2}k \frac{A^2}{4} = 200$$

$$k = \frac{200 \times 2 \times 4 \times 100 \times 100}{40 \times 40} = 10^4$$

$$= 10 \times 10^3$$

$$= 10 \text{ k N/m}$$

- 19.** For a ideal gas relation between its average speed (V_{avg}) and r.m.s. speed (V_{rms}) is

(Use : $\pi = \frac{22}{7}$)

$$V_{rms} = \left(1 + \frac{5}{x}\right)^{\frac{1}{2}} V_{avg}$$

Then value of 'x' is :

Ans. 28

Sol. $\sqrt{\frac{3RT}{M}} = \left(1 + \frac{5}{x}\right)^{\frac{1}{2}} \sqrt{\frac{8RT}{\pi M}}$ $\Rightarrow \frac{3 \times 22}{7 \times 8} = 1 + \frac{5}{x} \Rightarrow x = 28$

- 20.** An electric dipole is placed in an external electric field 4×10^{-4} N/C at angle 30° . Magnitude of charge of dipole is 10^{-2} C and separation between them is 0.2 mm. Find torque acting on dipole.

(1) 6×10^{-10} N-m (2) 14×10^{-8} N-m (3) 4×10^{-10} N-m (4) 8×10^{-10} N-m

Ans. (3)

Sol. $\vec{\tau} = \vec{P} \times \vec{\epsilon}$

$$P = qd = 10^{-2} \times 0.2 \times 10^{-3}$$

$$\tau = P\epsilon \sin 30$$

$$= 2 \times 10^{-6} \times 4 \times 10^{-4} \times \frac{1}{2}$$

$$= 4 \times 10^{-10} \text{ N-m}$$

- 21.** A solid sphere is Rolling on a flat horizontal surface. If the ratio of angular momentum to total kinetic energy is $\frac{\pi}{22}$, then find the angular speed (in rad/sec) with which sphere is moving?

Ans. 14

$$\text{Sol. } \frac{\text{Angular momentum}}{\text{Total kinetic energy}} = \frac{\left(\frac{2}{5}mR^2 + mR^2\right)\omega}{\frac{1}{2}mv^2 + \frac{1}{2}I\omega^2} = \frac{\pi}{22}$$

(Taking $v = \omega R$)

$$\frac{\frac{7}{5}mR^2 \cdot \omega}{\frac{7}{10}mv^2} = \frac{\pi}{22} \quad \Rightarrow \quad \omega = 14 \text{ rad/sec}$$

- 22.** Match the following lists.

- | | |
|---|------------|
| (A) Troposphere | (P) 300 km |
| (B) E part of stratosphere | (Q) 80 km |
| (C) F ₂ part of thermosphere | (R) 20 km |
| (D) D-part of stratosphere | (S) 100 km |
- (1) (A) \rightarrow R; (B) \rightarrow S; (C) \rightarrow P; (D) \rightarrow Q
 (2) (A) \rightarrow S; (B) \rightarrow R; (C) \rightarrow Q; (D) \rightarrow P
 (3) (A) \rightarrow Q; (B) \rightarrow S; (C) \rightarrow P; (D) \rightarrow R
 (4) (A) \rightarrow R; (B) \rightarrow P; (C) \rightarrow Q; (D) \rightarrow S

Ans. (1)

- 23.** Two metals A and B having work function $\phi_A = 9 \text{ eV}$ and $\phi_B = 4.5 \text{ eV}$. Find difference of threshold wavelength.

- (1) 1378 Å (2) 2100 Å (3) 1500 Å (4) 1100 Å

Ans. (1)


$$\text{Sol. } \lambda_A = \left(\frac{12400}{9} \right) \text{\AA} = 1377.77 \text{\AA}$$

$$\lambda_B = \left(\frac{12400}{4.5} \right) \text{\AA} = 2755.55 \text{\AA}$$

$$\lambda_B - \lambda_A = 1377.78 \text{\AA}$$

24. A bullet of mass 10 gm is fired with muzzle speed 600 m/s from 3 kg gun of barrel length 30 cm.

Find impulse on gun :



- (1) 60 Ns (2) 3 Ns (3) 8 Ns (4) 6 Ns

Ans. (4)

Sol. By momentum conservation

$$0 = 3(-v) + 0.01(600 - v)$$

$$v \approx 2 \text{ m/s}$$

$$\text{Impulse on gun} = 3 \times 2 = 6 \text{ Ns}$$

25. For the given radioactive decay



The Q-value is equal to:

- | | |
|------------------------|----------------------|
| (1) $294b + 4c - 298a$ | (2) $92b + 2c - 94a$ |
| (3) $294b + 4c + 298a$ | (4) $92b + 2c + 94a$ |


Ans. (1)

Sol. $Q = u_i - u_f$

$$Q = (B.E)_f - (B.E)_i$$

$$= 294b + 4c - 298a$$

25. For the given radioactive decay



The Q-value is equal to:

- | |
|------------------------|
| (1) $294b + 4c - 298a$ |
| (2) $92b + 2c - 94a$ |
| (3) $294b + 4c + 298a$ |
| (4) $92b + 2c + 94a$ |








Ans. (1)

Sol. $Q = u_i - u_f$

$$Q = (B.E)_f - (B.E)_i$$

$$= 294b + 4c - 298a$$

26. Which of the following represents wave form of output.



Ans. (3)

Sol. $Y = \overline{A\bar{B}} = \overline{A} + B$

